

Matrices

G V V Sharma*

CONTENTS

1	Definitions	1
1.1	Eigenvalues and Eigenvectors	1
1.2	Determinants	1
2	Rank of a Matrix	1
3	Inverse of a Matrix	2
4	Cayley-Hamilton Theorem	2
5	Matrix Polynomials	3
6	Linear Equations and Rank	3
7	Matrix Inverse	4

Abstract—This manual provides a simple introduction to matrices and their properties based on the NCERT textbooks from Class 6-12.

1 DEFINITIONS

1.1 Eigenvalues and Eigenvectors

1.1.1. The eigenvalue λ and the eigenvector \mathbf{x} for a matrix \mathbf{A} are defined as,

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x} \quad (1.1.1.1)$$

1.1.2. The eigenvalues are calculated by solving the equation

$$f(\lambda) = |\lambda\mathbf{I} - \mathbf{A}| = 0 \quad (1.1.2.1)$$

The above equation is known as the characteristic equation.

1.1.3. According to the Cayley-Hamilton theorem,

$$f(\lambda) = 0 \implies f(\mathbf{A}) = 0 \quad (1.1.3.1)$$

1.1.4. The trace of a square matrix is defined to be the sum of the diagonal elements.

$$\text{tr}(\mathbf{A}) = \sum_{i=1}^N a_{ii}. \quad (1.1.4.1)$$

where a_{ii} is the i th diagonal element of the matrix \mathbf{A} .

1.1.5. The trace of a matrix is equal to the sum of the eigenvalues

$$\text{tr}(\mathbf{A}) = \sum_{i=1}^N \lambda_i \quad (1.1.5.1)$$

1.2 Determinants

1.2.1. Let

$$\mathbf{A} = \begin{pmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{pmatrix}. \quad (1.2.1.1)$$

be a 3×3 matrix. Then,

$$|\mathbf{A}| = a_1 \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} - a_2 \begin{vmatrix} b_1 & c_1 \\ b_3 & c_3 \end{vmatrix} + a_3 \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}. \quad (1.2.1.2)$$

1.2.2. Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be the eigenvalues of a matrix \mathbf{A} . Then, the product of the eigenvalues is equal to the determinant of \mathbf{A} .

$$|\mathbf{A}| = \prod_{i=1}^n \lambda_i \quad (1.2.2.1)$$

1.2.3.

$$|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}| \quad (1.2.3.1)$$

1.2.4. If \mathbf{A} be an $n \times n$ matrix,

$$|k\mathbf{A}| = k^n |\mathbf{A}| \quad (1.2.4.1)$$

2 RANK OF A MATRIX

2.1. The rank of a matrix is defined as the number of nonzero rows obtained after row reduction.

2.2. An $n \times n$ matrix is invertible if and only if its rank is n .

*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

3 INVERSE OF A MATRIX

3.1. For a 2×2 matrix

$$\mathbf{A} = \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}, \quad (3.1.1)$$

the inverse is given by

$$\mathbf{A}^{-1} = \frac{1}{|\mathbf{A}|} \begin{pmatrix} b_2 & -b_1 \\ -a_2 & a_1 \end{pmatrix}, \quad (3.1.2)$$

3.2. For higher order matrices, the inverse should be calculated using row operations.

4 CAYLEY-HAMILTON THEOREM

4.1. If

$$\mathbf{A} = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{pmatrix}, \quad (4.1.1)$$

prove that

$$\mathbf{A}^3 - 6\mathbf{A}^2 + 7\mathbf{A} + 2\mathbf{I} = 0 \quad (4.1.2)$$

Solution: From (1.1.2.1), the characteristic equation is

$$\begin{vmatrix} 1-\lambda & 0 & 2 \\ 0 & 2-\lambda & 1 \\ 2 & 0 & 3-\lambda \end{vmatrix} = 0 \quad (4.1.3)$$

which can be expanded to obtain

$$(1-\lambda)(2-\lambda)(3-\lambda) + 2(-2(2-\lambda)) = 0 \quad (4.1.4)$$

yielding

$$\lambda^3 - 6\lambda^2 + 7\lambda + 2 = 0 \quad (4.1.5)$$

upon simplification. Using the Cayley-Hamilton theorem in (1.1.3.1), (4.1.2) is obtained

4.2. If

$$\mathbf{A} = \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix}, \quad (4.2.1)$$

show that

$$\mathbf{A}^2 - 5\mathbf{A} + 7\mathbf{I} = 0 \quad (4.2.2)$$

Solution: The characteristic equation is

$$|\mathbf{A} - \lambda\mathbf{I}| = 0 \quad (4.2.3)$$

$$\Rightarrow \begin{vmatrix} 3-\lambda & 1 \\ -1 & 2-\lambda \end{vmatrix} = 0 \quad (4.2.4)$$

$$\Rightarrow (3-\lambda)(2-\lambda) + 1 = 0 \quad (4.2.5)$$

$$\text{or, } \lambda^2 - 5\lambda + 7 = 0 \quad (4.2.6)$$

Using the Cayley-Hamilton theorem, (4.2.2) is obtained.

4.3. If \mathbf{A} is square matrix such that

$$\mathbf{A}^2 = \mathbf{A}, \quad (4.3.1)$$

then

$$g(\mathbf{A}) = (\mathbf{I} + \mathbf{A})^3 - 7\mathbf{A} \quad (4.3.2)$$

is equal to

- a) \mathbf{A}
- b) $\mathbf{I} - \mathbf{A}$
- c) \mathbf{I}
- d) $3\mathbf{A}$

Solution: From (4.3.2),

$$g(\lambda) = (1 + \lambda)^3 - 7\lambda \quad (4.3.3)$$

Also,

$$\mathbf{A}^2 = \mathbf{A} \implies \mathbf{A}^2 - \mathbf{A} = 0 \quad (4.3.4)$$

Using the Cayley-Hamilton theorem, the eigenvalues satisfy the characteristic equation

$$f(\lambda) = \lambda^2 - \lambda = 0 \quad (4.3.5)$$

$\therefore f(\lambda)$ is of degree 2 and $g(\lambda)$ is of degree 3, (4.3.3) can be expressed as

$$g(\lambda) = f(\lambda)q(\lambda) + a\lambda + b \quad (4.3.6)$$

where a, b are real numbers and $q(\lambda)$ is some polynomial. Thus,

$$g(0) = b = 1 \quad (4.3.7)$$

$$g(1) = a + b = 1 \quad (4.3.8)$$

$$\implies a = 0, b = 1 \quad (4.3.9)$$

Thus,

$$g(\mathbf{A}) = f(\mathbf{A})q(\mathbf{A}) + a\mathbf{A} + b\mathbf{I} \quad (4.3.10)$$

$$= \mathbf{I} \quad (4.3.11)$$

upon substituting from (4.3.5) and (4.3.9). Option c is the valid answer.

4.4. If

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 3 & -2 & 1 \\ 4 & 2 & 1 \end{pmatrix} \quad (4.4.1)$$

then show that

$$\mathbf{A}^3 - 23\mathbf{A} - 40\mathbf{I} = 0 \quad (4.4.2)$$

Solution: The Characteristic equation is given by

$$|\mathbf{A} - \lambda\mathbf{I}| = 0 \quad (4.4.3)$$

$$\Rightarrow \begin{vmatrix} 1-\lambda & 2 & 3 \\ 3 & -2-\lambda & 1 \\ 4 & 2 & 1-\lambda \end{vmatrix} = 0 \quad (4.4.4)$$

which can be expressed as

$$\begin{aligned} &\Rightarrow (1-\lambda)((-2-\lambda)(1-\lambda)-2) \\ &-2(3(1-\lambda)-4)+3(6+4(2+\lambda))=0 \end{aligned} \quad (4.4.5)$$

and simplified to obtain

$$\Rightarrow \lambda^3 - 23\lambda - 40 = 0. \quad (4.4.6)$$

Using the Cayley-Hamilton Theorem, (4.4.2) is obtained.

5 MATRIX POLYNOMIALS

5.1. Find $\mathbf{A}^2 - 5\mathbf{A} + 6\mathbf{I}$, if $\mathbf{A} = \begin{pmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{pmatrix}$

Solution: Factorizing the corresponding polynomial,

$$f(x) = x^2 - 5x + 6 \quad (5.1.1)$$

$$= (x-3)(x-2) \quad (5.1.2)$$

$$\Rightarrow \mathbf{A}^2 - 5\mathbf{A} + 6\mathbf{I} = (\mathbf{A} - 3\mathbf{I})(\mathbf{A} - 5\mathbf{I}) \quad (5.1.3)$$

$$\mathbf{A} - 3\mathbf{I} = \begin{pmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{pmatrix} + \begin{pmatrix} -3 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & -3 \end{pmatrix} \quad (5.1.4)$$

$$= \begin{pmatrix} -1 & 0 & 1 \\ 2 & -2 & 3 \\ 1 & -1 & -3 \end{pmatrix} \quad (5.1.5)$$

$$\mathbf{A} - 5\mathbf{I} = \begin{pmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{pmatrix} + \begin{pmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{pmatrix} \quad (5.1.6)$$

$$= \begin{pmatrix} 0 & 0 & 1 \\ 2 & -1 & 3 \\ 1 & -1 & -2 \end{pmatrix} \quad (5.1.7)$$

Multiplying the above expressions,

$$\begin{aligned} \mathbf{A}^2 - 5\mathbf{A} + 6\mathbf{I} &= \\ &\begin{pmatrix} -1 & 0 & 1 \\ 2 & -2 & 3 \\ 1 & -1 & -3 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \\ 2 & -1 & 3 \\ 1 & -1 & -2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & -1 & -3 \\ -1 & -1 & -10 \\ -5 & 4 & 4 \end{pmatrix} \end{aligned} \quad (5.1.8)$$

6 LINEAR EQUATIONS AND RANK

6.1. Two rails are represented by the equations

$$\begin{aligned} \begin{pmatrix} 1 & 2 \end{pmatrix} \mathbf{x} &= 4 \text{ and} \\ \begin{pmatrix} 2 & 4 \end{pmatrix} \mathbf{x} &= 12. \end{aligned} \quad (6.1.1)$$

Will the rails cross each other?

Solution: The above equations can be expressed as the matrix equation

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 4 \\ 12 \end{pmatrix} \quad (6.1.2)$$

The augmented matrix for the above equation is row reduced as follows

$$\begin{pmatrix} 1 & 2 & 4 \\ 2 & 4 & 12 \end{pmatrix} \xrightarrow{R_2 \leftarrow \frac{R_2}{2}} \begin{pmatrix} 1 & 2 & 4 \\ 1 & 2 & 6 \end{pmatrix} \quad (6.1.3)$$

$$\xrightarrow{R_2 \leftarrow R_2 - R_1} \begin{pmatrix} 1 & 2 & 4 \\ 0 & 0 & 2 \end{pmatrix} \quad (6.1.4)$$

\therefore row reduction of the 2×3 matrix

$$\begin{pmatrix} 1 & 2 & 4 \\ 2 & 4 & 12 \end{pmatrix} \quad (6.1.5)$$

results in a matrix with 2 nonzero rows, its rank is 2. Similarly, the rank of the matrix

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \quad (6.1.6)$$

is 1, from 6.1.4.

$$\therefore \text{rank} \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \neq \text{rank} \begin{pmatrix} 1 & 2 & 4 \\ 2 & 4 & 12 \end{pmatrix}, \quad (6.1.7)$$

(6.1.1) has no solution.

6.2. Check whether the pair of equations

$$\begin{aligned} \begin{pmatrix} 1 & 3 \end{pmatrix} \mathbf{x} &= 6 \text{ and} \\ \begin{pmatrix} 2 & -3 \end{pmatrix} \mathbf{x} &= 12 \end{aligned} \quad (6.2.1)$$

is consistent.

Solution: The above equations can be expressed as the matrix equation

$$\begin{pmatrix} 1 & 3 \\ 2 & -3 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 6 \\ 12 \end{pmatrix} \quad (6.2.2)$$

The augmented matrix for the above equation is row reduced as follows

$$\begin{pmatrix} 1 & 3 & 6 \\ 2 & -3 & 12 \end{pmatrix} \xrightarrow{R_2 \leftarrow R_2 - 2R_1} \quad (6.2.3)$$

$$\begin{pmatrix} 1 & 3 & 6 \\ 0 & 1 & 0 \end{pmatrix} \xrightarrow{R_1 \leftarrow R_1 - 3R_2} \begin{pmatrix} 1 & 0 & 6 \\ 0 & 1 & 0 \end{pmatrix} \quad (6.2.4)$$

$$\Rightarrow \mathbf{x} = \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (6.2.5)$$

which is the solution of 6.1.1.

6.3. Find whether the following pair of equations has no solution, unique solution or infinitely many solutions:

$$\begin{aligned} (5 \quad -8) \mathbf{x} &= -1 \text{ and} \\ (3 \quad -\frac{24}{5}) \mathbf{x} &= -\frac{3}{5} \end{aligned} \quad (6.3.1)$$

Solution: The above equations can be expressed as the matrix equation

$$\begin{pmatrix} 5 & -8 \\ 3 & -\frac{24}{5} \end{pmatrix} \mathbf{x} = -\begin{pmatrix} 1 \\ \frac{3}{5} \end{pmatrix} \quad (6.3.2)$$

The augmented matrix for the above equation is row reduced as follows

$$\begin{pmatrix} 5 & -8 & -1 \\ 3 & -\frac{24}{5} & -\frac{3}{5} \end{pmatrix} \xrightarrow{R_2 \leftarrow 5R_2} \begin{pmatrix} 5 & -8 & 1 \\ 15 & -24 & -3 \end{pmatrix} \quad (6.3.3)$$

$$\xrightarrow{R_2 \leftarrow R_2 - 3R_1} \begin{pmatrix} 5 & -8 & 1 \\ 0 & 0 & 0 \end{pmatrix} \quad (6.3.4)$$

$$\therefore \text{rank} \begin{pmatrix} 5 & -8 \\ 3 & -\frac{24}{5} \end{pmatrix} = \text{rank} \begin{pmatrix} 5 & -8 & 1 \\ 3 & -\frac{24}{5} & -\frac{3}{5} \end{pmatrix} \quad (6.3.5)$$

$$= 1 < \dim \begin{pmatrix} 5 & -8 \\ 3 & -\frac{24}{5} \end{pmatrix} = 2, \quad (6.3.6)$$

(6.3.1) has infinitely many solutions.

7 MATRIX INVERSE

7.1. Using elementary transformations, find the inverse of $\begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix}$

Solution: Given that

$$\mathbf{A} = \begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix} \quad (7.1.1)$$

The augmented matrix $[\mathbf{A}|\mathbf{I}]$ is as given below:-

$$\left(\begin{array}{cc|cc} 1 & -1 & 1 & 0 \\ 2 & 3 & 0 & 1 \end{array} \right) \quad (7.1.2)$$

We apply the elementary row operations on $[\mathbf{A}|\mathbf{I}]$ as follows :-

$$[\mathbf{A}|\mathbf{I}] = \left(\begin{array}{cc|cc} 1 & -1 & 1 & 0 \\ 2 & 3 & 0 & 1 \end{array} \right) \quad (7.1.3)$$

$$\xrightarrow{R_2 \leftarrow R_2 - 2R_1} \left(\begin{array}{cc|cc} 1 & -1 & 1 & 0 \\ 0 & 5 & -2 & 1 \end{array} \right) \quad (7.1.4)$$

$$\xrightarrow{R_2 \leftarrow \frac{R_2}{5}} \left(\begin{array}{cc|cc} 1 & -1 & 1 & 0 \\ 0 & 1 & -\frac{2}{5} & \frac{1}{5} \end{array} \right) \quad (7.1.5)$$

$$\xrightarrow{R_2 \leftarrow R_1 + R_2} \left(\begin{array}{cc|cc} 1 & 0 & \frac{3}{5} & \frac{1}{5} \\ 0 & 1 & -\frac{2}{5} & \frac{1}{5} \end{array} \right) \quad (7.1.6)$$

By performing elementary transformations on augmented matrix $[\mathbf{A}|\mathbf{I}]$, we obtained the augmented matrix in the form $[\mathbf{I}|\mathbf{A}]$. Hence we can conclude that the matrix \mathbf{A} is invertible and

$$\mathbf{A}^{-1} = \begin{pmatrix} \frac{3}{5} & \frac{1}{5} \\ -\frac{2}{5} & \frac{1}{5} \end{pmatrix} \quad (7.1.7)$$

7.2. Using elementary transformations, find the inverse of $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$

Solution: Given that

$$\mathbf{A} = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \quad (7.2.1)$$

The augmented matrix $[\mathbf{A}|\mathbf{I}]$ is as given below:-

$$\left(\begin{array}{cc|cc} 2 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{array} \right) \quad (7.2.2)$$

We apply the elementary row operations on $[\mathbf{A}|\mathbf{I}]$ as follows :-

$$[\mathbf{A}|\mathbf{I}] = \left(\begin{array}{cc|cc} 2 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{array} \right) \quad (7.2.3)$$

$$\xrightarrow{R_1 \leftarrow R_1 - R_2} \left(\begin{array}{cc|cc} 1 & 0 & 1 & -1 \\ 1 & 1 & 0 & 1 \end{array} \right) \quad (7.2.4)$$

$$\xrightarrow{R_2 \leftarrow R_2 - R_1} \left(\begin{array}{cc|cc} 1 & 0 & 1 & -1 \\ 0 & 1 & -1 & 2 \end{array} \right) \quad (7.2.5)$$

By performing elementary transformations on augmented matrix $[\mathbf{A}|\mathbf{I}]$, we obtained the augmented matrix in the form $[\mathbf{I}|\mathbf{A}]$. Hence we can conclude that the matrix \mathbf{A} is invertible and inverse of the matrix is

$$\mathbf{A}^{-1} = \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix} \quad (7.2.6)$$

7.3. Obtain the inverse of the following matrix using elementary operations

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{pmatrix}.$$

Solution: Given that

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{pmatrix}, \quad (7.3.1)$$

The augmented matrix $[\mathbf{A}|\mathbf{I}]$ is

$$\left(\begin{array}{ccc|ccc} 0 & 1 & 2 & 1 & 0 & 0 \\ 1 & 2 & 3 & 0 & 1 & 0 \\ 3 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \quad (7.3.2)$$

Applying elementary row operations on $[\mathbf{A}|\mathbf{I}]$,

$$[\mathbf{A}|\mathbf{I}] = \left(\begin{array}{ccc|ccc} 0 & 1 & 2 & 1 & 0 & 0 \\ 1 & 2 & 3 & 0 & 1 & 0 \\ 3 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \quad (7.3.3)$$

$$\xleftrightarrow{R_1 \leftrightarrow R_2} \left(\begin{array}{ccc|ccc} 1 & 2 & 3 & 0 & 1 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 \\ 3 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \quad (7.3.4)$$

$$\xleftrightarrow{R_3 \leftarrow R_3 - 3R_1} \left(\begin{array}{ccc|ccc} 1 & 2 & 3 & 0 & 1 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 \\ 0 & -5 & -8 & 0 & -3 & 1 \end{array} \right) \quad (7.3.5)$$

$$\xleftrightarrow{R_1 \leftarrow R_1 - 2R_2} \left(\begin{array}{ccc|ccc} 1 & 0 & -1 & -2 & 1 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 \\ 0 & -5 & -8 & 0 & -3 & 1 \end{array} \right) \quad (7.3.6)$$

$$\xleftrightarrow{R_3 \leftarrow R_3 + 5R_2} \left(\begin{array}{ccc|ccc} 1 & 0 & -1 & -2 & 1 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 \\ 0 & 0 & 2 & 5 & -3 & 1 \end{array} \right) \quad (7.3.7)$$

$$\xleftrightarrow{R_3 \leftarrow R_3/2} \left(\begin{array}{ccc|ccc} 1 & 0 & -1 & -2 & 1 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & \frac{5}{2} & -\frac{3}{2} & \frac{1}{2} \end{array} \right) \quad (7.3.8)$$

$$\xleftrightarrow{R_1 \leftarrow R_1 + R_3} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & \frac{5}{2} & -\frac{3}{2} & \frac{1}{2} \end{array} \right) \quad (7.3.9)$$

$$\xleftrightarrow{R_2 \leftarrow R_2 - 2R_3} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 0 & -4 & 3 & -1 \\ 0 & 0 & 1 & \frac{5}{2} & -\frac{3}{2} & \frac{1}{2} \end{array} \right) \quad (7.3.10)$$

By performing elementary transformations on augmented matrix $[\mathbf{A}|\mathbf{I}]$, we obtained the augmented matrix in the form $[\mathbf{I}|\mathbf{A}]$. Hence we can conclude that the matrix \mathbf{A} is invertible and inverse of the matrix is

$$\mathbf{A}^{-1} = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ -4 & 3 & -1 \\ \frac{5}{2} & -\frac{3}{2} & \frac{1}{2} \end{pmatrix} \quad (7.3.11)$$

7.4. Find \mathbf{P}^{-1} , if it exists, given

$$\mathbf{P} = \begin{pmatrix} 10 & -2 \\ -5 & 1 \end{pmatrix}.$$

Solution: Using row reduction,

$$\begin{pmatrix} 10 & -2 \\ -5 & 1 \end{pmatrix} \xleftrightarrow{R_2 \leftarrow R_2 + \frac{R_1}{2}} \begin{pmatrix} 10 & -2 \\ 0 & 0 \end{pmatrix} \quad (7.4.1)$$

Since we obtain a zero row, \mathbf{P}^{-1} does not exist.