

Points and Vectors

G V V Sharma*

CONTENTS

1	Definitions	1
1.1	2×1 vectors	1
1.2	3×1 vectors	1
2	Area of a Triangle	2
3	Angle Between Vectors	3

Abstract—This manual provides an introduction to vectors and their properties, based on the NCERT textbooks from Class 6-12.

1 DEFINITIONS

1.1 2×1 vectors

1.1.1. Let

$$\mathbf{A} \equiv \vec{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \quad (1.1.1.1)$$

$$\equiv a_1 \vec{i} + a_2 \vec{j}, \quad (1.1.1.2)$$

$$\mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}, \quad (1.1.1.3)$$

be 2×1 vectors. Then, the determinant of the 2×2 matrix

$$\mathbf{M} = \begin{pmatrix} \mathbf{A} & \mathbf{B} \end{pmatrix} \quad (1.1.1.4)$$

is defined as

$$|\mathbf{M}| = \begin{vmatrix} \mathbf{A} & \mathbf{B} \end{vmatrix} \quad (1.1.1.5)$$

$$= \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1 b_2 - a_2 b_1 \quad (1.1.1.6)$$

1.1.2. The area of the triangle with vertices $\mathbf{A}, \mathbf{B}, \mathbf{C}$ is given by the absolute value of

$$\frac{1}{2} |\mathbf{A} - \mathbf{B} \quad \mathbf{A} - \mathbf{C}| \quad (1.1.2.1)$$

1.1.3. The transpose of \mathbf{A} is defined as

$$\mathbf{A}^\top = \begin{pmatrix} a_1 & a_2 \end{pmatrix} \quad (1.1.3.1)$$

1.1.4. The *inner product* or *dot product* is defined as

$$\mathbf{A}^\top \mathbf{B} \equiv \mathbf{A} \cdot \mathbf{B} \quad (1.1.4.1)$$

$$= \begin{pmatrix} a_1 & a_2 \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = a_1 b_1 + a_2 b_2 \quad (1.1.4.2)$$

1.1.5. *norm* of \mathbf{A} is defined as

$$\|\mathbf{A}\| \equiv |\vec{A}| \quad (1.1.5.1)$$

$$= \sqrt{\mathbf{A}^\top \mathbf{A}} = \sqrt{a_1^2 + a_2^2} \quad (1.1.5.2)$$

Thus,

$$\|\lambda \mathbf{A}\| \equiv |\lambda \vec{A}| \quad (1.1.5.3)$$

$$= |\lambda| \|\mathbf{A}\| \quad (1.1.5.4)$$

1.1.6. The angle between two vectors is given by

$$\theta = \cos^{-1} \frac{\mathbf{A}^\top \mathbf{B}}{\|\mathbf{A}\| \|\mathbf{B}\|} \quad (1.1.6.1)$$

1.1.7. The *direction vector* of the line joining two points \mathbf{A}, \mathbf{B} is given by

$$\mathbf{m} = \mathbf{A} - \mathbf{B} \quad (1.1.7.1)$$

1.1.8. The *normal vector* to \mathbf{m} is defined by

$$\mathbf{m}^\top \mathbf{n} = 0 \quad (1.1.8.1)$$

1.2 3×1 vectors

1.2.1. Let

$$\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \equiv a_1 \vec{i} + a_2 \vec{j} + a_3 \vec{k}, \quad (1.2.1.1)$$

$$\mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}, \quad (1.2.1.2)$$

and

$$\mathbf{A}_{ij} = \begin{pmatrix} a_i \\ a_j \end{pmatrix}, \mathbf{B}_{ij} = \begin{pmatrix} b_i \\ b_j \end{pmatrix}, \quad (1.2.1.3)$$

*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

1.2.2. The *cross product* or *vector product* of \mathbf{A}, \mathbf{B} is defined as

$$\mathbf{A} \times \mathbf{B} = \begin{pmatrix} \mathbf{A}_{23} \times \mathbf{B}_{23} \\ \mathbf{A}_{31} \times \mathbf{B}_{31} \\ \mathbf{A}_{12} \times \mathbf{B}_{12} \end{pmatrix} \quad (1.2.2.1)$$

1.2.3. Verify that

$$\mathbf{A} \times \mathbf{B} = -\mathbf{B} \times \mathbf{A} \quad (1.2.3.1)$$

1.2.4. The area of a triangle with vertices $\mathbf{A}, \mathbf{B}, \mathbf{C}$ is given by

$$\frac{1}{2} \|(\mathbf{A} - \mathbf{B}) \times (\mathbf{A} - \mathbf{C})\| \quad (1.2.4.1)$$

2 AREA OF A TRIANGLE

2.1. Find the area of a triangle whose vertices are

$$\mathbf{A} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} -4 \\ 6 \end{pmatrix} \text{ and } \mathbf{C} = \begin{pmatrix} -3 \\ -5 \end{pmatrix}.$$

Solution:

2.2. Find the area of a triangle formed by the vertices $\mathbf{A} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 4 \\ 7 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 7 \\ -4 \end{pmatrix}.$

Solution:

2.3. Find the area of a triangle formed by the points $\mathbf{P} = \begin{pmatrix} -1.5 \\ 3 \end{pmatrix}, \mathbf{Q} = \begin{pmatrix} 6 \\ -2 \end{pmatrix}, \mathbf{R} = \begin{pmatrix} -3 \\ 4 \end{pmatrix}.$

Solution:

2.4. Find the area of a triangle having the points

$$\mathbf{A} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \quad (2.4.1)$$

as its vertices.

Solution:

3 ANGLE BETWEEN VECTORS

3.1. Find the angle between the vectors

$$\begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}, \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} \quad (3.1.1)$$

Solution: Let

$$\mathbf{a} = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} \quad (3.1.2)$$

Angle between the vectors is given by,

$$\theta = \cos^{-1} \left(\frac{\mathbf{a}^T \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|} \right) \quad (3.1.3)$$

$$\|\mathbf{a}\| = \sqrt{1^2 + (-2)^2 + 3^2} = \sqrt{14} \quad (3.1.4)$$

$$\|\mathbf{b}\| = \sqrt{3^2 + (-2)^2 + 1^2} = \sqrt{14} \quad (3.1.5)$$

$$\mathbf{a}^T \mathbf{b} = (1)(3) + (-2)(-2) + (3)(1) = 10 \quad (3.1.6)$$

$$\theta = \cos^{-1} \left(\frac{10}{(\sqrt{14})(\sqrt{14})} \right) \quad (3.1.7)$$

$$= \cos^{-1} \left(\frac{10}{14} \right) \quad (3.1.8)$$

$$(3.1.9)$$