

# Points and Vectors

G V V Sharma\*

## CONTENTS

<b>1</b>	<b>Definitions</b>	<b>1</b>
1.1	$2 \times 1$ vectors . . . . .	1
1.2	$3 \times 1$ vectors . . . . .	1
<b>2</b>	<b>Area of a Triangle</b>	<b>2</b>
<b>3</b>	<b>Angle Between Vectors</b>	<b>2</b>

**Abstract**—This book provides a computational approach to school geometry based on the NCERT textbooks from Class 6-12. Links to sample Python codes are available in the text.

Download python codes using

svn co <https://github.com/gadepall/school/trunk/ncert/computation/codes>

1.1.2. The area of the triangle with vertices  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  is given by the absolute value of

$$\frac{1}{2} |\mathbf{A} - \mathbf{B} \quad \mathbf{A} - \mathbf{C}| \quad (1.1.2.1)$$

1.1.3. The transpose of  $\mathbf{A}$  is defined as

$$\mathbf{A}^\top = \begin{pmatrix} a_1 & a_2 \end{pmatrix} \quad (1.1.3.1)$$

1.1.4. The *inner product* or *dot product* is defined as

$$\mathbf{A}^\top \mathbf{B} \equiv \mathbf{A} \cdot \mathbf{B} \quad (1.1.4.1)$$

$$= \begin{pmatrix} a_1 & a_2 \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = a_1 b_1 + a_2 b_2 \quad (1.1.4.2)$$

1.1.5. *norm* of  $\mathbf{A}$  is defined as

$$\|\mathbf{A}\| \equiv |\vec{A}| \quad (1.1.5.1)$$

$$= \sqrt{\mathbf{A}^\top \mathbf{A}} = \sqrt{a_1^2 + a_2^2} \quad (1.1.5.2)$$

Thus,

$$\|\lambda \mathbf{A}\| \equiv |\lambda \vec{A}| \quad (1.1.5.3)$$

$$= |\lambda| \|\mathbf{A}\| \quad (1.1.5.4)$$

1.1.6. The angle between two vectors is given by

$$\theta = \cos^{-1} \frac{\mathbf{A}^\top \mathbf{B}}{\|\mathbf{A}\| \|\mathbf{B}\|} \quad (1.1.6.1)$$

## 1 DEFINITIONS

### 1.1 $2 \times 1$ vectors

1.1.1. Let

$$\mathbf{A} \equiv \vec{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \quad (1.1.1.1)$$

$$\equiv a_1 \vec{i} + a_2 \vec{j}, \quad (1.1.1.2)$$

$$\mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}, \quad (1.1.1.3)$$

be  $2 \times 1$  vectors. Then, the determinant of the  $2 \times 2$  matrix

$$\mathbf{M} = \begin{pmatrix} \mathbf{A} & \mathbf{B} \end{pmatrix} \quad (1.1.1.4)$$

is defined as

$$|\mathbf{M}| = |\mathbf{A} \quad \mathbf{B}| \quad (1.1.1.5)$$

$$= \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1 b_2 - a_2 b_1 \quad (1.1.1.6)$$

### 1.2 $3 \times 1$ vectors

1.2.1. Let

$$\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \equiv a_1 \vec{i} + a_2 \vec{j} + a_3 \vec{k}, \quad (1.2.1.1)$$

$$\mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}, \quad (1.2.1.2)$$

and

$$\mathbf{A}_{ij} = \begin{pmatrix} a_i \\ a_j \end{pmatrix}, \mathbf{B}_{ij} = \begin{pmatrix} b_i \\ b_j \end{pmatrix}, \quad (1.2.1.3)$$

\*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

1.2.2. The *cross product* or *vector product* of  $\mathbf{A}, \mathbf{B}$  is defined as

$$\mathbf{A} \times \mathbf{B} = \begin{pmatrix} \mathbf{A}_{23} \times \mathbf{B}_{23} \\ \mathbf{A}_{31} \times \mathbf{B}_{31} \\ \mathbf{A}_{12} \times \mathbf{B}_{12} \end{pmatrix} \quad (1.2.2.1)$$

1.2.3. Verify that

$$\mathbf{A} \times \mathbf{B} = -\mathbf{B} \times \mathbf{A} \quad (1.2.3.1)$$

1.2.4. The area of a triangle with vertices  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  is given by

$$\frac{1}{2} \|(\mathbf{A} - \mathbf{B}) \times (\mathbf{A} - \mathbf{C})\| \quad (1.2.4.1)$$

## 2 AREA OF A TRIANGLE

2.1. Find the area of a triangle whose vertices are

$$\mathbf{A} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} -4 \\ 6 \end{pmatrix} \text{ and } \mathbf{C} = \begin{pmatrix} -3 \\ -5 \end{pmatrix}.$$

**Solution:**

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} - \begin{pmatrix} -4 \\ 6 \end{pmatrix} = \begin{pmatrix} 5 \\ -7 \end{pmatrix} \quad (2.1.1)$$

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} -1 \\ 11 \end{pmatrix} \quad (2.1.2)$$

Hence, the desired area is

$$\frac{1}{2} \begin{vmatrix} 5 & -1 \\ -7 & 11 \end{vmatrix} = \frac{1}{2} (55 - 7) = 24 \quad (2.1.3)$$

2.2. Find the area of a triangle formed by the vertices  $\mathbf{A} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 4 \\ 7 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 7 \\ -4 \end{pmatrix}$ .

**Solution:**

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} 1 \\ -5 \end{pmatrix} \quad (2.2.1)$$

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} -2 \\ 6 \end{pmatrix} \quad (2.2.2)$$

Hence, the desired area is

$$\frac{1}{2} \begin{vmatrix} 1 & -2 \\ -5 & 6 \end{vmatrix} = \frac{1}{2} (6 - 10) = 2 \quad (2.2.3)$$

after taking the absolute value.

2.3. Find the area of a triangle formed by the points

$$\mathbf{P} = \begin{pmatrix} -1.5 \\ 3 \end{pmatrix}, \mathbf{Q} = \begin{pmatrix} 6 \\ -2 \end{pmatrix}, \mathbf{R} = \begin{pmatrix} -3 \\ 4 \end{pmatrix}.$$

**Solution:**

$$\mathbf{P} - \mathbf{Q} = \begin{pmatrix} -7.5 \\ 5 \end{pmatrix} \quad (2.3.1)$$

$$\mathbf{P} - \mathbf{R} = \begin{pmatrix} 1.5 \\ -1 \end{pmatrix} \quad (2.3.2)$$

Hence, the desired area is the absolute value of

$$\frac{1}{2} \begin{vmatrix} -7.5 & 1.5 \\ 5 & -1 \end{vmatrix} = \frac{1}{2} (7.5 - 7.5) = 0 \quad (2.3.3)$$

This means that the points are on a straight line.

2.4. Find the area of a triangle having the points

$$\mathbf{A} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \quad (2.4.1)$$

as its vertices.

**Solution:**

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} 0 \\ -1 \\ -2 \end{pmatrix}, \mathbf{A} - \mathbf{C} = \begin{pmatrix} -1 \\ -2 \\ 0 \end{pmatrix} \quad (2.4.2)$$

and

$$\begin{vmatrix} -1 & -2 \\ -2 & 0 \end{vmatrix} = -4 \quad (2.4.3)$$

$$\begin{vmatrix} -2 & 0 \\ 0 & -1 \end{vmatrix} = 2 \quad (2.4.4)$$

$$\begin{vmatrix} 0 & -1 \\ -1 & -2 \end{vmatrix} = -1 \quad (2.4.5)$$

$$(2.4.6)$$

From (1.2.2.1),

$$\frac{1}{2} (\mathbf{A} - \mathbf{B}) \times (\mathbf{A} - \mathbf{C}) = \frac{1}{2} \begin{pmatrix} -4 \\ 2 \\ -1 \end{pmatrix} \quad (2.4.7)$$

and from (1.2.2.1), the area of the triangle is

$$\frac{1}{2} \left\| \begin{pmatrix} -4 \\ 2 \\ -1 \end{pmatrix} \right\| = \frac{1}{2} \sqrt{4^2 + 2^2 + 1^2} = \frac{1}{2} \sqrt{21} \quad (2.4.8)$$

## 3 ANGLE BETWEEN VECTORS

3.1. Find the angle between the vectors

$$\begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}, \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} \quad (3.1.1)$$

**Solution:** Let

$$\mathbf{a} = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} \quad (3.1.2)$$

Angle between the vectors is given by,

$$\theta = \cos^{-1} \left( \frac{\mathbf{a}^T \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|} \right) \quad (3.1.3)$$

$$\|\mathbf{a}\| = \sqrt{1^2 + (-2)^2 + 3^2} = \sqrt{14} \quad (3.1.4)$$

$$\|\mathbf{b}\| = \sqrt{3^2 + (-2)^2 + 1^2} = \sqrt{14} \quad (3.1.5)$$

$$\mathbf{a}^T \mathbf{b} = (1)(3) + (-2)(-2) + (3)(1) = 10 \quad (3.1.6)$$

$$\theta = \cos^{-1} \left( \frac{10}{(\sqrt{14})(\sqrt{14})} \right) \quad (3.1.7)$$

$$= \cos^{-1} \left( \frac{10}{14} \right) \quad (3.1.8)$$

$$(3.1.9)$$