
OPTIMIZATION

Through High School Math

G. V. V. Sharma



Copyright ©2022 by G. V. V. Sharma.

<https://creativecommons.org/licenses/by-sa/3.0/>

and

<https://www.gnu.org/licenses/fdl-1.3.en.html>

Contents

Introduction	iii
1 Problem Formulation	1
2 Convex Functons	5
2.1 Definition	5
2.2 Examples	7
3 Gradient Descent	11
3.1 Definition	11
3.2 Examples	12
4 Convex Optimization	25
4.1 Lagrange Multipliers	25
4.2 Inequality Constraints	28
4.3 KKT Conditions	29
5 Geometric Programming	31
6 Linear Programming	33
6.1 Introduction	33
6.2 Applications	46

6.3	Miscellenaeous	58
7	Quadratic Programming	69
8	Semi-Definite Programming	71
A	Convex Functions	77
B	Line	81
C	Manual	83

Introduction

This book introduces optimization through high school math.

Chapter 1

Problem Formulation

1.1. Express the problem of finding the distance of the point $\mathbf{P} = \begin{pmatrix} 3 \\ -5 \end{pmatrix}$ from the line

$$L : \begin{pmatrix} 3 & -4 \end{pmatrix} \mathbf{x} = 26 \quad (1.1.1)$$

as an optimization problem.

Solution: The given problem can be expressed as

$$\min_{\mathbf{x}} g(\mathbf{x}) = \|\mathbf{x} - \mathbf{P}\|^2 \quad (1.1.2)$$

$$\text{s.t.} \quad \mathbf{n}^T \mathbf{x} = c \quad (1.1.3)$$

where

$$\mathbf{n} = \begin{pmatrix} 3 \\ -4 \end{pmatrix} \quad (1.1.4)$$

$$c = 26 \quad (1.1.5)$$

1.2. Explain Problem 1.1 through a plot and find a graphical solution.

1.3. Solve (1.1.2) using `cvxpy`.

Solution: The following code yields

```
codes/opt/line_dist_cvx.py
```

$$\mathbf{x}_{\min} = \begin{pmatrix} 2.64 \\ -4.52 \end{pmatrix}, \quad (1.3.1)$$

$$g(\mathbf{x}_{\min}) = 0.6 \quad (1.3.2)$$

1.4. Convert (1.1.2) to an unconstrained optimization problem.

Solution: L in (1.1.1) can be expressed in terms of the direction vector \mathbf{m} as

$$\mathbf{x} = \mathbf{A} + \lambda \mathbf{m}, \quad (1.4.1)$$

where \mathbf{A} is any point on the line and

$$\mathbf{m}^T \mathbf{n} = 0 \quad (1.4.2)$$

Substituting (1.4.1) in (1.1.2), an unconstrained optimization problem

$$\min_{\lambda} f(\lambda) = \|\mathbf{A} + \lambda \mathbf{m} - \mathbf{P}\|^2 \quad (1.4.3)$$

is obtained.

1.5. Solve (1.4.3).

Solution:

$$f(\lambda) = (\lambda \mathbf{m} + \mathbf{A} - \mathbf{P})^T (\lambda \mathbf{m} + \mathbf{A} - \mathbf{P}) \quad (1.5.1)$$

$$\begin{aligned} &= \lambda^2 \|\mathbf{m}\|^2 + 2\lambda \mathbf{m}^T (\mathbf{A} - \mathbf{P}) \\ &\quad + \|\mathbf{A} - \mathbf{P}\|^2 \end{aligned} \quad (1.5.2)$$

$$\because f^{(2)}\lambda = 2\|\mathbf{m}\|^2 > 0 \quad (1.5.3)$$

the minimum value of $f(\lambda)$ is obtained when

$$f^{(1)}(\lambda) = 2\lambda \|\mathbf{m}\|^2 + 2\mathbf{m}^T (\mathbf{A} - \mathbf{P}) = 0 \quad (1.5.4)$$

$$\implies \lambda_{\min} = -\frac{\mathbf{m}^T (\mathbf{A} - \mathbf{P})}{\|\mathbf{m}\|^2} \quad (1.5.5)$$

Choosing \mathbf{A} such that

$$\mathbf{m}^T (\mathbf{A} - \mathbf{P}) = 0, \quad (1.5.6)$$

substituting in (1.5.5),

$$\lambda_{\min} = 0 \quad \text{and} \quad (1.5.7)$$

$$\mathbf{A} - \mathbf{P} = \mu \mathbf{n} \quad (1.5.8)$$

for some constant μ . (1.5.8) is a consequence of (1.4.2) and (1.5.6). Also, from (1.5.8),

$$\mathbf{n}^T (\mathbf{A} - \mathbf{P}) = \mu \|\mathbf{n}\|^2 \quad (1.5.9)$$

$$\implies \mu = \frac{\mathbf{n}^T \mathbf{A} - \mathbf{n}^T \mathbf{P}}{\|\mathbf{n}\|^2} = \frac{c - \mathbf{n}^T \mathbf{P}}{\|\mathbf{n}\|^2} \quad (1.5.10)$$

from (1.1.3). Substituting $\lambda_{\min} = 0$ in (1.4.3),

$$\min_{\lambda} f(\lambda) = \|\mathbf{A} - \mathbf{P}\|^2 = \mu^2 \|\mathbf{n}\|^2 \quad (1.5.11)$$

upon substituting from (1.5.8). The distance between \mathbf{P} and L is then obtained from (1.5.11) as

$$\|\mathbf{A} - \mathbf{P}\| = |\mu| \|\mathbf{n}\| \quad (1.5.12)$$

$$= \frac{|\mathbf{n}^T \mathbf{P} - c|}{\|\mathbf{n}\|} \quad (1.5.13)$$

after substituting for μ from (1.5.10). Using the corresponding values from Problem (1.1) in (1.5.13),

$$\min_{\lambda} f(\lambda) = 0.6 \quad (1.5.14)$$

Chapter 2

Convex Functions

2.1. Definition

2.1.1. The following python script plots

$$f(\lambda) = a\lambda^2 + b\lambda + d \tag{2.1.1.1}$$

for

$$a = \|\mathbf{m}\|^2 > 0 \tag{2.1.1.2}$$

$$b = \mathbf{m}^T (\mathbf{A} - \mathbf{P}) \tag{2.1.1.3}$$

$$c = \|\mathbf{A} - \mathbf{P}\|^2 \tag{2.1.1.4}$$

where \mathbf{A} is the intercept of the line L in (1.1.1) on the x-axis and the points

$$\mathbf{U} = \begin{pmatrix} \lambda_1 \\ f(\lambda_1) \end{pmatrix}, \mathbf{V} = \begin{pmatrix} \lambda_2 \\ f(\lambda_2) \end{pmatrix} \quad (2.1.1.5)$$

$$\mathbf{X} = \begin{pmatrix} t\lambda_1 + (1-t)\lambda_2 \\ f[t\lambda_1 + (1-t)\lambda_2] \end{pmatrix}, \quad (2.1.1.6)$$

$$\mathbf{Y} = \begin{pmatrix} t\lambda_1 + (1-t)\lambda_2 \\ tf(\lambda_1) + (1-t)f(\lambda_2) \end{pmatrix} \quad (2.1.1.7)$$

for

$$\lambda_1 = -3, \lambda_2 = 4, t = 0.3 \quad (2.1.1.8)$$

in Fig. 2.1. Geometrically, this means that any point \mathbf{Y} between the points \mathbf{U}, \mathbf{V} on the line UV is always above the point \mathbf{X} on the curve $f(\lambda)$. Such a function f is defined to be convex function

opt/codes/1.2.py

2.1.2. Show that

$$f[t\lambda_1 + (1-t)\lambda_2] \leq tf(\lambda_1) + (1-t)f(\lambda_2) \quad (2.1.2.1)$$

for $0 < t < 1$. This is true for any convex function.

2.1.3. Show that

$$(2.1.2.1) \implies f^{(2)}(\lambda) > 0 \quad (2.1.3.1)$$




Figure 2.1: $f(\lambda)$ versus λ

2.1.4. Show that a convex function has a unique minimum.

2.2. Examples

Determine if the following are convex functions

2.1 (a) Determine whether the function $f(x) = (2x - 1)^2 + 3$ is convex or not.

Solution: The given function is

$$f(x) = (2x - 1)^2 + 3 \quad (2.1.1)$$

$$= 4x^2 + 4x + 4 \quad (2.1.2)$$

$$\therefore a = 4, > 0 \quad (2.1.3)$$

Hence, the function in equation (2.1.1) is convex from A.1. The figure is as shown in Fig2.1.1




Figure 2.1.1:

2.2 At what points in the interval $(0, 2\pi)$ does the function $\sin 2x$ attain its maximum value.

2.3 Find the absolute maximum and minimum values of the function f given by

$$f(x) = \cos^2 x + \sin x, \quad x \in [0, \pi] \quad (2.3.1)$$

Chapter 3

Gradient Descent

3.1. Definition

3.1.1. Find a numerical solution for (1.4.3)

(1.4.3)

Solution: A numerical solution for (1.4.3) is obtained as

$$\lambda_{n+1} = \lambda_n - \mu f'(\lambda_n) \quad (3.1.1.1)$$

$$= \lambda_n - \mu (2a\lambda_n + b) \quad (3.1.1.2)$$

where λ_0 is an initial guess and μ is a variable parameter. The choice of these parameters is very important since they decide how fast the algorithm converges.

3.1.2. Write a program to implement (3.1.1.2).

Solution: Download and execute

`opt/codes/gd.py`

3.1.3. Find a closed form solution for λ_n in (3.1.1.2) using the one sided Z transform.

3.1.4. Find the condition for which (3.1.1.2) converges, i.e.

$$\lim_{n \rightarrow \infty} |\lambda_{n+1} - \lambda_n| = 0 \quad (3.1.4.1)$$

3.2. Examples

3.1 Find the maximum and minimum values of

- (a) Find the minimum value of the function $f(x) = (2x - 1)^2 + 3$ using Gradient Descent method.

Solution: The given function has a minimum value as shown in Figure 3.1.1.

$$f'(x) = 8x - 4 \quad (3.1.1)$$

The minimum value of the function is calculated using Gradient Descent method as below

$$x_{n+1} = x_n - \alpha \nabla f(x_n) \quad (3.1.2)$$

Choosing

- i. $\alpha = 0.001$
- ii. precision = 0.0000001
- iii. n = 10000000
- iv. $x_0 = -5$

$$x_{min} = \frac{1}{2}, f(x)_{min} = 3 \quad (3.1.3)$$




Figure 3.1.1:

3.2 Find the maximum profit that a company can make if the profit function is given by

$$f(x) = 41 - 72x + 18x^2 \quad (3.2.1)$$

Solution: Considering

$$\lambda (41 - 72x_1 - 18x_1^2) + (1 - \lambda) (41 - 72x_2 - 18x_2^2) \geq \quad (3.2.2)$$

$$41 - 72 (\lambda x_1 + (1 - \lambda)x_2) - 18 (\lambda x_1 + (1 - \lambda)x_2)^2, \quad (3.2.3)$$

we obtain

$$18x_1^2 (\lambda^2 - \lambda) + 18x_2^2 (\lambda^2 - \lambda) + 36x_1x_2 (\lambda^2 - \lambda) \geq 0 \quad (3.2.4)$$

$$x_1^2 (\lambda^2 - \lambda) + x_2^2 (\lambda^2 - \lambda) + 2x_1x_2 (\lambda^2 - \lambda) \geq 0 \quad (3.2.5)$$

$$-\lambda (1 - \lambda) (x_1 - x_2)^2 \geq 0 \quad (3.2.6)$$

$$\implies \lambda (1 - \lambda) (x_1 - x_2)^2 \leq 0 \quad (3.2.7)$$

which is false for all $\lambda \in (0, 1)$. Hence the given function $f(x)$ is concave. Using the gradient ascent method,

$$x_n = x_{n-1} + \mu \frac{df(x)}{dx} \quad (3.2.8)$$

Since

$$\frac{df(x)}{dx} = -72 - 36x, \quad (3.2.9)$$

substituting (3.2.9) in (3.2.8),

$$x_n = x_{n-1} + \mu(-72 - 36x_{n-1}) \quad (3.2.10)$$

Choosing

$$x_0 = 1, \alpha = 0.001, \text{precision} = 0.00000001, \quad (3.2.11)$$

$$f_{\max} \approx 113, x_{\max} \approx -2.0, \quad (3.2.12)$$

which is verified in Fig. 3.2.1.




Figure 3.2.1:

3.3 Find both the maximum value and the minimum value of

$$f(x) = 3x^4 - 8x^3 + 12x^2 - 48x + 25 = 0 \quad x \in (0, 3) \quad (3.3.1)$$

Solution:




Figure 3.3.1:

$$\frac{df(x)}{dx} = 12x^3 - 24x^2 + 24x - 48 \quad (3.3.2)$$

The minimum can be found using

$$x_{n+1} = x_n - \alpha \frac{df(x)}{dx} \quad (3.3.3)$$

$$= x_n - \alpha(12x_n^3 - 24x_n^2 + 24x_n - 48) \quad (3.3.4)$$

where

(a) $\alpha = 0.001$

(b) x_{n+1} is current value

(c) x_n is previous value

(d) precession = 0.00000001

(e) maximum iterations = 100000000

as

$$f_{min} = -39 \quad (3.3.5)$$

$$x_{min} = 2 \quad (3.3.6)$$

3.4 At what points in the interval $(0, 2\pi)$ does the function $\sin 2x$ attain its maximum value.

Solution: Since

$$f(x) = \sin 2x, \quad (3.4.1)$$

$$f'(x) = 2 \cos 2x \quad (3.4.2)$$




Figure 3.4.1:

Using gradient ascent,

$$x_{n+1} = x_n + \alpha \nabla f(x_n) \quad (3.4.3)$$

$$= x_n + \alpha(2 \cos 2x) \quad (3.4.4)$$

Choosing

$$x_0 = 0.5, \alpha = 0.001, \text{precision} = 0.00000001, \quad (3.4.5)$$

$$f_{max} = 1.0000, x_{max} = 0.7854. \quad (3.4.6)$$

3.5 Find the maximum value of $2x^3 - 24x + 107$ in the interval $[1, 3]$. Find the maximum value of the same function in $[-3, -1]$.

Solution: Using gradient ascent method,

$$x_n = x_{n-1} + \mu \frac{df(x)}{dx} \quad (3.5.1)$$

where

$$\frac{df(x)}{dx} = 6x^2 - 24 \quad (3.5.2)$$

yielding

$$x_n = x_{n-1} + \mu(6x_{n-1}^2 - 24) \quad (3.5.3)$$

Choosing

$$x_0 = 1, \mu = 0.001 \text{ and precision} = 0.00000001, \quad (3.5.4)$$

$$f_{max} \approx 139, x_{max} \approx -2.0 \quad (3.5.5)$$

3.6 It is given that at $x=1$, the function $x^4 - 62x^2 + ax + 9$ attains its maximum value,




Figure 3.5.1:

on the interval $[0,2]$. Find the value of a .

Solution: Differentiating the given function,

$$\nabla f(x) = 4x^3 - 124x + a \quad (3.6.1)$$

Since f attains its maximum value on the interval $[0,2]$ at $x = 1$,

$$\nabla f(1) = 0 \implies a = 120 \quad (3.6.2)$$

Using gradient descent,

$$x_{n+1} = x_n + \alpha \nabla f(x_n) \quad (3.6.3)$$

$$= x_n + \alpha (4x_n^3 - 124x_n + 120) \quad (3.6.4)$$

and choosing

$$x_0 = 0.5, \alpha = 0.001 \text{ and precision} = 0.00000001, \quad (3.6.5)$$

$$f_{max} = 68, x_{max} = 1 \quad (3.6.6)$$

3.7 Find the absolute maximum and minimum values of the function f given by

$$f(x) = \cos^2 x + \sin x, \quad x \in [0, \pi] \quad (3.7.1)$$

Solution: The derivative of the given function is

$$\nabla f(x) = \cos x - 2 \sin x \cos x \quad (3.7.2)$$

The maxima is calculated by

$$x_{n+1} = x_n + \alpha \nabla f(x_n) \quad (3.7.3)$$

$$= x_n + \alpha (\cos x_n - 2 \sin x_n \cos x_n) \quad (3.7.4)$$

where

(a) $x_0 = 0.5$




Figure 3.6.1:

(b) $\alpha = 0.001$

(c) precision = 0.00000001

yielding

$$f_{max} = 1.25, x_{max} = 0.52. \quad (3.7.5)$$




Figure 3.7.1:

The minima is found by

$$x_{n+1} = x_n - \alpha \nabla f(x_n) \quad (3.7.6)$$

$$= x_n - \alpha (\cos x_n - 2 \sin x_n \cos x_n) \quad (3.7.7)$$

Chapter 4

Convex Optimization

4.1. Lagrange Multipliers

4.1.1. Find

$$\min_{\mathbf{x}} g(\mathbf{x}) = \|\mathbf{x} - \mathbf{P}\|^2 = r^2 \quad (4.1.1.1)$$

$$\text{s.t. } h(\mathbf{x}) = \mathbf{n}^T \mathbf{x} - c = 0 \quad (4.1.1.2)$$

by plotting the circles $g(\mathbf{x})$ for different values of r along with the line $g(\mathbf{x})$.

Solution: The following code plots Fig. 4.1.1.1

```
opt/codes/concirc.py
```

4.1.2. By solving the quadratic equation obtained from (4.1.1.1), show that

$$\min_{\mathbf{x}} r = \frac{3}{5}, \mathbf{x}_{\min} = \mathbf{Q} = \begin{pmatrix} 2.64 \\ -4.52 \end{pmatrix} \quad (4.1.2.1)$$

In Fig. 4.1.1.1, it can be seen that \mathbf{Q} is the point of contact of the line L with the circle of minimum radius $r = \frac{3}{5}$.




Figure 4.1.1.1: Finding $\min_{\mathbf{x}} g(\mathbf{x})$

4.1.3. Show that

$$\nabla h(\mathbf{x}) = \begin{pmatrix} 3 \\ -4 \end{pmatrix} = \mathbf{n} \quad (4.1.3.1)$$

where

$$\nabla = \begin{pmatrix} \frac{\partial}{\partial x_1} \\ \frac{\partial}{\partial x_2} \end{pmatrix} \quad (4.1.3.2)$$

4.1.4. Show that

$$\nabla g(\mathbf{x}) = 2 \left\{ \mathbf{x} - \begin{pmatrix} 3 \\ -5 \end{pmatrix} \right\} = 2 \{\mathbf{x} - \mathbf{P}\} \quad (4.1.4.1)$$

4.1.5. From Fig. 4.1.1.1, show that

$$\nabla g(\mathbf{Q}) = \lambda \nabla h(\mathbf{Q}), \quad (4.1.5.1)$$

Solution: In Fig. 4.1.1.1, PQ is the normal to the line L , represented by $h(\mathbf{x})$. \therefore the normal vector of L is in the same direction as PQ , for some constant k ,

$$(\mathbf{Q} - \mathbf{P}) = k\mathbf{n} \quad (4.1.5.2)$$

which is the same as (4.1.5.1) after substituting from (4.1.3.1). and (4.1.4.1).

4.1.6. Use (4.1.5.1) and $\mathbf{h}(\mathbf{Q}) = 0$ from (4.1.1.2) to obtain \mathbf{Q} .

Solution: From the given equations, we obtain

$$(\mathbf{Q} - \mathbf{P}) - \lambda \mathbf{n} = 0 \quad (4.1.6.1)$$

$$\mathbf{n}^T \mathbf{Q} - c = 0 \quad (4.1.6.2)$$

which can be simplified to obtain

$$\begin{pmatrix} \mathbf{I} & -\mathbf{n} \\ \mathbf{n}^T & 0 \end{pmatrix} \begin{pmatrix} \mathbf{Q} \\ \lambda \end{pmatrix} = \begin{pmatrix} \mathbf{P} \\ c \end{pmatrix} \quad (4.1.6.3)$$

The following code computes the solution to (4.1.6.3)

```
opt/codes/lagmul.py
```

4.1.7. Define

$$C(\mathbf{x}, \lambda) = g(\mathbf{x}) - \lambda h(\mathbf{x}) \quad (4.1.7.1)$$

and show that \mathbf{Q} can also be obtained by solving the equations

$$\nabla C(\mathbf{x}, \lambda) = 0. \quad (4.1.7.2)$$

What is the sign of λ ? C is known as the Lagrangian and the above technique is known as the Method of Lagrange Multipliers.

4.1.8. Obtain \mathbf{Q} using gradient descent.

Solution:

```
opt/codes/gd_lagrange.py
```

4.2. Inequality Constraints

4.2.1. Modify the code in problem 4.1.1 to find a graphical solution for minimising

$$f(\mathbf{x}) \quad (4.2.1.1)$$

with constraint

$$g(\mathbf{x}) \geq 0 \quad (4.2.1.2)$$

Solution: This problem reduces to finding the radius of the smallest circle in the

shaded area in Fig. 1.1 . It is clear that this radius is 0.

```
opt/codes/2.4.py
```




Figure 1.1: Smallest circle in the shaded region is a point.

4.3. KKT Conditions

Chapter 5

Geometric Programming

- 5.1 Find the maximum area of an isosceles triangle inscribed in the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ with its vertex at one end of the major axis.

Solution:

- 5.2 A tank with rectangular base and rectangular sides, open at the top is to be constructed so that its depth is 2 m and volume is $8m^3$. If building of tank costs Rs 70 per sq metres for the base and Rs 45 per square metre for sides. What is the cost of least expensive tank?

Solution:

Let l, b and h be the length, width and height of a tank. The volume of tank is given by,

$$V = lbh \quad (5.2.1)$$

$$h = 2 \quad (5.2.2)$$

Cost of Building

$$R_b = 70/m^2 \tag{5.2.3}$$

$$R_s = 45/m^2 \tag{5.2.4}$$

The given problem can then be formulated as

$$S = \min_{l,b} R_b(lb) + R_s(4(l+b)) \tag{5.2.5}$$

$$\text{s.t } lb = 4 \tag{5.2.6}$$

which is a disciplined geometric programming (DGP) problem that can be solved using cvxpy. DGP is a subset of log-log-convex program (LLCP).

Chapter 6

Linear Programming

6.1. Introduction

6.1.1 Maximize

$$Z = 3x + 4y \tag{6.1.1.1}$$

subject to the constraints:

$$x + 4y \leq 4, \tag{6.1.1.2}$$

$$x \geq 0, y \geq 0 \tag{6.1.1.3}$$

Solution:

(a) Using cvxpy method: The given problem can be formulated as

$$\max_{\mathbf{x}} Z = \begin{pmatrix} 3 & 4 \end{pmatrix} \mathbf{x} \tag{6.1.1.4}$$

$$\begin{pmatrix} 1 & 4 \\ -1 & 0 \\ 0 & -1 \end{pmatrix} \mathbf{x} \preceq \begin{pmatrix} 4 \\ 0 \\ 0 \end{pmatrix} \tag{6.1.1.5}$$

Solving using cvxpy, we get

$$\max_{\mathbf{x}} Z = 12, \mathbf{x} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \quad (6.1.1.6)$$

(b) Using Corner point method: The corner points of the inequalities are:

$$\mathbf{A} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (6.1.1.7)$$

$$\mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (6.1.1.8)$$

$$\mathbf{x} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \quad (6.1.1.9)$$

Substituting above values of corner points in Equation (6.1.1.1) to get the value of Z , as shown in the Table 6.1.1.2

Corner Point	Corresponding Z value
$\mathbf{A} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$	4
$\mathbf{B} \begin{pmatrix} 0 \\ 0 \end{pmatrix}$	0
$\mathbf{x} \begin{pmatrix} 4 \\ 0 \end{pmatrix}$	12

Table 6.1.1.2:

From the table 6.1.1.2, it is clear that the optimum value and optimum point are similar to what we found in (6.1.1.6).

The relevant figure is as shown in 6.1.1.1




Figure 6.1.1.1:

6.1.2 Minimise

$$Z = -3x + 4y \quad (6.1.2.1)$$

such that

$$x + 2y < 8, \quad (6.1.2.2)$$

$$3x + 2y < 12, \quad (6.1.2.3)$$

$$x > 0, y > 0 \quad (6.1.2.4)$$

Solution: The given problem can be formulated as




Figure 6.1.2.1:

$$\min_{\mathbf{x}} Z = \begin{pmatrix} -3 & 4 \end{pmatrix} \mathbf{x} \quad (6.1.2.5)$$

$$\begin{pmatrix} 1 & 2 \\ 3 & 2 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} \succeq \begin{pmatrix} 8 \\ 12 \\ 0 \\ 0 \end{pmatrix} \quad (6.1.2.6)$$

Solving above equations using cvxpy, we get

$$\min_{\mathbf{x}} Z = -12 \quad (6.1.2.7)$$

$$\mathbf{x} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \quad (6.1.2.8)$$

6.1.3 Maximize $Z = 5x + 3y$ such that

$$3x + 5y \leq 15, \quad (6.1.3.1)$$

$$5x + 2y \leq 10, \quad (6.1.3.2)$$

$$x \geq 0, y \geq 0. \quad (6.1.3.3)$$

Solution: The given problem can be expressed as

$$Z = \begin{pmatrix} 5 & 3 \end{pmatrix} \mathbf{x} \quad (6.1.3.4)$$

$$s.t. \quad \begin{pmatrix} 3 & 5 \\ 5 & 2 \end{pmatrix} \mathbf{x} \leq \begin{pmatrix} 15 \\ 10 \end{pmatrix} \quad (6.1.3.5)$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} \geq \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (6.1.3.6)$$

$$(6.1.3.7)$$

Using cvxpy, the solution is

$$\mathbf{x} = \begin{pmatrix} \frac{20}{19} \\ \frac{45}{19} \end{pmatrix}, Z_{max} = \frac{235}{19} \quad (6.1.3.8)$$

6.1.4 Minimize $Z = 3x + 5y$ such that

$$x + 3y \geq 3 \quad (6.1.4.1)$$

$$x + y \geq 2 \quad (6.1.4.2)$$

$$x \geq 0, y \geq 0. \quad (6.1.4.3)$$

Solution: The given problem can be expressed as

$$Z = \min_{\mathbf{x}} \begin{pmatrix} 3 & 5 \end{pmatrix} \mathbf{x} \quad (6.1.4.4)$$

$$\begin{pmatrix} 1 & 3 \\ 1 & 1 \end{pmatrix} \mathbf{x} \succeq \begin{pmatrix} 3 \\ 2 \end{pmatrix} \quad (6.1.4.5)$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} \succeq \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (6.1.4.6)$$

Solving using cvxpy, we get,

$$\mathbf{x} = \begin{pmatrix} \frac{3}{2} \\ \frac{1}{2} \end{pmatrix}, Z_{min} = 7 \quad (6.1.4.7)$$

6.1.5

6.1.6 Minimize $Z=x+2y$ subject to

$$2x + 3y \geq 3 \quad (6.1.6.1)$$

$$x + 2y \geq 6 \quad (6.1.6.2)$$

$$x, y \geq 0. \quad (6.1.6.3)$$

Solution: The optimization problem can be defined as




Figure 6.1.6.1:

$$P = \min_{\mathbf{x}} \begin{pmatrix} 1 & 2 \end{pmatrix} \mathbf{x} \quad (6.1.6.4)$$

$$\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \mathbf{x} \succeq \begin{pmatrix} 3 \\ 6 \end{pmatrix} \quad (6.1.6.5)$$

$$x, y \geq \mathbf{0} \quad (6.1.6.6)$$

From Fig. 6.1.6.1, the feasible region vertices are

$$\begin{pmatrix} 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (6.1.6.7)$$

yielding

$$\begin{pmatrix} 1 & 2 \end{pmatrix} \begin{pmatrix} 0 \\ 3 \end{pmatrix} = 6 \quad (6.1.6.8)$$

$$\begin{pmatrix} 1 & 2 \end{pmatrix} \begin{pmatrix} 6 \\ 0 \end{pmatrix} = 6 \quad (6.1.6.9)$$

$$(6.1.6.10)$$

Thus, the minimum value of Z is 6.

6.1.7 Minimise and Maximise

$$Z = 5x + 10y \quad (6.1.7.1)$$

subject to

$$x + 2y \leq 120 \quad (6.1.7.2)$$

$$x + y \geq 60 \quad (6.1.7.3)$$

$$x - 2y \geq 0 \quad (6.1.7.4)$$

$$x \geq 0, y \geq 0 \quad (6.1.7.5)$$

Solution: The given problem can be formulated as




Figure 6.1.7.1:

$$\min_{\mathbf{x}} \mathbf{Z} = \begin{pmatrix} 5 & 10 \end{pmatrix} \mathbf{x} \quad (6.1.7.6)$$

$$\max_{\mathbf{x}} \mathbf{Z} = \begin{pmatrix} 5 & 10 \end{pmatrix} \mathbf{x} \quad (6.1.7.7)$$

$$s.t. \quad \begin{pmatrix} -1 & -2 \\ 1 & 1 \\ 1 & -2 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} \succeq \begin{pmatrix} -120 \\ 60 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad (6.1.7.8)$$

$$(6.1.7.9)$$

Solving above equations using cvxpy,

$$\min_{\mathbf{x}} Z = 300, \mathbf{x} = \begin{pmatrix} 60 \\ 0 \end{pmatrix} \quad (6.1.7.10)$$

$$\max_{\mathbf{x}} Z = 600, \mathbf{x} = \begin{pmatrix} 60 \\ 30 \end{pmatrix} \quad (6.1.7.11)$$

6.1.8

6.1.9 Maximise

$$Z = -x + 2y \quad (6.1.9.1)$$

subject to the constraints

$$x + y \geq 5 \quad (6.1.9.2)$$

$$x + 2y \geq 6 \quad (6.1.9.3)$$

$$x \geq 3, y \geq 0. \quad (6.1.9.4)$$

Solution: The given problem can be expressed as

$$z = \max_{\mathbf{x}} \begin{pmatrix} -1 & 2 \end{pmatrix} \mathbf{x} \quad (6.1.9.5)$$

$$s.t. \quad \begin{pmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 5 \\ 6 \\ 3 \\ 0 \end{pmatrix} \quad (6.1.9.6)$$




Figure 6.1.9.1:

By providing the objective function and constraints to cvxpy, the optimal value gives infinity as result and the problem is unbounded. This is verified from Fig. 6.1.9.1.

6.1.10 Maximize

$$Z = x + y \quad (6.1.10.1)$$

subject to

$$x - y \leq -1 \quad (6.1.10.2)$$

$$-x + y \leq 0 \quad (6.1.10.3)$$

$$x, y \geq 0 \quad (6.1.10.4)$$

Solution: From Fig. 6.1.10.1, the given problem has no optimal solution. This is




Figure 6.1.10.1:

verified from cvxpy by considering the following optimization problem.

$$z = \max_x \begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} \quad (6.1.10.5)$$

$$s.t. \quad \begin{pmatrix} 1 & -1 \\ -1 & 0 \\ 0 & -1 \\ -1 & 0 \end{pmatrix} x \preceq \begin{pmatrix} -1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad (6.1.10.6)$$

6.2. Applications

6.2.1 Reshma wishes to mix two types of food P and Q in such a way that the vitamin contents of the mixture contain at least 8 units of vitamin A and 11 units of vitamin B. Food P costs Rs 60/kg and Food Q costs Rs 80/kg. Food P contains 3 units/kg of Vitamin A and 5 units / kg of Vitamin B while food Q contains 4 units/kg of Vitamin A and 2 units/kg of vitamin B. Determine the minimum cost of the mixture.

Solution: Let the mixture contain x kg of food and y kg of food. The given information can be compiled in a table as and which can be expressed in vector form as

	Vitamin A(units/kg)	Vitamin B(units/kg)	Cost(Rs/kg)
Food P	3	5	60
Food Q	4	2	80
Requirement(units/kg)	8	11	

Table 6.2.1.1:

$$P = \min_{\mathbf{x}} \begin{pmatrix} 60 & 80 \end{pmatrix} \mathbf{x} \quad (6.2.1.1)$$

$$\begin{pmatrix} 3 & 4 \\ 5 & 2 \end{pmatrix} \mathbf{x} \succeq \begin{pmatrix} 8 \\ 11 \end{pmatrix} \quad (6.2.1.2)$$

$$\mathbf{x} \succeq \mathbf{0} \quad (6.2.1.3)$$

Solving using cvxpy, we get

$$P_{min} = 159.99999999 \quad (6.2.1.4)$$

$$\mathbf{x} = \begin{pmatrix} 2.11436236 \\ 0.41422823 \end{pmatrix} \quad (6.2.1.5)$$

6.2.2 One kind of cake requires 200g of flour and 25g of fat, and another kind of cake requires 100g of flour and 50g of fat. Find the maximum number of cakes which can be made from 5kg of flour and 1 kg of fat assuming that there is no shortage of the other ingredients used in making the cakes.

Solution: Let x, y be the number of cakes of first kind and second kind that can be made from the given amount of flour and fat respectively. From the given information,

Kind of cake	No. of cakes	Flour (in gm)	Fat (in gm)
$Cake_1$	x	200	25
$Cake_2$	y	100	50

Table 6.2.2.1:

$$200x + 100y \leq 5000 \quad (6.2.2.1)$$

$$100x + 50y \leq 1000 \quad (6.2.2.2)$$

Let P be the maximum number of cakes that can be made from the given amount of flour and fat. The problem can be formulated as

$$P = \max_{\mathbf{x}} \begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} \quad (6.2.2.3)$$

$$\begin{pmatrix} 200 & 100 \\ 100 & 50 \end{pmatrix} \mathbf{x} \leq \begin{pmatrix} 5000 \\ 1000 \end{pmatrix} \quad (6.2.2.4)$$

$$\mathbf{x} \geq \mathbf{0} \quad (6.2.2.5)$$

Solving the above equations using `cvxpy`, we get

$$P_{max} = 30, \mathbf{x} = \begin{pmatrix} 20 \\ 10 \end{pmatrix} \quad (6.2.2.6)$$

6.2.3 A factory makes tennis rackets and cricket bats. A tennis racket takes 1.5 hours of machine time and 3 hours of craftman's time in its making while a cricket bat takes 3 hour of machine time and 1 hour of craftman's time. In a day, the factory has the availability of not more than 42 hours of machine time and 24 hours of craftsman's time.

- (a) What number of rackets and bats must be made if the factory is to work at full capacity?
- (b) If the profit on a racket and on a bat is Rs 20 and Rs 10 respectively, find the maximum profit of the factory when it works at full capacity.

Solution: The given information is summarized in Table 6.2.3.1. From the given

Item	Number	Machine hours	Craftman's hours	Profit
Tennis Rackets	x	1.5	3	Rs.20
Cricket Bats	y	3	1	Rs.10
Maximum time available		42	24	

Table 6.2.3.1:




Figure 6.2.3.1:

information, the optimization problem can be expressed as

$$Z = \max_{\mathbf{x}} \begin{pmatrix} 20 & 10 \end{pmatrix} \mathbf{x} \tag{6.2.3.1}$$

$$s.t. \quad \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix} \mathbf{x} \preceq \begin{pmatrix} 28 \\ 24 \end{pmatrix} \tag{6.2.3.2}$$

$$\mathbf{x} \succeq \mathbf{0} \tag{6.2.3.3}$$

From Fig. 6.2.3.1, the values at the corner points are obtained in Table 6.2.3.2.

Corner points	Value of Z
(0,14)	140
(4,12)	200
(8,0)	160

Table 6.2.3.2:

At full capacity,

$$\mathbf{x} = \begin{pmatrix} 4 \\ 12 \end{pmatrix}, \quad (6.2.3.4)$$

$$\Rightarrow Z = \begin{pmatrix} 20 & 10 \end{pmatrix} \begin{pmatrix} 4 \\ 12 \end{pmatrix} \quad (6.2.3.5)$$

$$= 200 \quad (6.2.3.6)$$

This is verified using cvxpy.

6.2.4 A manufacturer produces nuts and bolts. It takes 1 hour of work on machine A and 3 hours on machine B to produce a package of nuts. It takes 3 hours on machine A and 1 hour on machine B to produce a package of bolts. He earns a profit of Rs17.50 per package on nuts and Rs 7.00 per package on bolts.

How many packages of each should be produced each day so as to maximise his profit, if he operates his machines for at the most 12 hours a day?.

Solution: Table 6.2.4.1 summarizes the given information. The optimization problem

Symbol	Name	Machine A	Machine B	Profit
x	nuts	1x	3x	17.5x
y	bolts	3y	1y	7y
Sum	x+y	x+3y	3x+y	17.5x+7y
t	time	12 h	12 h	

Table 6.2.4.1:

is formulated as

$$z = \begin{pmatrix} 17.5 & 7 \end{pmatrix} \mathbf{x} \quad (6.2.4.1)$$

$$\begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} \mathbf{x} \preceq \begin{pmatrix} 12 \\ 12 \end{pmatrix} \quad (6.2.4.2)$$

Fig. 6.2.4.1 represents the given constraints from which, the corner points of the feasible region intersected by above two lines are

$$\mathbf{0}, \mathbf{A} = \begin{pmatrix} 0 \\ 4 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \text{ and } \mathbf{C} = \begin{pmatrix} 3 \\ 3 \end{pmatrix} \quad (6.2.4.3)$$

The values of Z at these points are listed in Table 6.2.4.2. Thus, the maximum profit

Corner Point	z=17.5x+7y	Remarks
O= $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	0	
A= $\begin{pmatrix} 0 \\ 4 \end{pmatrix}$	28	
C= $\begin{pmatrix} 3 \\ 3 \end{pmatrix}$	73.5	Maximum
B= $\begin{pmatrix} 4 \\ 0 \end{pmatrix}$	70	

Table 6.2.4.2:




Figure 6.2.4.1:

is

$$z = 73.50, \mathbf{x} = \begin{pmatrix} 3 \\ 3 \end{pmatrix}. \quad (6.2.4.4)$$

6.2.5 A factory manufactures two types of screws , A and B. Each type screw requires the use of two machines , an automatic and a hand operated. It takes 4 minutes on the automatic and 6 minutes on hand operated machines to manufacture a package of screws A, while it takes 6 minutes on automatic and 3 minutes on the hand operated

machines to manufacture a package of screws B. Each machine is available for at the most 4hrs on any day. The manufacturer can sell a package of screws A at a profit of Rs. 7 and screws B at a profit of Rs. 10. Assuming that he can sell all the screws he manufactures, how many packages of each type should the factory owner produce in a day in order to maximise his profit? Determine the maximum profit.

Solution: The given information is summarized in Table 6.2.5.1 resulting in the

Item	Number	Machine A	Machine B	Profit
Screw A	x	4(min)	6(min)	7
SCREW B	y	6(min)	3(min)	10
Max.Time available		4 hrs	4 hrs	

Table 6.2.5.1:

following optimization problem.

$$z = \max_{\mathbf{x}} \begin{pmatrix} 7 & 10 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad (6.2.5.1)$$

$$s.t. \quad \begin{pmatrix} 2 & 3 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \preceq \begin{pmatrix} 120 \\ 80 \end{pmatrix} \quad (6.2.5.2)$$

$$\mathbf{x} \succeq \mathbf{0} \quad (6.2.5.3)$$

The maximum profit is obtained as

$$\begin{pmatrix} 2 & 3 & 120 \\ 2 & 1 & 80 \end{pmatrix} \longrightarrow R_2 - R_1 \begin{pmatrix} 2 & 3 & 120 \\ 0 & -2 & -40 \end{pmatrix} \longrightarrow R_2 * \frac{-1}{2} \quad (6.2.5.4)$$

$$\begin{pmatrix} 2 & 3 & 120 \\ 0 & 1 & 20 \end{pmatrix} \longrightarrow R_1 - 3R_2 \begin{pmatrix} 2 & 0 & 60 \\ 0 & 1 & 20 \end{pmatrix} \longrightarrow \frac{R_1}{2} \begin{pmatrix} 1 & 0 & 30 \\ 0 & 1 & 20 \end{pmatrix} \quad (6.2.5.5)$$

yielding the maximum profit

$$z = 410 \quad (6.2.5.6)$$

at

$$\mathbf{x} = \begin{pmatrix} 30 \\ 20 \end{pmatrix}. \quad (6.2.5.7)$$

This is verified by Table 6.2.5.2, where the corner points are obtained from Fig. 6.2.5.1.




Figure 6.2.5.1:

Corner points	Value of Z
(0,40)	400
(30,20)	410
(40,0)	280

Table 6.2.5.2:

6.2.6

6.2.7

6.2.8 A merchant plans to sell two types of personal computers, a desktop model and a portable model that will cost Rs 25000 and Rs 40000 respectively. He estimates that the total monthly demand of computers will not exceed 250 units. Determine the number of units of each type of computers which the merchant should stock to get maximum profit if he does not want to invest more than Rs 70 lakhs and if his profit on the desktop model is Rs 4500 and on portable model is Rs 5000.

Solution: Table 6.2.8.1 summarizes the given information. The optimization problem

Item	Number	Cost	Profit
Desktop	x	25000	4500
Portable Computers	y	40000	5000
Max Investment		7000000	

Table 6.2.8.1:

can then be summarized as

$$Z = \begin{pmatrix} 4500 & 5000 \end{pmatrix} \mathbf{x} \quad (6.2.8.1)$$

$$s.t. \quad \begin{pmatrix} 1 & 1 \\ 5 & 8 \end{pmatrix} \mathbf{x} \preceq \begin{pmatrix} 250 \\ 1400 \end{pmatrix} \quad (6.2.8.2)$$

$$\mathbf{x} \succeq \mathbf{0} \quad (6.2.8.3)$$




Figure 6.2.8.1:

From Fig. 6.2.8.1, the corner points are listed in Table 6.2.8.2 yielding the solution

$$Z = 1150000, \mathbf{x} = \begin{pmatrix} 200 \\ 50 \end{pmatrix} \quad (6.2.8.4)$$

Corner points	Value of Z
(250,0)	112500
(200,50)	1150000
(0,175)	875000

Table 6.2.8.2:

6.2.9

6.2.10 There are two types of fertilisers F_1 and F_2 . F_1 consists of 10% Nitrogen and 6% Phosphoric acid and F_2 consists of 5% Nitrogen and 10% Phosphoric acid. After testing the soil conditions, a farmer finds that she needs atleast 14 kg of nitrogen and 14 kg of phosphoric acid for her crop. If F_1 costs Rs 6/kg and F_2 costs Rs 5/kg, determine how much of each type of fertiliser should be used so that nutrient requirements are met at a minimum cost. What is the minimum cost?

Solution: The optimization problem can be framed from Table 6.2.10.1 as




Figure 6.2.10.1:

Fertiliser	Nitrogen	Phosphoric Acid
F_1	10%	6%
F_2	5%	10%
Total	14 kg	14 kg

Table 6.2.10.1:

$$Z = \min_{\mathbf{x}} \begin{pmatrix} 6 & 5 \end{pmatrix} \mathbf{x} \quad (6.2.10.1)$$

$$\begin{pmatrix} 2 & 1 \\ 3 & 5 \end{pmatrix} \mathbf{x} \preceq \begin{pmatrix} 280 \\ 700 \end{pmatrix} \quad (6.2.10.2)$$

$$\mathbf{x} \succeq \mathbf{0} \quad (6.2.10.3)$$

yielding

$$Z_{min} = Rs.1000, \mathbf{x} = \begin{pmatrix} 100 \\ 80 \end{pmatrix} \quad (6.2.10.4)$$

6.3. Miscellaneous

6.3.1 A dietician has to develop a special diet using two foods P and Q . Each packet (containing 30 g) of food P contains 12 units of calcium, 4 units of iron, 6 units of cholesterol and 6 units of vitamin A. Each packet of the same quantity of food Q contains 3 units of calcium, 20 units of iron, 4 units of cholesterol and 3 units of vitamin A. The diet requires atleast 240 units of calcium, atleast 460 units of iron and at most 300 units of cholesterol. How many packets of each food should be used to maximise the amount of vitamin A in the diet? What is the maximum amount of vitamin A in the diet?

6.3.2 A farmer mixes two brands P and Q of cattle feed. Brand P , costing Rs 250 per bag,

contains 3 units of nutritional element A, 2.5 units of element B and 2 units of element C. Brand Q costing Rs 200 per bag contains 1.5 units of nutritional element A, 11.25 units of element B, and 3 units of element C. The minimum requirements of nutrients A, B and C are 18 units, 45 units and 24 units respectively. Determine the number of bags of each brand which should be mixed in order to produce a mixture having a minimum cost per bag? What is the minimum cost of the mixture per bag?

6.3.3 A dietician wishes to mix together two kinds of food X and Y in such a way that the mixture contains at least 10 units of vitamin A, 12 units of vitamin B and 8 units of vitamin C. The vitamin contents of one kg food is given below:

Food	Vitamin A	Vitamin B	Vitamin C
X	1	2	3
Y	2	2	1

Table 6.3.3.1:

One kg of food X costs Rs 16 and one kg of food Y costs Rs 20. Find the least cost of the mixture which will produce the required diet?

6.3.4 A manufacturer makes two types of toys A and B. Three machines are needed for this purpose and the time (in minutes) required for each toy on the machines is given below:

Types of Toys	Machines		
	I	II	III
A	12	18	6
B	6	18	9

Table 6.3.4.1:

Each machine is available for a maximum of 6 hours per day. If the profit on each toy of type A is Rs 7.50 and that on each toy of type B is Rs 5, show that 15 toys of type

A and 30 of type B should be manufactured in a day to get maximum profit.

Solution: The given information can be framed as the optimization problem




Figure 6.3.4.1:

$$Z = \max_{\mathbf{x}} \begin{pmatrix} 7.50 & 5 \end{pmatrix} \mathbf{x} \quad (6.3.4.1)$$

$$\begin{pmatrix} 2 & 1 \\ 3 & 0 \\ 2 & 3 \end{pmatrix} \mathbf{x} \preceq \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \quad (6.3.4.2)$$

$$\mathbf{x} \succeq \mathbf{0} \quad (6.3.4.3)$$

Solving the above equations using cvxpy, we obtain

$$Z_{max} = Rs.262.50, \mathbf{x} = \begin{pmatrix} 15 \\ 30 \end{pmatrix} \quad (6.3.4.4)$$

6.3.5 An aeroplane can carry a maximum of 200 passengers. A profit of Rs 1000 is made on each executive class ticket and a profit of Rs 600 is made on each economy class ticket. The airline reserves at least 20 seats for executive class. However, at least 4 times as many passengers prefer to travel by economy class than by the executive class. Determine how many tickets of each type must be sold in order to maximise the profit for the airline. What is the maximum profit?

Solution: Let P be the maximum number of tickets of each type must be sold in order to maximise the profit for the airline . The problem can be formulated as

$$P = \max_{\mathbf{x}} \begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} \quad (6.3.5.1)$$

$$s.t. \quad \begin{pmatrix} -1 & -1 \\ 1 & 0 \\ -4 & 1 \end{pmatrix} \mathbf{x} \succeq \begin{pmatrix} 200 \\ 20 \\ 0 \end{pmatrix} \quad (6.3.5.2)$$

yielding

$$P_{max} = 136000, \mathbf{x} = \begin{pmatrix} 40 \\ 160 \end{pmatrix} \quad (6.3.5.3)$$

6.3.6 Two godowns A and B have grain capacity of 100 quintals and 50 quintals respectively.

They supply to 3 ration shops, D, E and F whose requirements are 60, 50 and 40




Figure 6.3.5.1:

quintals respectively. The cost of transportation per quintal from the godowns to the shops are given in the following table:

How should the supplies be transported in order that the transportation cost is minimum? What is the minimum cost?

Solution: Let's assume that

- (a) A supplies x quintals grain to ration shop D.
- (b) A supplies y quintals grain to ration shop E.
- (c) A will supply remaining grains $100-x-y$ quintals to F.

XUZ		
From/To	A	B
D	6	4
E	3	2
F	2.5	3

Table 6.3.6.1:

- (d) B will supply $60-x$ quintals grain to ration shop D.
- (e) B will supply $50-y$ quintals grain to ration shop E.
- (f) B will supply $x+y-60$ quintals grain to ration shop F.

Total transportation cost is given by :

$$P = 2.5x + 1.5y + 410 \quad (6.3.6.1)$$

Now, Since godown A can supply maximum 60 quintals to ration shop D and 50 quintals to ration shop E and have maximum 100 quintals capacity to supply.

Also, if godown A supplies all 40 quintals to ration shop F, then remaining 60 quintals will be supplied to ration shop D and E and x and y is amount of grains. It can never be negative. This leads to the following conditions

$$x + y \leq 100 \quad (6.3.6.2)$$

$$x \leq 60 \quad (6.3.6.3)$$

$$y \leq 50 \quad (6.3.6.4)$$

$$-x - y \leq -60 \quad (6.3.6.5)$$

$$x \geq 0 \quad (6.3.6.6)$$

$$y \geq 0 \quad (6.3.6.7)$$

The optimization problem can then be expressed as

$$P = \max_{\mathbf{x}} \begin{pmatrix} 2.5 & 1.5 \end{pmatrix} \mathbf{x} + 410 \quad (6.3.6.8)$$

$$s.t. \quad \begin{pmatrix} 1 & 1 \\ -1 & -1 \\ -1 & 0 \\ 0 & -1 \end{pmatrix} \mathbf{x} \preceq \begin{pmatrix} 100 \\ -60 \\ -60 \\ -50 \end{pmatrix} \quad (6.3.6.9)$$

yielding

$$P = 510, \mathbf{x} = \begin{pmatrix} 10 \\ 50 \end{pmatrix} \quad (6.3.6.10)$$

Hence,

- (a) The minimum transportation cost is : 510 /-
- (b) A supplies 10 quintals grain to ration shop D.
- (c) A supplies 50 quintals grain to ration shop E.
- (d) A supplies 40 quintals grain to ration shop F.
- (e) A supplies 50 quintals grain to ration shop D.
- (f) A supplies 0 quintals grain to ration shop E.
- (g) A supplies 0 quintals grain to ration shop F.

6.3.7 An oil company has two depots A and B with capacities of 7000 L and 4000 L respectively. The company is to supply oil to three petrol pumps, D, E and F whose requirements are 4500L, 3000L and 3500L respectively. The distances (in km) between the depots and the petrol pumps is given in the following table:

Distance in (km.)		
From / To	A	B
D	7	3
E	6	4
F	3	2

Table 6.3.7.1:

Assuming that the transportation cost of 10 litres of oil is Re 1 per km, how should the delivery be scheduled in order that the transportation cost is minimum? What is the minimum cost?

6.3.8 A fruit grower can use two types of fertilizer in his garden, brand P and brand Q. The amounts (in kg) of nitrogen, phosphoric acid, potash, and chlorine in a bag of each brand are given in the table. Tests indicate that the garden needs at least 240 kg of phosphoric acid, at least 270 kg of potash and at most 310 kg of chlorine

If the grower wants to minimise the amount of nitrogen added to the garden, how many bags of each brand should be used? What is the minimum amount of nitrogen added in the garden?

kg per bag		
	Brand <i>P</i>	Brand <i>Q</i>
Nitrogen	3	3.5
Phosphoric acid	1	2
Potash	3	1.5
Chlorine	1.5	2

Table 6.3.8.1:

Solution: The given information is summarized in Table 6.3.8.2.

Kg per bag	Brand P	Brand Q
Nitrogen	3	3.5
Phosphoric acid	1	2
Potash	3	1.5
Chlorine	1.5	2

Table 6.3.8.2:




Figure 6.3.8.1:

The given problem can be expressed as

$$P = \min_{\mathbf{x}} \begin{pmatrix} 3 & 3.5 \end{pmatrix} \mathbf{x} \quad (6.3.8.1)$$

$$\begin{pmatrix} 2 & 1 \\ 1 & 2 \\ -1.5 & -2 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 180 \\ 240 \\ -310 \end{pmatrix} \quad (6.3.8.2)$$

yielding

$$P_{min} = 470, \mathbf{x} = \begin{pmatrix} 40 \\ 100 \end{pmatrix} \quad (6.3.8.4)$$

This can be verified using Fig. 6.3.8.1.

6.3.9 Refer to Question 6.3.8.1 If the grower wants to maximise the amount of nitrogen added to the garden, how many bags of each brand should be added? What is the maximum amount of nitrogen added?

6.3.10 A toy company manufactures two types of dolls, A and B. Market research and available resources have indicated that the combined production level should not exceed 1200 dolls per week and the demand for dolls of type B is at most half of that for dolls of type A. Further, the production level of dolls of type A can exceed three times the production of dolls of other type by at most 600 units. If the company makes profit of Rs 12 and Rs 16 per doll respectively on dolls A and B, how many of each should be produced weekly in order to maximise the profit?

Chapter 7

Quadratic Programming

Chapter 8

Semi-Definite Programming

8.1 Find the normal to the curve $2y + x^2 = 3$ passing through (2,2).

Solution: The parameters of the given conic are

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \mathbf{u} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, f = -3 \quad (8.1.1)$$

If \mathbf{x} be the point of contact on the conic, the optimization problem can be formulated as

$$\mathbf{q} = \min_{\mathbf{x}} \|\mathbf{x} - \mathbf{p}\|^2 \quad (8.1.2)$$

$$s.t. \quad \mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} + f = 0 \quad (8.1.3)$$

where

$$\mathbf{p} = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad (8.1.4)$$

Since

$$\|\mathbf{x} - \mathbf{p}\|^2 = \|\mathbf{x}\|^2 - 2\mathbf{p}^\top \mathbf{x} + \|\mathbf{p}\|^2 \quad (8.1.5)$$

$$= \mathbf{y}^\top \mathbf{C} \mathbf{y} \quad (8.1.6)$$

where

$$\mathbf{C} = \begin{pmatrix} \mathbf{I} & -\mathbf{p} \\ -\mathbf{p}^\top & \|\mathbf{p}\|^2 \end{pmatrix} \mathbf{y} = \begin{pmatrix} \mathbf{x} \\ 1 \end{pmatrix} \quad (8.1.7)$$

and (8.1.3) can be expressed as

$$\mathbf{y}^\top \mathbf{A} \mathbf{y} = 0, \quad (8.1.8)$$

where

$$\mathbf{A} = \begin{pmatrix} \mathbf{V} & \mathbf{u} \\ \mathbf{u}^\top & f \end{pmatrix}, \quad (8.1.9)$$

Using SDR (Semi Definite Relaxation), (8.1.2) can be expressed as

$$\min_{\mathbf{X}} tr(\mathbf{C}\mathbf{X}) \quad (8.1.10)$$

$$s.t. \quad tr(\mathbf{A}\mathbf{X}) = 0, \quad (8.1.11)$$

$$\mathbf{X} \succeq \mathbf{0} \quad (8.1.12)$$

yielding

$$\mathbf{x} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}. \quad (8.1.13)$$

Thus, the equation of the normal is

$$\begin{pmatrix} 1 & -1 \end{pmatrix} \mathbf{x} = 0 \quad (8.1.14)$$

8.2 Find the normal to the curve $x^2 = 4y$ passing through (1,2). The parameters of the given conic are

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \mathbf{u} = \begin{pmatrix} 0 \\ -2 \end{pmatrix}, f = 0 \quad (8.2.1)$$

If \mathbf{x} be the point of contact on the conic, the optimization problem can be formulated as

$$\mathbf{q} = \min_{\mathbf{x}} \|\mathbf{x} - \mathbf{p}\|^2 \quad (8.2.2)$$

$$s.t. \quad \mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} + f = 0 \quad (8.2.3)$$

where

$$\mathbf{p} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad (8.2.4)$$

Since

$$\|\mathbf{x} - \mathbf{p}\|^2 = \|\mathbf{x}\|^2 - 2\mathbf{p}^\top \mathbf{x} + \|\mathbf{p}\|^2 \quad (8.2.5)$$

$$= \mathbf{y}^\top \mathbf{C} \mathbf{y} \quad (8.2.6)$$

where

$$\mathbf{C} = \begin{pmatrix} \mathbf{I} & -\mathbf{p} \\ -\mathbf{p}^\top & \|\mathbf{p}\|^2 \end{pmatrix} \mathbf{y} = \begin{pmatrix} \mathbf{x} \\ 1 \end{pmatrix} \quad (8.2.7)$$

and (8.2.3) can be expressed as

$$\mathbf{y}^\top \mathbf{A} \mathbf{y} = 0, \quad (8.2.8)$$

where

$$\mathbf{A} = \begin{pmatrix} \mathbf{V} & \mathbf{u} \\ \mathbf{u}^\top & f \end{pmatrix}, \quad (8.2.9)$$

Using SDR (Semi Definite Relaxation), (8.2.2) can be expressed as

$$\min_{\mathbf{X}} tr(\mathbf{C}\mathbf{X}) \quad (8.2.10)$$

$$s.t. \quad tr(\mathbf{A}\mathbf{X}) = 0, \quad (8.2.11)$$

$$\mathbf{X} \succeq \mathbf{0} \quad (8.2.12)$$

yielding

$$\mathbf{x} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}. \quad (8.2.13)$$

Thus, the equation of the normal is

$$\begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} = 3 \quad (8.2.14)$$

Appendix A

Convex Functions

A.1 A single variable function f is said to be convex if

$$f[\lambda x_1 + (1 - \lambda)x_2] \leq \lambda f(x_1) + (1 - \lambda)f(x_2), \quad (\text{A.1.1})$$

for $0 < \lambda < 1$ and $x_1, x_2 \in \mathbb{R}$.

For a generic quadratic function $ax^2 + bx + c$, let us determine the sufficient condition for it to be convex. Let

$$f(x) = ax^2 + bx + c \quad (\text{A.1.2})$$

Substituting LHS of inequality from (A.1.1) in (A.1.2)

$$\begin{aligned} f[\lambda x_1 + (1 - \lambda)x_2] &= f[x_2 + \lambda(x_1 - x_2)] \\ \implies a[x_2 + \lambda(x_1 - x_2)]^2 &+ b[x_2 + \lambda(x_1 - x_2)] + c \\ \implies ax_2^2 + a\lambda^2 x_1^2 + a\lambda^2 x_2^2 - 2a\lambda^2 x_1 x_2 \\ &+ 2a\lambda x_1 x_2 - 2a\lambda x_2^2 + bx_2 + b\lambda x_1 - b\lambda x_2 + c \quad (\text{A.1.3}) \end{aligned}$$

Substituting RHS of inequality from (A.1.1) in (A.1.2)

$$\begin{aligned}
\lambda f(x_1) + (1 - \lambda) f(x_2) &= a\lambda x_1^2 + b\lambda x_1 + \lambda c \\
&\quad + (1 - \lambda) (ax_2^2 + bx_2 + c) \\
\implies a\lambda x_1^2 + b\lambda x_1 + ax_2^2 + bx_2 + c - a\lambda x_2^2 - b\lambda x_2 &\quad (\text{A.1.4})
\end{aligned}$$

Combining (A.1.3) and (A.1.4) with inequality and simplifying

$$\begin{aligned}
a\lambda^2 x_1^2 + a\lambda^2 x_2^2 - 2a\lambda^2 x_1 x_2 + 2a\lambda x_1 x_2 - 2a\lambda x_2^2 &\leq \\
a\lambda x_1^2 - a\lambda x_2^2 &\quad (\text{A.1.5})
\end{aligned}$$

$$\begin{aligned}
\implies a\lambda^2 x_1^2 + a\lambda^2 x_2^2 - 2a\lambda^2 x_1 x_2 + 2a\lambda x_1 x_2 - a\lambda x_2^2 - a\lambda x_1^2 &\leq 0 \\
\implies x_1^2 (a\lambda^2 - a\lambda) + x_2^2 (a\lambda^2 - a\lambda) - 2x_1 x_2 (a\lambda^2 - a\lambda) &\leq 0 \\
\implies (a\lambda^2 - a\lambda) (x_1 - x_2)^2 &\leq 0 \\
\implies a\lambda(1 - \lambda) (x_1 - x_2)^2 &\geq 0 \quad (\text{A.1.6})
\end{aligned}$$

For the inequality in (A.1.6) to be true,

$$a \geq 0 \because \lambda, 1 - \lambda \geq 0, (x_1 - x_2)^2 \geq 0 \quad (\text{A.1.7})$$

However, $a \neq 0$, since it is a quadratic function. Hence $a > 0$, for $f(x)$ to be convex.

A.2 The quadratic form

$$q(\mathbf{x}) \triangleq \mathbf{x}^\top \mathbf{A} \mathbf{x} + \mathbf{b}^\top \mathbf{x} + c \quad (\text{A.2.1})$$

is convex iff \mathbf{A} is positive semi-definite.

Solution: Consider two points \mathbf{x}_1 and \mathbf{x}_2 , and a real constant $0 \leq \mu \leq 1$. Then,

$$\begin{aligned} & \mu f(\mathbf{x}_1) + (1 - \mu) f(\mathbf{x}_2) - f(\mu \mathbf{x}_1 + (1 - \mu) \mathbf{x}_2) \\ &= (\mu - \mu^2) \mathbf{x}_1^\top \mathbf{A} \mathbf{x}_1 + \left(1 - \mu - (1 - \mu)^2\right) \mathbf{x}_2^\top \mathbf{A} \mathbf{x}_2 \\ & \quad - 2\mu(1 - \mu) \mathbf{x}_1^\top \mathbf{A} \mathbf{x}_2 \end{aligned} \tag{A.2.2}$$

$$= \mu(1 - \mu) \left(\mathbf{x}_1^\top \mathbf{A} \mathbf{x}_1 - 2\mathbf{x}_1^\top \mathbf{A} \mathbf{x}_2 + \mathbf{x}_2^\top \mathbf{A} \mathbf{x}_2 \right) \tag{A.2.3}$$

$$= \mu(1 - \mu) (\mathbf{x}_1 - \mathbf{x}_2)^\top \mathbf{A} (\mathbf{x}_1 - \mathbf{x}_2) \tag{A.2.4}$$

Since \mathbf{x}_1 and \mathbf{x}_2 are arbitrary, it follows from (A.2.4) that

$$\mu f(\mathbf{x}_1) + (1 - \mu) f(\mathbf{x}_2) \geq f(\mu \mathbf{x}_1 + (1 - \mu) \mathbf{x}_2) \tag{A.2.5}$$

iff \mathbf{A} is positive semi-definite, as required.

A.3 Let

$$\mathbf{M} \triangleq \begin{pmatrix} \mathbf{m}_1 & \mathbf{m}_2 \end{pmatrix} \tag{A.3.1}$$

$$\boldsymbol{\lambda} \triangleq \begin{pmatrix} \lambda_1 \\ -\lambda_2 \end{pmatrix} \tag{A.3.2}$$

$$\tag{A.3.3}$$

The function

$$f(\boldsymbol{\lambda}) \triangleq \|\mathbf{M}\boldsymbol{\lambda} - \mathbf{x}\|^2 \tag{A.3.4}$$

is convex.

Solution: (A.3.4) can be expressed as

$$(\mathbf{M}\boldsymbol{\lambda} - \mathbf{x})^\top (\mathbf{M}\boldsymbol{\lambda} - \mathbf{x}) \quad (\text{A.3.5})$$

Consider $\boldsymbol{\lambda}_1$ and $\boldsymbol{\lambda}_2$ and let $0 \leq \mu \leq 1$. Then,

$$f(\mu\boldsymbol{\lambda}_1 + (1 - \mu)\boldsymbol{\lambda}_2) = \|\mathbf{M}(\mu\boldsymbol{\lambda}_1 + (1 - \mu)\boldsymbol{\lambda}_2) - \mathbf{x}\| \quad (\text{A.3.6})$$

$$= \|\mu(\mathbf{M}\boldsymbol{\lambda}_1 - \mathbf{x}) + (1 - \mu)(\mathbf{M}\boldsymbol{\lambda}_2 - \mathbf{x})\| \quad (\text{A.3.7})$$

$$\leq \mu \|\mathbf{M}\boldsymbol{\lambda}_1 - \mathbf{x}\| + (1 - \mu) \|\mathbf{M}\boldsymbol{\lambda}_2 - \mathbf{x}\| \quad (\text{A.3.8})$$

Where (A.3.8) follows from the triangle inequality.

Appendix B

Line

B.1 The shortest distance between the lines whose vector equations are

$$L_1 : \mathbf{x} = \mathbf{x}_1 + \lambda_1 \mathbf{m}_1 \quad (\text{B.1.1})$$

$$L_2 : \mathbf{x} = \mathbf{x}_2 + \lambda_2 \mathbf{m}_2 \quad (\text{B.1.2})$$

is given by

$$\|\mathbf{A} - \mathbf{B}\| = \left\| \frac{\mathbf{m}_1^\top \mathbf{x} \mathbf{m}_1}{\|\mathbf{m}_1\|^2} - \mathbf{x} \right\| \quad (\text{B.1.3})$$

Solution: Let \mathbf{A} and \mathbf{B} be points on lines L_1 and L_2 respectively such that AB is normal to both lines. Define

$$\mathbf{M} \triangleq \begin{pmatrix} \mathbf{m}_1 & \mathbf{m}_2 \end{pmatrix} \quad (\text{B.1.4})$$

$$\boldsymbol{\lambda} \triangleq \begin{pmatrix} \lambda_1 \\ -\lambda_2 \end{pmatrix} \quad (\text{B.1.5})$$

$$\mathbf{x} \triangleq \mathbf{x}_2 - \mathbf{x}_1 \quad (\text{B.1.6})$$

Then, we have the following equations:

$$\mathbf{A} = \mathbf{x}_1 + \lambda_1 \mathbf{m}_1 \quad (\text{B.1.7})$$

$$\mathbf{B} = \mathbf{x}_2 + \lambda_2 \mathbf{m}_2 \quad (\text{B.1.8})$$

From (B.1.7) and (B.1.8), define the real-valued function f as

$$f(\boldsymbol{\lambda}) \triangleq \|\mathbf{A} - \mathbf{B}\| \quad (\text{B.1.9})$$

$$= \|\mathbf{M}\boldsymbol{\lambda} - \mathbf{x}\| \quad (\text{B.1.10})$$

From (A.3.4), since f is convex, differentiating

$$\|\mathbf{M}\boldsymbol{\lambda} - \mathbf{x}\|^2 = \boldsymbol{\lambda}^\top \mathbf{M}^\top \mathbf{M} \boldsymbol{\lambda} - 2\boldsymbol{\lambda}^\top \mathbf{M}^\top \mathbf{x} + \|\mathbf{x}\|^2 \quad (\text{B.1.11})$$

with respect to $\boldsymbol{\lambda}$ and equating to zero,

$$\mathbf{M}^\top (\mathbf{M}\boldsymbol{\lambda} - \mathbf{x}) = \mathbf{0} \quad (\text{B.1.12})$$

yielding

$$\mathbf{M}^\top \mathbf{M} \boldsymbol{\lambda} = \mathbf{M}^\top \mathbf{x} \quad (\text{B.1.13})$$

Appendix C

Manual

