

Quadratic Programming

1 12th Maths - Chapter 6

This is Problem-23 from Exercise 6.6

1. Find the equation of the normal to the curve $x^2 = 4y$ and passing through the point $(1, 2)$.

Solution: The given equation of the curve can be written as

$$g(\mathbf{x}) = \mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (1)$$

where

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (2)$$

$$\mathbf{u} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \quad (3)$$

$$f = 0 \quad (4)$$

We are given that

$$\mathbf{h} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad (5)$$

This can be formulated as optimization problem as below:

$$\min_{\mathbf{x}} \quad f(\mathbf{x}) = \|\mathbf{x} - \mathbf{h}\|^2 \quad (6)$$

$$\text{s.t.} \quad g(\mathbf{x}) = \mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (7)$$

It is already proved that the optimization problem is nonconvex. The constraints throw an error when *cvxpy* is used.

We will use Lagrange multipliers method to find the optimum value.
Define

$$H(\mathbf{x}, \lambda) = f(\mathbf{x}) - \lambda g(\mathbf{x}) \quad (8)$$

and we find that

$$\nabla f(\mathbf{x}) = 2(\mathbf{x} - \mathbf{h}) \quad (9)$$

$$\nabla g(\mathbf{x}) = 2(\mathbf{V}\mathbf{x} + \mathbf{u}) \quad (10)$$

We have to find $\lambda \in \mathbb{R}$ such that

$$\nabla H(\mathbf{x}, \lambda) = 0 \quad (11)$$

$$\implies 2(\mathbf{x} - \mathbf{h}) - 2\lambda(\mathbf{V}\mathbf{x} + \mathbf{u}) = 0 \quad (12)$$

$$\implies \mathbf{x} - \mathbf{h} = \lambda(\mathbf{V}\mathbf{x} + \mathbf{u}) \quad (13)$$

$$\implies (\mathbf{I} - \lambda\mathbf{V})\mathbf{x} = \lambda\mathbf{u} + \mathbf{h} \quad (14)$$

$$\implies \begin{pmatrix} 1-\lambda & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} = \lambda \begin{pmatrix} 0 \\ -2 \end{pmatrix} + \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad (15)$$

$$\implies \begin{pmatrix} 1-\lambda & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 1 \\ -2\lambda + 2 \end{pmatrix} \quad (16)$$

We have 2 cases to consider here.

(a) When $\lambda \neq 1$. Writing augmented matrix,

$$\left(\begin{array}{ccc|c} 1-\lambda & 0 & 1 & 1 \\ 0 & 1 & -2\lambda+2 & -2\lambda+2 \end{array} \right) \xleftarrow{R_1 \leftarrow \frac{R_1}{1-\lambda}} \left(\begin{array}{ccc|c} 1 & 0 & \frac{1}{1-\lambda} & \frac{1}{1-\lambda} \\ 0 & 1 & -2\lambda+2 & -2\lambda+2 \end{array} \right) \quad (17)$$

Then, we get

$$\mathbf{x}_m = \begin{pmatrix} \frac{1}{1-\lambda} \\ -2\lambda+2 \end{pmatrix} \quad (18)$$

Substituting this value in (7)

$$\begin{aligned}
& \left(\frac{1}{1-\lambda} \quad -2\lambda + 2 \right) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{1-\lambda} \\ -2\lambda + 2 \end{pmatrix} \\
& \quad + 2 \begin{pmatrix} 0 & -2 \end{pmatrix} \begin{pmatrix} \frac{1}{1-\lambda} \end{pmatrix} = 0 \\
& \implies \left(\frac{1}{1-\lambda} \quad 0 \right) \begin{pmatrix} \frac{1}{1-\lambda} \\ -2\lambda + 2 \end{pmatrix} - 4(-2\lambda + 2) = 0 \\
& \implies 8(\lambda^3 - 3\lambda^2 + 3\lambda - 1) + 1 = 0 \\
& \implies (\lambda^3 - 3\lambda^2 + 3\lambda - 1) = -\frac{1}{8} \\
& \implies (\lambda - 1)^3 = -\frac{1}{8} \\
& \implies \lambda - 1 = -\frac{1}{2} \\
& \implies \lambda = \frac{1}{2} \quad (19)
\end{aligned}$$

Substituting the value of λ in (18)

$$\mathbf{x}_m = \mathbf{q} = \begin{pmatrix} \frac{1}{1-\frac{1}{2}} \\ -2\left(\frac{1}{2}\right) + 2 \end{pmatrix} \quad (20)$$

$$= \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad (21)$$

(b) When $\lambda = 1$.

$$(16) \implies \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (22)$$

This is an invalid solution.

Given the point of contact \mathbf{q} , the equation to the normal is given by

$$(\mathbf{V}\mathbf{q} + \mathbf{u})^\top \mathbf{R}(\mathbf{x} - \mathbf{q}) = 0 \quad (23)$$

$$\Rightarrow \left(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} + \begin{pmatrix} 0 \\ -2 \end{pmatrix} \right)^\top \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \left(\mathbf{x} - \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right) = 0 \quad (24)$$

$$\Rightarrow (2 \quad -2) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \left(\mathbf{x} - \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right) = 0 \quad (25)$$

$$\Rightarrow (2 \quad 2) \left(\mathbf{x} - \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right) = 0 \quad (26)$$

$$\Rightarrow (1 \quad 1) \mathbf{x} = 3 \quad (27)$$

The relevant figure is shown in 1




Figure 1