

The Neurophysiology of Cybersickness: A Systematic Review of Neuromarkers, Neurostimulation, and Reporting Practices in Virtual Reality

Gaël Van der Lee, François Cabestaing, Hakim Si-Mohammed

Abstract—Visually induced motion sickness (VIMS) is a prevalent adverse response to virtual reality (VR) exposure, and its assessment has relied primarily on retrospective subjective questionnaires that cannot support continuous monitoring or real-time detection. This PRISMA-guided systematic review synthesizes 92 studies on the neurophysiology of VIMS in VR, retrieved from PubMed, Web of Science, and Scopus through March 2025, spanning electroencephalography, functional near-infrared spectroscopy, and non-invasive neurostimulation.

The EEG literature converges on increased low-frequency (delta and theta) power as the most consistent spectral correlate of VIMS. Alpha-band findings have been inconsistent across prior work; meta-analytic comparison of hardware platforms reported here attributes a substantial portion of that inconsistency to VR display type, with head-mounted-display and screen-based studies differing systematically in the direction of alpha and beta responses. Beyond spectral power, the corpus documents event-related potential evidence of attentional and inhibitory-control costs, fNIRS hemodynamic correlates, resting-state EEG predictors of individual susceptibility, and early closed-loop mitigation systems driven by neurophysiological detection.

A systematic examination of the classification and prediction literature reveals recurrent reporting gaps (undisclosed decision thresholds, absent discriminative metrics, and validation schemes that place the same participant’s data in both training and test sets) that limit the interpretability of headline performance figures. Together with hardware-related heterogeneity, baseline-selection variability, and terminological inconsistency, these gaps constrain cross-study synthesis. The causal status of observed neural associations, cross-subject generalization of detection models, and the neurodynamics of adaptation and recovery remain open empirical questions.

Index Terms—Cybersickness, EEG, Human-centered computing, Human computer interaction (HCI), Interaction paradigms, Virtual reality.

I. INTRODUCTION

Virtual Reality (VR) technology has permeated numerous sectors beyond entertainment, becoming an integral tool in fields such as healthcare, professional training, and education [1]. VR has been characterized as “the illusion of participation in a synthetic environment rather than external observation of such an environment,” [2] relying on stereoscopic head-tracked displays, hand/body tracking, and binaural sound to constitute an immersive, multisensory experience, though no single definition has been universally adopted [3].

A barrier to widespread VR adoption is the onset of cybersickness (CS) [4]. Cybersickness is a form of

visually induced motion sickness (VIMS) [5] characterized by symptoms such as nausea, dizziness, and disorientation that arise during or after immersion in a virtual environment [4]. The relationship between cybersickness, VIMS, simulator sickness, and motion sickness is treated in detail in Section II; the present review uses VIMS as the umbrella term when introducing constructs and uses the term each cited study employs when reporting its findings. VIMS affects a substantial portion of the user population, with studies reporting prevalence between 60% and 95% and premature session termination in 6–12.9% of participants [4]; it also degrades user experience and impairs task performance [6]. Assessment of VIMS has historically relied on subjective post-hoc questionnaires such as the Simulator Sickness Questionnaire [7]. Such instruments are retrospective, prone to recall bias, and do not capture the continuous nature of the experience [8], [9]. This limitation creates a need for objective measures grounded in the user’s physiology and for interventions that can modulate the user’s state during exposure rather than after the fact.

Two converging lines of research have responded by examining the brain. The first records neural activity, primarily electroencephalography (EEG) and functional near-infrared spectroscopy (fNIRS), to identify signals that index the onset, severity, or trajectory of VIMS. The second applies non-invasive brain stimulation, including galvanic vestibular stimulation (GVS), transcranial direct-current stimulation (tDCS), and transcranial alternating-current stimulation (tACS), to modulate the underlying processes that produce sickness. We use the term *neuromarkers* to refer to signals derived from non-invasive recording of brain activity in this context. Non-invasive brain stimulation is explored as a complementary intervention approach. In a VR context, neuromarkers can carry information that subjective questionnaires cannot: continuous, time-resolved indices of user state that can in principle drive adaptive content, comfort thresholds, and closed-loop mitigation [10], [11].

In a prior review, Chang et al. [8] synthesized the EEG literature on cybersickness and characterized the EEG analysis pipeline used in that research. The present systematic review differs in four respects. First, the scope extends beyond EEG to include functional near-infrared spectroscopy as an additional neuromarker modality and non-invasive neurostimulation as a complementary intervention approach.

Second, the corpus is larger and more recent: 92 studies retrieved through March 2025, compared with 33 EEG studies published before September 2021. Third, beyond characterizing the analysis pipeline, this review applies meta-analytic tests for confounders across the corpus, identifying VR display hardware as a source of the inconsistent alpha-band findings reported in prior work. Fourth, this review situates the empirical findings within the broader context of the field: the theoretical frameworks invoked to explain VIMS, the terminology used across the cybersickness, VIMS, simulator sickness, and motion sickness literatures, and a critical appraisal of recurring methodological shortcomings.

A. Contributions

The contributions of this review are threefold. First, we provide a systematic synthesis of 92 VR studies spanning non-invasive neuroimaging (EEG and fNIRS) and non-invasive neurostimulation, covering spectral correlates, event-related potential evidence, hemodynamic findings, individual susceptibility, and early closed-loop mitigation systems within a unified neuromarker frame. Second, we report meta-analytic evidence that VR display hardware modulates the EEG band-power responses associated with VIMS, attributing a substantial portion of the alpha-band inconsistency reported in the prior literature to hardware differences rather than to differences in VIMS per se. Third, we systematically examine the methodological practices of the classification and prediction literature and the broader corpus, characterizing recurrent reporting gaps, hardware and protocol heterogeneity, and baseline-selection practices that constrain cross-study synthesis and the assessment of generalization claims.

B. Paper Organization

The remainder of this paper is organized as follows. Section II reviews the relevant theoretical frameworks for VIMS and the principal neuroimaging and stimulation modalities. Section III describes the review methodology and search strategy. Section IV presents the corpus, the per-modality synthesis, and the corpus-level meta-analytic findings. Section V discusses the methodological challenges identified in this review and directions for future work. Section VI concludes.

II. THEORETICAL AND METHODOLOGICAL BACKGROUND

A. Terminology and Etiological Frameworks

Before surveying the theories proposed to explain cybersickness, it is useful to delimit the construct itself and to distinguish it from related forms of sickness and from perceptual phenomena that often co-occur with it.

Motion sickness is the broadest construct, encompassing adverse responses to both physical and visually specified motion [12], [13]. Within it, visually induced motion sickness (VIMS) designates sickness elicited primarily by visual motion, including cases where physical body motion is absent or inconsistent with the visual scene [14], [15]. Cybersickness is most often treated as the VR-specific instance of VIMS;

simulator sickness, gaming sickness, and related labels denote overlapping exposure contexts rather than distinct symptom categories [16]–[18]. This review adopts the above hierarchy when introducing constructs, but reports each included study using the terminology its authors chose, in order to avoid imposing a uniform vocabulary on a literature that does not yet share one.

Traditionally, these constructs have been measured using subjective questionnaires. The dominant instrument is the Simulator Sickness Questionnaire (SSQ), which yields nausea, oculomotor, and disorientation subscales [7]; it is frequently substituted by the single-item Fast Motion Sickness Scale (FMS) for fast, continuous, in-task ratings [19]. Broader diagnostic criteria for motion sickness and VIMS extend these subscales to gastrointestinal, thermoregulatory, arousal, dizziness or vertigo, headache, and visual or oculomotor domains [15].

Several etiological frameworks have been proposed to explain cybersickness and the wider VIMS phenomenon; they are not mutually exclusive and often emphasize complementary mechanisms.

The most commonly used framework is the *sensory conflict theory*, which attributes sickness to contradictory visual, vestibular, and proprioceptive signals, or between current sensory input and the prior exposure history encoded by the observer [20]–[22]. Two refinements within this family further specify the nature of the conflict: *subjective vertical mismatch* frames it as a discrepancy between the sensed and expected gravitational vertical direction [23], while *rest-frame conflict* locates it in the observer’s difficulty identifying a stable spatial reference [24]. This family is particularly pertinent to VR, where HMDs routinely present visual self-motion while vestibular and proprioceptive signals indicate little or no corresponding physical movement.

A second framework, *postural instability theory*, locates the cause of sickness not in sensory mismatch but in a sustained inability to maintain postural control under unfamiliar sensorimotor demands [25], [26]. The theory is most directly testable in VR paradigms involving standing, walking, or large-field optic flow, where body sway and control dynamics can be measured objectively and have been reported to precede or accompany subjective sickness [27]. A related strand of recent VR work returns to the role of visually induced self-motion itself, showing that sickness intensifies when it violates the user’s expected motion or control state [28], [29].

A further set of proposals addresses narrower mechanisms or different levels of explanation. *Vergence-accommodation conflict* identifies a mismatch between binocular convergence and the fixed focal distance of stereoscopic displays as a source of oculomotor discomfort [30], while *oculomotor theories* more broadly relate sickness to specific eye-movement patterns, such as nystagmus or ocular torsion, evoked by the visual stimulus [31]. At an evolutionary level, Treisman’s *toxin-defense hypothesis* explains why disturbed sensory integration should give rise to nausea and emesis, but does not specify the proximate sensory conditions that trigger them [32]. Finally, *predictive-coding formulations* recast sensory mismatch as a persistent prediction error between expected

and observed input, offering a unifying cognitive framework for the preceding theories [33].

B. Neural Recording and Modulation Modalities

While VIMS has historically been assessed through subjective reports, a growing body of work uses neural recordings, and more recently non-invasive brain stimulation, to characterize the user's state during VR exposure or to actively modulate it.

Electroencephalography (EEG) is the most widely used recording modality in this literature. Its high temporal resolution supports the study of spectral power, event-related potentials, and connectivity dynamics throughout the development of sickness, and its portability makes it compatible with HMD-based and other interactive VR paradigms [27], [34]. These strengths are offset by noise, limited spatial specificity, particularly for deep vestibular structures, and by methodological sensitivities, such as motion artifacts, interference, reference choice, and baseline selection, which constrain the interpretation of reported effects.

Functional near-infrared spectroscopy (fNIRS) is an emerging optical technique that measures cortical hemodynamic changes via near-infrared light [35], [36]. It offers an intermediate trade-off between temporal and spatial resolution and tolerates moderate movement, making it well suited to VR paradigms such as HMD-based roller-coaster exposure, VR locomotion, or stereoscopic viewing [11], [37]–[39]. Because it relies on hemodynamic rather than electrical signals, its temporal resolution is substantially lower than that of EEG, and its optical penetration restricts measurement to superficial cortical regions.

Beyond recording, non-invasive neurostimulation has begun to be used to modulate sickness-relevant processes during VR or visually induced motion exposure. Reported approaches include galvanic vestibular stimulation, transcranial direct-current stimulation, transcranial alternating-current stimulation, and related techniques [40]. By directly acting on vestibular or cortical activity, these methods provide a causal complement to the correlational evidence offered by EEG and fNIRS.

III. REVIEW METHODOLOGY

A. Search Strategy

A systematic literature search across three databases, PubMed, Web of Science, and Scopus, retrieved records published through the 27th of March 2025. No earlier date restriction was imposed, so the search captures the full historical record of the field up to that date. The query targeted the intersection of three concept blocks: brain-directed measurement or modulation methods, cybersickness and related sickness constructs, and immersive virtual reality. The full query was:

```
((("Brain Computer Interface" OR BCI OR EEG OR
electroencephalogram OR ERP OR MEG OR
magnetoencephalography OR fMRI OR ECoG OR
fNIRS OR PET OR "positron emission
tomography" OR neurostimulation OR "brain
```




Fig. 1: PRISMA flow diagram for the present review.

```
stimulation" OR "non-invasive brain
stimulation" OR tACS OR "transcranial
alternating current stimulation" OR tDCS
OR "transcranial direct current
stimulation" OR TMS OR "transcranial
magnetic stimulation" OR "galvanic
vestibular stimulation" OR GVS)
AND (Cybersickness OR "simulator sickness" OR
"visually induced motion sickness" OR VIMS
OR "motion sickness")
AND ("virtual reality" OR VR OR "head-mounted
display" OR HMD OR "immersive environment
") )
```

B. Study Selection and Eligibility Criteria

Records identified by the search were screened in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 statement; the full selection process is summarized in the PRISMA flow diagram (Figure 1). The combined search returned 443 records: 57 from PubMed, 147 from Web of Science, and 239 from Scopus. Duplicate removal across the three sources reduced this set to 219 unique records.

Each of the 219 unique records underwent manual title-and-abstract screening against the following eligibility criteria. A record was included if it met all of the following:

- The study employed a neuroimaging modality (EEG, ERP, fNIRS, fMRI, MEG, ECoG, PET) or a brain-stimulation modality (tDCS, tACS, TMS, GVS) to measure or modulate brain activity.
- The study addressed VIMS or a closely related construct (simulator sickness, or motion sickness in a virtual environment) within a virtual reality or otherwise immersive context.

- The study reported original empirical results from human participants, including reanalyses of previously released public datasets.

A record was excluded if it met any of the following:

- The work was non-empirical (literature reviews, meta-analyses, editorials, letters, or conference abstracts without an accompanying full paper).
- The study did not address VIMS or a related sickness construct as a primary outcome.
- The full text could not be retrieved.

Title-and-abstract screening excluded 119 records, leaving 100 reports sought for full-text retrieval. Of these, 8 could not be obtained as full text and were excluded at the retrieval stage. The remaining 92 articles comprise the final corpus analyzed in this review. PET returned no eligible studies; MEG returned a single record, which was excluded at the title-and-abstract stage as not addressing VIMS or a related sickness construct; fMRI returned no records at all. The absence of these modalities from the final corpus therefore reflects the current literature rather than a scope restriction imposed by the search.

C. Data Extraction and Thematic Synthesis

For each of the 92 included articles, a structured data extraction process was performed. This process was guided by a predefined protocol designed to ensure consistency and to facilitate subsequent comparison and synthesis. The following data points were extracted from each paper:

- 1) **Summary of Contribution:** A concise summary of the study's primary objective and its main contribution.
- 2) **Sickness Measurement:** The methods used to measure or label VIMS, distinguishing between subjective questionnaires (e.g., SSQ, FMS, VRSQ) and objective physiological or behavioral metrics (e.g., galvanic skin response, postural sway).
- 3) **Methodological Pipeline:** A breakdown of the neurophysiological pipeline, including: (a) the measurement or stimulation modality, (b) the key feature extraction techniques (e.g., power spectral density, event-related potentials, functional connectivity), and (c) the primary analysis or classification methods (e.g., statistical tests, machine-learning models).
- 4) **Theoretical Framework:** The explicitly stated or implicitly assumed theory of VIMS (e.g., sensory conflict, postural instability) that guided the study's design or interpretation.
- 5) **Quantitative Performance:** Reported performance measures, such as classification accuracy, correlation coefficients, root-mean-square error, or statistical significance values.
- 6) **Empirical Study Details:** For empirical studies, specific details of the experimental protocol, including the VR environment, display hardware, sickness-inducing stimulus, and number of participants.
- 7) **Reported Shortcomings:** Limitations, caveats, or weaknesses identified by the authors or apparent from the study's design.

Following extraction, each paper was assigned to a primary research-objective category to support quantitative synthesis. An initial set of categories was defined *a priori* based on established research paradigms and refined iteratively during extraction. The final categories, which structure the Results section, are: Classification (Binary, Likert, Continuous), Correlates (Brain Regions, Rhythms), Mitigation (A/B Testing, Real-time, Neurostimulation), User Differences, Dataset Provision, Recovery, and studies on the Effects of VIMS on other cognitive or performance metrics.

IV. RESULTS

The 92 corpus studies address four main research objectives: classification or prediction of VIMS from physiological signals (38 studies), identification of neural and physiological correlates (25), development of mitigation strategies with a closed-loop or brain-state-interventional design that engages neurophysiological measurement (8), and characterization of individual susceptibility (3). Eighteen residual studies (further mitigation work without a closed-loop or brain-state-interventional component, cognitive effects, recovery, dataset contributions, and other framings) are integrated into the relevant subsections, with full discussion deferred to Section V. This section adds two cross-cutting layers on top of these objectives: Section IV-A describes the corpus as a whole, and Section IV-F reports meta-analytic observations across the 92 studies.

Activity in the field has accelerated in recent years: 63.0% of the corpus (58 of 92 studies) was published from 2021 onward, with a corresponding shift toward classification, mitigation, and susceptibility objectives (Fig. 2). Per-study sample sizes have also increased over the same period (Mann-Whitney U on studies published before 2021 versus from 2021 onward, Cliff's $\delta = -0.337$, $p = 0.007$, medium effect).

Table I lists all 92 studies with their modality, sample size, virtual reality hardware, stimulus paradigm, VIMS measure, primary objective, and topic.

A. Corpus Characteristics

The 92 included studies cluster on EEG-based investigations of young adults in head-mounted VR, but vary widely on baseline condition, montage, and preprocessing. The dispersion documented in this subsection is the precondition for the meta-analytic observations reported in Section IV-F.

1) *Neuroimaging and Physiological Recording Modalities:* EEG is the predominant modality (77 of 92 studies, 83.7%), followed by fNIRS (4 of 92, 4.3%). The remaining 11 studies (12.0%) record neither EEG nor fNIRS: 9 deliver non-invasive stimulation without concurrent brain recording (GVS in 5, tDCS in 2, tACS in 1, bone-conducted vibration in 1), and 2 record only peripheral physiology. Across the full corpus, including studies that also record EEG, non-invasive stimulation is used in 11 studies (GVS in 5, tDCS in 3, tACS in 2, bone-conducted vibration in 1). No corpus study uses fMRI, MEG, or PET; Section III states the search-query and screening implications of this absence.

TABLE I: All 92 studies included in this systematic review. Modality: EEG, fNIRS = neuroimaging; GVS = galvanic vestibular stimulation; tDCS/tACS = transcranial direct/alternating current stimulation; ECG, GSR, EOG, EMG, EGG = peripheral physiology; posturography = balance / center-of-pressure recording. CS measure abbreviations: SSQ = Simulator Sickness Questionnaire; VRSQ = VR Sickness Questionnaire; FMS = Fast Motion-Sickness Scale; FMS+SSQ = repeated FMS sampling plus SSQ assessment; MSSQ = Motion Sickness Susceptibility Questionnaire; cont. = continuous self-report; Likert = single-item Likert; custom = study-specific scale; behav. = behavioral measure only; none = no cybersickness measure reported; studies listed as cont., Likert, or behav. without a named questionnaire did not report a specific instrument. Sample size N gives subjects analyzed. Topic gives a short noun phrase describing the focus of each paper.

Ref.	Modality	N	VR device	Stimulus	CS measure	Objective	Topic
[41]	EEG	2	HMD	Mixed	none	Mitigation	EEG-informed scene-simplification mitigation
[42]	EEG+EOG	17	HMD	Driving (passive)	SSQ	Correlates	Alpha connectivity and P3 in susceptibility
[43]	EEG+EOG	33	Screen	Passive viewing	FMS	Correlates	EEG time-frequency correlates of VIMS severity
[44]	EEG	31	HMD	Mixed	SSQ	Correlates	Spectral EEG correlates of VIMS in HMD
[45]	EEG+ECG+GSR+EOG	27	HMD	Active navigation	custom	Classification	Spiking-network EEG+HRV cybersickness prediction
[40]	tACS+GSR	37	HMD	Passive viewing	cont.	Mitigation	10 Hz tACS over vestibular cortex
[46]	EEG	68	HMD	Mixed	SSQ	Other	Sex differences in EEG neurofeedback sessions
[47]	EEG+tDCS+ECG+EOG	41	HMD	Passive viewing	SSQ	Mitigation	Placebo tDCS in EEG neurofeedback for cybersickness
[48]	EEG+ECG	39	HMD	Active navigation	SSQ	Cognition	VR-parameter manipulation during EEG neurofeedback
[49]	EEG	15	HMD	Passive viewing	SSQ	Classification	Resting-state EEG entropy classification
[50]	EEG	12	Screen	Passive viewing	SSQ	Mitigation	Rest-frame mitigation with EEG correlates
[51]	EEG+ECG+EOG	20	Screen	Passive viewing	SSQ	Correlates	Heartbeat-evoked potentials as cybersickness correlates
[52]	EEG	19	Projector	Driving (passive)	cont.	Correlates	ICA-based motion sickness EEG component clusters
[53]	EEG	19	Projector	Driving (passive)	cont.	Correlates	Continuous motion sickness EEG dynamics
[54]	EEG	20	HMD	Other	SSQ	Correlates	blur-induced visual discomfort in HMDs
[27]	EEG	16	HMD	Active navigation	SSQ	Correlates	EEG during VR sickness while walking
[55]	EEG	32	HMD	Mixed	SSQ	Dataset	EEG cybersickness dataset and benchmark
[56]	EEG	32	HMD	Active navigation	SSQ	Classification	EEG-based comparison of VR locomotion techniques
[9]	EEG+ECG+EOG+EGG+ posturography	20	HMD	Active navigation	Likert	Classification	Multimodal EEG+physiology severity classification
[57]	EEG+ECG+GSR	25	Other	Passive viewing	SSQ	Classification	Convolutional-transformer EEG cybersickness detection
[58]	EEG	23	HMD	Passive viewing	SSQ	Classification	Multi-domain EEG feature comparison
[59]	EMG	24	HMD	Passive viewing	MSSQ	Correlates	VEMP changes after VR vection exposure
[60]	GVS	47	HMD	Passive viewing	SSQ	Mitigation	Omnidirectional GVS for 360° video viewing
[61]	EEG+EOG	23	HMD	Passive viewing	SSQ	Classification	LSTM regression of continuous VRMS severity
[62]	EEG	15	HMD	Mixed	SSQ	Classification	Multi-scale EEG brain-network classification
[63]	EEG	15	HMD	Passive viewing	SSQ	Classification	Bilateral asymmetry features for VRMS detection
[64]	EEG+EOG	22	HMD	Passive viewing	SSQ	Classification	Siam-LSTM with EEG functional brain networks
[65]	EEG+posturography	22	HMD	Active navigation	VRSQ	Classification	Dual-pathway raw EEG and network model
[66]	EEG	40	Screen	Passive viewing	SSQ	Indiv. diff.	EEG band-power correlates of cybersickness susceptibility
[67]	EEG	24	HMD	Passive viewing	cont.	Classification	Deep-learning EEG classification of cybersickness
[68]	EEG	9	Screen	Active navigation	SSQ	Other	Gamma-band EEG visualization of sickness onset
[69]	EEG+ECG+GSR+EOG	45	CAVE	Active navigation	SSQ	Correlates	EEG and autonomic correlates during navigation
[70]	EEG+ECG+GSR+EOG+ EGG	57	Projector	Active navigation	SSQ	Correlates	Multimodal physiology of navigation cybersickness
[71]	EEG	202	HMD	Passive viewing	Likert	Classification	EEG-to-video transfer for cybersickness prediction
[72]	EEG+EOG	30	HMD	Passive viewing	SSQ	Correlates	Source-localized alpha cybersickness correlates after VR
[73]	EEG	16	Projector	Driving (passive)	cont.	Classification	ICA-EEG regression of motion-sickness level
[74]	EEG	6	CAVE	Driving (passive)	cont.	Classification	Genetic feature selection for EEG classification
[34]	EEG	43	HMD	Passive viewing	cont.	Correlates	EEG band-power correlates with continuous joystick cybersickness
[75]	EEG+ECG+GSR	40	HMD; Screen	Passive viewing	SSQ	Classification	EEG and physiology fusion for SSQ estimation
[76]	EEG+EOG+posturography	20	Projector	Driving (passive)	custom	Classification	Multimodal ensemble for VIMS severity
[77]	EEG	18	HMD	Passive viewing	behav.	Classification	Wavelet-packet EEG features for VRMS
[78]	tDCS	20	HMD	Passive viewing	cont.	Mitigation	Cathodal parietal tDCS for cybersickness
[79]	EEG+EOG	22	HMD	Passive viewing	FMS	Classification	EEG vs. autonomic features for VR sickness

Continued on next page

Table 1 continued from previous page

Ref.	Modality	N	VR device	Stimulus	CS measure	Objective	Topic
[80]	EEG+EOG	20	HMD	Passive viewing	FMS	Indiv. diff.	EEG biomarker of VRMS resistance
[81]	EEG	39	HMD	Mixed	FMS	Classification	Resting EEG prediction of in-car VRMS
[82]	EEG+tACS	18	HMD	Mixed	FMS	Mitigation	Online 18 Hz tACS over P3
[83]	EEG	130	HMD	Passive viewing	custom	Classification	LSTM prediction of impending motion sickness
[84]	EEG	16	TV	Passive viewing	SSQ	Other	Test-retest reliability of EEG sickness markers
[85]	EEG	9	Projector	Driving (passive)	custom	Correlates	EEG dynamics during VR motion sickness
[86]	EEG	6	CAVE	Driving (passive)	cont.	Classification	BCI estimation of motion-sickness level
[87]	EEG+ECG	25	HMD	Passive viewing	cont.	Classification	Real-time EEG+ECG cybersickness warning
[88]	EEG	9	CAVE	Driving (active)	cont.	Correlates	Wearable EEG and cardiovascular correlates
[89]	EEG	8	Screen	Driving (active)	cont.	Correlates	Low-channel EEG markers for VIMS
[90]	EEG	36	HMD	Passive viewing	SSQ	Classification	CNN-ECA-LSTM continuous cybersickness regression
[91]	EEG	36	HMD	Passive viewing	SSQ	Classification	EEG spectral and network regression of severity
[92]	EEG	25	HMD	Passive viewing	SSQ	Classification	Wavelet-packet GRU for VR motion sickness
[93]	EEG	9	Screen	Active navigation	SSQ	Classification	Time-domain EEG features for cybersickness
[6]	EEG+EOG	29	HMD	Passive viewing	SSQ	Cognition	P3b ERP attention markers during VR sickness
[94]	EEG+ECG+EOG	8	HMD	Active navigation	custom	Classification	Multimodal symptom-wise cybersickness markers
[95]	EEG	18	HMD	Passive viewing	cont.	Correlates	EEG correlates of display-parameter cybersickness
[96]	EEG	40	Screen	Passive viewing	SSQ	Correlates	EEG microstates and functional connectivity
[97]	ECG+GSR	13	HMD	Mixed	VRSQ	Classification	Symbolic ML on peripheral biosignals
[33]	EEG	14	HMD	Passive viewing	SSQ	Correlates	VIMS effective connectivity and predictive coding
[98]	EEG	33	HMD	Passive viewing	SSQ	Correlates	EEG across speed, complexity, and stereoscopy
[99]	EEG	9	Screen	Active navigation	SSQ	Classification	Rule-based EEG severity classification
[100]	GVS+EGG	19	HMD	Active navigation	SSQ	Mitigation	Simulator-sickness GVS mitigation during flight VR
[101]	GVS+balance	140	HMD	Passive viewing	custom	Correlates	GVS probing of vestibular contribution
[102]	EEG+EMG	28	HMD	Passive viewing	MSSQ	Classification	BioVRSea EEG+EMG motion sickness prediction
[103]	EEG+EOG	20	HMD	Mixed	FMS	Classification	EEG preprocessing trade-offs for cybersickness
[104]	EEG+GSR	22	HMD	Passive viewing	SSQ	Classification	Explainable AI on multimodal physiology
[105]	EEG	25	HMD	Passive viewing	SSQ	Classification	PLV connectivity CNN-LSTM for VIMS
[106]	GVS+GSR	20	HMD	Passive viewing	SSQ	Mitigation	Wearable motion-synchronized GVS
[107]	EEG	22	HMD; Screen	Passive viewing	SSQ	Correlates	EEG comparison of HMD versus screen VR
[108]	EEG	31	HMD	Passive viewing	SSQ	Correlates	EEG across VR motion parameter variations
[109]	tDCS+ECG+posturography	20	HMD	Passive viewing	SSQ	Mitigation	Anodal tDCS over right TPJ
[39]	fNIRS+ECG+posturography	11	HMD	Passive viewing	behav.	Correlates	fNIRS and posturography for 3D sickness
[110]	EEG+ECG+EGG	28	HMD	Mixed	FMS+SSQ	Indiv. diff.	EEG aperiodic activity for susceptibility profiling
[10]	EEG	53	HMD	Passive viewing	SSQ	Mitigation	Real-time EEG-driven factor-specific mitigation
[111]	EEG	6	HMD	Active navigation	SSQ	Mitigation	Gaze-contingent peripheral blur mitigation
[112]	Other	108	Projector	Mixed	SSQ	Mitigation	Bone-conducted vibration vestibular stimulation
[113]	GVS	40	HMD	Active navigation	FMS+SSQ	Mitigation	Noisy GVS for nauseogenic VR
[114]	EEG	6	Projector	Driving (passive)	cont.	Classification	Continuous EEG-based motion-sickness estimation
[115]	EEG	40	TV	Passive viewing	SSQ	Recovery	EEG recovery dynamics after VR exposure
[116]	EEG	17	HMD	Active navigation	SSQ	Cognition	Inhibition-related ERPs after VIMS
[117]	EEG	12	HMD	Other	SSQ	Cognition	Full-body exergame, simulator sickness, EEG theta
[118]	EEG	16	HMD	Passive viewing	SSQ	Classification	SE-CNN regression of VR motion sickness from EEG
[38]	fNIRS	1	HMD	Active navigation	SSQ	Mitigation	fNIRS-triggered FOV-narrowing mitigation
[11]	fNIRS	10	HMD	Mixed	SSQ	Mitigation	Wearable fNIRS for adaptive FOV restriction
[119]	EEG+ECG	64	HMD	Passive viewing	SSQ	Classification	Interpretable spiking-network EEG prediction
[120]	EEG	25	HMD	Mixed	FMS+SSQ	Mitigation	Multisensory VIMS mitigation with EEG correlates
[37]	fNIRS	30	HMD	Passive viewing	SSQ	Correlates	fNIRS angular-gyrus correlates of cybersickness
[121]	EEG+EOG	15	HMD	Passive viewing	none	Classification	S-transform EEG features for cybersickness




Fig. 2: Distribution of reviewed papers by year and primary research objective.

2) *Sample Sizes and Demographics*: The 92 studies span 2001–2025. Across the corpus, 2687 participants were analyzed (recruited: 2822), with a mean of 29.2 ± 29.0 participants analyzed per study (median: 22, range: 1–202). Of the 92 studies, 62 report a mean age (yielding a corpus mean of 23.7 ± 2.7 years, indicating a focus on young-adult populations); 15 report a minimum–maximum range without a mean, and 15 report no parseable age information. Fig. 3 shows the age distribution across studies.

EEG channel counts vary widely (median 30, range 1–256, mean 39.6 ± 55.8 , $n = 72$ studies reporting). The most frequent montages are 32-channel (18 studies), 64-channel (9), 14-channel (8), 16-channel (5), and 30- or 256-channel (4 each); a long tail of low-channel configurations (4 to 16 channels) is consistent with portable in-VR recording.

Fig. 4 shows the distribution of EEG channel configurations across studies. Channel count is positively correlated with reported within-subject classification accuracy ($r = 0.403$, $p = 0.037$, $n = 27$); this observation is revisited in Section IV-F.

Concurrent physiological recording is reported in 36 of 92 studies (39.1%). Most commonly used are electro-oculography (18 studies, 19.6%), electrocardiography (16, 17.4%), and galvanic skin response (9, 9.8%). Present but less common are electromyography (2, 2.2%), electrogastrography (4, 4.3%), photoplethysmography (7, 7.6%), respiration (2, 2.2%), skin temperature (4, 4.3%), and blood pressure (1, 1.1%).

3) *Experimental Design and Data Collection Protocols*: Study designs predominantly employ within-subject comparisons (78 of 92, 84.8%), with 7.6% employing between-subject designs exclusively and 7.6% using both. The majority of protocols are single-session (mean 1.6 ± 2.1 sessions, median 1, $n = 91$). Session duration averages 36.3 ± 32.6 minutes (median 25.8, range 1.8–180, $n = 76$), and the VIMS-inducing stimulus block within a session averages 15.2 ± 15.4 minutes (median 10.0, range 0.5–90, $n = 86$).

Virtual reality is delivered through head-mounted displays in 71.7% of studies (66 of 92), with screens (9.8%) as the next most common configuration, followed by projection systems (8.7%), CAVE (4.3%), television (2.2%), and other or




Fig. 3: Age distribution across studies showing mean age (black dots) with age ranges (dotted bars) and standard deviations when available (solid bars). The age range shows the minimum and maximum participant ages, while the standard deviation shows the typical spread around the mean age.

mixed-modality setups (3.3% combined). Virtual environments cluster on navigation (17 studies, 18.5%), roller-coaster experiences (16, 17.4%), games (15, 16.3%), driving (13, 14.1%), passive video (10, 10.9%), boating or flight (7, 7.6%), and space simulators (5, 5.4%); categories are non-exclusive, as a study may employ more than one type of environment. The joint distribution of device type and environment is shown in Fig. 5; the hardware mix bears on the meta-analytic findings in Section IV-F.

4) *VIMS Assessment Methods*: VIMS is measured through questionnaire instruments in 70 of 92 studies (76.1%), continuous ratings during exposure in 20 (21.7%), single-item Likert scales in 15 (16.3%), and behavioral indices in 8 (8.7%); these categories sum above 100% because 19 studies (20.7%)




Fig. 4: Distribution of EEG channel configurations across VIMS studies.




Fig. 5: Heatmap of the distribution of VR device types and virtual environments across studies. Studies employing multiple devices or environments contribute toward each.

use more than one method on the same exposure. Across the corpus, the Simulator Sickness Questionnaire is the most common instrument (57 of 92 studies, 62.0%), followed by the Fast Motion Sickness Scale (6, 6.5%), the Motion Sickness Susceptibility Questionnaire (2, 2.2%), and the Virtual Reality Sickness Questionnaire (2, 2.2%); 6 studies (6.5%) use other instruments, and 19 (20.7%) either use no questionnaire or do not name one.

5) *Baseline Conditions and Preprocessing Heterogeneity:* Baseline conditions vary across the corpus: 39 of 92 studies (42.4%) use a task baseline, 16 (17.4%) use an eyes-open resting baseline, 8 (8.7%) use a pre-stimulus segment of the same recording, 5 (5.4%) use an eyes-closed resting baseline, and the remainder either do not specify a baseline (3, 3.3%) or do not report one (21 missing, 22.8%, including the neurostimulation-only studies). The 5 eyes-closed cases are a minority of the corpus, but alpha-band reactivity to eye closure overlaps with the alpha modulations attributed

to VIMS; Section V develops this critique and discusses recommended alternatives.

Across the 77 EEG corpus studies, band-pass and notch-filter reporting is heterogeneous: 11 (14.3%) do not report a band-pass filter, and 59 (76.6%) do not state a notch-filter frequency. Of the 66 that report a band-pass setting, 59 give numeric low and high cutoffs, summarized in Fig. 6; among these, low cutoffs span 0–8 Hz and high cutoffs 15–100 Hz, and no configuration predominates in the corpus.

Band-pass Filter Frequency Ranges Across Studies




Fig. 6: Band-pass filter frequency ranges across studies showing center frequency (black dots) with frequency ranges (black error bars) and notch filter frequencies (red crosses). The frequency range shows the low and high cutoff frequencies of the band-pass filters, while red crosses indicate notch filter frequencies used to remove specific interference (e.g., 50/60 Hz power line noise).

B. Neural and Physiological Correlates of VIMS

Across the 77 EEG studies, 4 fNIRS studies, and the subset using event-related potential analyses, the corpus converges

on a low-frequency power increase during VIMS; higher-frequency and lateralized findings are less consistent at the corpus level and, as shown in Section IV-F, are partly attributable to differences in VR hardware platform.

1) *EEG Spectral Power*: The EEG spectrum is conventionally divided into delta (1–4 Hz), theta (4–8 Hz), alpha (8–13 Hz), beta (13–30 Hz), and gamma (>30 Hz) bands, though the exact cutoffs applied to each band vary across individual studies [122]. Across the studies reporting per-band changes during VIMS, the most consistent direction is an increase in low-frequency power. Of the 30 studies reporting a delta-band change, 16 (53.3%) report an increase, 6 a mixed pattern, 5 no change, and 3 a decrease. Of the 35 studies reporting a theta-band change, 20 (57.1%) report an increase, 6 a decrease, 5 a mixed pattern, and 4 no change. Increases in delta and theta power have been associated with the onset and increasing severity of cybersickness across diverse VR stimuli and hardware [33], [34], [43], [66], [70].

Alpha-band findings diverge across the corpus. Of the 39 studies reporting an alpha change, 15 (38.5%) report an increase, 11 (28.2%) report a decrease, 9 (23.1%) report a mixed pattern, and 4 (10.3%) report no change. Studies that report alpha suppression, particularly over central, parietal, and occipital cortices, interpret it as a marker of increased cognitive load under sensory conflict or as a consequence of postural-control demand [27], [53], [69]. Studies that report an alpha increase attribute it to sensory gating or to drowsiness during prolonged exposure. Section IV-F shows that the direction of the alpha change tracks VR display hardware, and reframes part of this corpus-level heterogeneity as hardware-attributable.

Higher-frequency bands show less consistent patterns than delta and theta. Of the 36 studies reporting a beta-band change, 11 (30.6%) report an increase, 10 (27.8%) a decrease, 10 a mixed pattern, and 5 no change. Individual reports range from a beta decrease [70] to a beta increase [108], and one study identifies beta as the most informative band for channel selection in classification [123]. Of the 21 studies reporting a gamma-band change, 7 (33.3%) report an increase, 6 no change, 5 a mixed pattern, and 3 a decrease. The full distribution of rhythm changes across anatomical regions is shown in Fig. 7; as with alpha, the beta findings are partly attributable to display hardware (Section IV-F).

Test-retest reliability for delta, theta, and alpha power changes in frontal and central areas has been reported in 1 corpus study [84], and delta-band normalization has been used as a marker of the post-exposure recovery time course [115]; the recovery literature is discussed in Section V.

Beyond band-power, a subset of studies has examined additional spectral and information-theoretic features (entropy, complexity measures, amplitude modulation, mean theta-band frequency) [58], [88], [89], [103], [124]. Functional connectivity is investigated in 16 studies using heterogeneous metrics (coherence, phase-locking value, transfer entropy, Granger causality), and no single direction of change emerges at the corpus level. An interpretable signal comes from the subset that examines specific visual-vestibular pathways, which report reduced or altered coupling during VIMS,




Fig. 7: Brain rhythm changes across anatomical regions during VIMS.

consistent with the sensory-conflict account: transfer entropy between cortical regions decreases at VIMS onset [33], connectivity between visual area V5 and vestibular regions is reduced [96], and Granger-causal analysis indicates altered directed interaction within the visual-vestibular network [42]. A separate group of studies instead uses connectivity features, such as phase-locking value, as inputs to deep classification models without assigning a directional interpretation [65], [105]. The causal relevance of these patterns is supported by intervention studies in which symptom mitigation correlates with normalization of the same EEG bands [120].

2) *Event-Related Potentials and Evoked Responses*: Beyond ongoing oscillations, a subset of studies has used event-related potentials (ERPs) to characterize how VIMS

affects discrete cognitive processes. These studies indicate that VIMS is accompanied by measurable changes in attentional allocation and response inhibition. VIMS severity is associated with a reduction in the amplitude of the P3b component in a dual-task paradigm, a marker of diminished attentional resources [6]. VIMS has also been shown to impair response inhibition, as reflected by a larger N2 amplitude and a smaller, delayed P3 component during an oddball task [116]. Read alongside the sensory-conflict and cognitive-load accounts introduced in Section II, these findings are consistent with VIMS consuming processing resources that would otherwise support task performance.

Susceptibility-related ERP patterns recur in studies that compare resistant and susceptible participants. VIMS-susceptible participants exhibit larger parietal N2 amplitudes and reduced theta-band phase synchronization compared to resistant counterparts [125], and individuals with high sickness levels show higher P3 amplitudes during sensory-mismatch conditions [42]. A separate ERP approach has linked the subjective experience of disorientation to a suppression of fronto-central heartbeat-evoked potentials, providing a link between interoceptive processing and a key symptom of cybersickness [51]. These susceptibility markers are revisited in Section IV-D alongside the resting-state-prediction literature.

3) *Hemodynamic Responses (fNIRS)*: Four fNIRS studies in the corpus report convergent regional hemodynamics during VIMS, providing a small but coherent body of evidence on the cortical regions involved. Bilateral angular-gyrus deactivation has been reported in association with subjective sickness scores [37], and increases in total hemoglobin concentration have been linked directly to VIMS onset [11], [38]. Reduced cortical activation during low-resolution stereoscopic exposure has also been documented [39].

The direction of the hemodynamic effect is not uniform across the small fNIRS literature: regional perfusion increases in some cortical areas coexist with decreases in others. The two fNIRS-based closed-loop mitigation systems reported in the corpus [11], [38] are taken up in Section IV-E.

C. Classification and Prediction of VIMS

Of the 92 corpus studies, 38 have classification or prediction of VIMS as their primary objective; a broader 41 use classification or regression methods as part of mixed-objective work. Corpus-level claims in this subsection are restricted to the 38-study primary-objective subset. Across these 38 studies, accuracy is the dominant reported performance metric, but sensitivity/specificity, AUC/F1, and the operational definition of the sick/non-sick label are reported in fewer than half of the subset.

1) *Methodological Approaches and Feature Engineering*: Computational approaches across the 38-study subset divide between traditional machine learning (15 of 38, 39.5%), deep learning (14, 36.8%), and regression (6, 15.8%); thresholding (1) and other approaches (2) account for the remainder. Traditional pipelines rely on hand-crafted features (Power Spectral Density, wavelet-derived energy ratios [77], spectral




Fig. 8: Distribution of classification method categories across VIMS studies.

entropy and skew [58], time-domain statistics such as signal variance [93]) fed to Support Vector Machines, k-Nearest Neighbors, or Linear Discriminant Analysis [49], [126]; dimensionality reduction is addressed through Correlation Feature Selection [123] or genetic algorithms [127].

Deep-learning pipelines learn features directly from the data. Convolutional Neural Networks extract spatial and spectral features from EEG time series, spectrograms, or topographic scalp maps [67], with recurrent layers (Long Short-Term Memory [83], Gated Recurrent Units [92]) capturing temporal dynamics. Other architectures include spiking neural networks [119], shallow CNNs designed for real-time detection in mitigation systems [10], hybrid CNN-LSTM models [105], dual-pathway designs that process raw EEG and functional-connectivity inputs in parallel [65], and channel-attention mechanisms [62], [63]. Multimodal fusion combines EEG with peripheral physiological signals [104], with electromyography from postural muscles (reported as more informative than concurrent EEG in one study) [102], or with VR video features via transfer learning [71]. One paper investigates explainable-AI interpretation [104].

2) *Reported Performance and Discrimination Metrics*: Model validation predominantly employs k-fold cross-validation (21 of 38, 55.3%), with train-test splits (6), leave-one-subject-out cross-validation (1), and other or unspecified schemes (10) making up the remainder. The distribution of methods is summarized in Fig. 8 and Fig. 9.

Reported within-subject classification accuracies are high but rest on a small denominator: across the 8 of 38 studies that report a within-subject accuracy, the mean is $90.7\% \pm 8.4\%$ (median 91.5%, range 73.3–100.0%). Cross-subject accuracies are reported by 22 of 38 studies, with a mean of $88.1\% \pm 11.6\%$ (median 88.9%, range 55.0–99.7%). The two distributions are compared in Fig. 10, and the temporal trend of reported accuracies is shown in Fig. 11.

These headline accuracies are reported on heterogeneous discrimination metrics. Of the 38 classification studies, 13




Fig. 9: Detailed breakdown of specific methods within each classification category.

(34.2%) report a headline accuracy with no sensitivity, specificity, AUC, or F1; only 7 (18.4%) report sensitivity and specificity, and 10 (26.3%) report AUC or F1. Ten further studies (26.3%) report only regression metrics (R^2 , RMSE) without a classification accuracy, and 2 (5.3%) report no quantitative performance.

3) *Classification Tasks*: Across the 38-study subset, binary classification of a “sick” versus “not sick” state is the most common task. Several studies report cross-subject accuracies above 95% on this binary task [49], [62], [63], [65], [67], [121]. The sick/non-sick label that grounds these accuracies varies in how it is operationalized: 19 of 38 classification studies disclose the threshold or its basis, of which 8 use a numeric cutoff and the remainder use self-reported binary labels, pre/post differences, or total-questionnaire-score cutoffs. The other 19 studies do not document the threshold at all.

A second task is multi-level classification, which categorizes VIMS into several discrete levels of severity. Three-level classification has been demonstrated with multimodal fusion [9], [76] and with EEG alone using rule-based induction [99] or a genetic algorithm combined with a Support Vector Machine [86].

A third task is continuous prediction or regression of a VIMS severity score, measured by correlation coefficients and error metrics. Deep-learning models have demonstrated high correlations between predicted and actual SSQ scores [61], [64], [75], [90], [91], [118], while earlier work used Support Vector Regression [73], [114] and linear regression [128].

D. Individual Differences and Susceptibility

Resting-state neural features have been used to predict VIMS susceptibility before exposure, and during exposure susceptible and resistant individuals show distinct neural signatures. Demographic moderators are reported in a small subset of the corpus and warrant cautious interpretation given the small effect sizes documented in the broader literature.

Pre-exposure prediction of VIMS has been demonstrated in 3 corpus studies. A spiking neural network classified subsequent sickness with 85.9% accuracy from pre-exposure resting-state EEG alone [119]. Resting-state EEG beta power in the left parietal region has been reported as a significant predictor of subsequent motion-sickness ratings in an in-car VR platform [81], and aperiodic components of the EEG signal have been reported as predictive of susceptibility to rotational motion [110]. These findings point to pre-existing neural traits that influence susceptibility, although the evidence base remains small.

During VIMS exposure, resistant and susceptible individuals show distinct neural signatures. Resistant individuals exhibit increased theta activity in the left parietal cortex, a candidate biomarker of resilience [80]. Susceptible individuals show larger parietal N2 ERP amplitudes and reduced theta-band phase synchronization [125], and higher P3 amplitudes during sensory-mismatch conditions [42]. One study reports that the cybersickness condition produces increased delta and decreased alpha power in both susceptible and non-susceptible groups, but susceptible participants additionally show lower absolute power in the beta and gamma bands at baseline [66]. An earlier study using a long-duration driving task found that the ratio of theta to total EEG power in frontal and parietal lobes differed between sick and non-sick groups, proposing this ratio as a physiological index of sensitivity [129].

Sex and gender differences are reported in 3 of 92 corpus studies, all from a single laboratory using a VR-neurofeedback paradigm [46]–[48]. Across all 3 studies, women reported greater cybersickness than men; in 2 of the 3, higher cybersickness was additionally associated with poorer sensorimotor-rhythm neurofeedback performance [47], [48]. An additional 11 corpus studies use male-only samples (precluding sex comparisons) and 15 do not report participant sex composition, leaving 63 mixed-sex studies that report composition without analyzing sex effects.

Individual variability in VIMS susceptibility has measurable neural correlates and small demographic moderators. At




Fig. 10: Classification accuracy distributions by method category for within-subject and cross-subject validation approaches.




Fig. 11: Classification accuracy over time, comparing within-subject and cross-subject validation approaches.

present, the corpus is underpowered for either to be operationalized as a predictive biomarker. The mitigation literature reviewed next acts on the neural signals identified here.

E. Mitigation and Intervention Strategies

Mitigation research in the corpus splits into two threads: closed-loop adaptive VR that modifies the virtual environment in response to neurophysiologically detected sickness (3 corpus studies), and interventions that target the user's brain state directly (5 corpus studies). Both threads are early-stage, and the small literature already documents one methodological obstacle specific to the closed-loop setting.

1) *Closed-Loop Adaptive VR*: Three corpus studies implement neurophysiology-driven closed-loop adaptation: the literature is in early-stage development. One system uses a two-stage shallow convolutional network to detect

cybersickness from EEG and infer a likely causal factor, then selectively modifies the implicated VR parameter (navigation speed or scene complexity), producing a significant reduction in user-reported sickness compared to a control condition [10]. Two further systems use fNIRS to monitor total hemoglobin concentration; on detecting increases correlated with sickness, the system dynamically reduces the user's field of view, demonstrating the feasibility of an fNIRS-based closed-loop pipeline [11], [38].

2) *Interventions Modulating Brain State*: The corpus includes 5 studies that investigate interventions intended to act on the underlying physiological state of the user, most using concurrent neurophysiological recording to verify engagement. Providing synchronized auditory and physical motion cues during a simulated bicycle ride reduced VIMS scores relative to conditions with mismatched or absent cues, with the effect reflected in concurrent EEG recordings [120].

Transcranial alternating-current stimulation has been applied online during VR exposure to target oscillations implicated in VIMS. Random-phase tACS at 18 Hz over the left parietal cortex, chosen to disrupt a sickness-correlated oscillation identified from EEG, reduced reported nausea relative to sham and control stimulation while the user remained in VR [82]. In a separate double-blind trial, 10 Hz tACS modeled to reach the vestibular cortex bilaterally reduced cybersickness nausea during a VR roller-coaster in a frequency-dependent, placebo-insensitive manner, paralleled by a modulation of galvanic skin response [40]. Because the effect in both studies depended on the stimulation frequency, these results provide causal evidence that the targeted oscillations contribute to VIMS rather than only accompanying it.

Two further studies, both from the same laboratory, investigated neurofeedback protocols in which users were trained to volitionally modulate their own brain activity. The presence of cybersickness symptoms was found to directly reduce the user's ability to perform the neurofeedback training task [48], and a placebo transcranial direct current

stimulation procedure, while having no significant effect on reported sickness symptoms, degraded performance on a concurrent neurofeedback task [47]. These findings illustrate a methodological challenge for any neuromarker-driven closed-loop intervention: the physiological state being targeted can interfere with the user's ability to engage with the intervention itself.

F. Corpus-Level Methodological Observations

Meta-analytic comparison across the 92 corpus studies indicates that VR display hardware is associated with EEG band-power responses and that stimulus-exposure duration is confounded with hardware. The findings in this subsection are observational; the interpretive critique and reporting recommendations they motivate are developed in Section V.

Alpha-band VIMS responses differ between head-mounted-display and screen-based studies. The Mann–Whitney U test asks whether the ordered alpha-change codes differ between the two display classes; Cliff's δ gives the direction and size of that separation. This comparison yields Cliff's $\delta = 0.875$ ($p = 0.004$), indicating that head-mounted-display studies are more likely than screen-based studies to report alpha increases. Among head-mounted-display studies with a reported directional change, alpha increases occur four times as often as decreases (12 of 24 versus 3 of 24); screen-based studies in the same comparison report decreases only. This pattern reframes the alpha-band heterogeneity reported in Section IV-B as partly hardware-attributable, rather than as a pure intersubject or task effect. The same head-mounted-display versus screen-based comparison for beta yields Cliff's $\delta = 0.664$ ($p = 0.015$), indicating the same direction of association beyond alpha. No corresponding between-device differences are detected for delta, theta, or gamma.

Stimulus-exposure duration is itself confounded with hardware. The Mann–Whitney comparison for exposure time shows that HMD studies use shorter exposure periods (mean 12.9 ± 14.8 minutes, $n = 62$) than non-HMD studies (mean 23.6 ± 15.1 minutes, $n = 21$; $U = 324.0$, $p < 0.001$, Cliff's $\delta = -0.502$, large effect). The hardware confound therefore covaries with how long the user is exposed, not only with the display optics; comparisons of band-power changes between HMD and screen studies necessarily mix the two effects.

These corpus-level observations, together with the inline reporting-gap counts in Section IV-C and the baseline-heterogeneity findings in Section IV-A, seed the design and reporting recommendations developed in Section V.

V. DISCUSSION

The prior review of cybersickness EEG research by Chang et al. [8] reported delta-band increases, beta-band decreases, and inconsistent alpha-band findings; the present corpus refines this picture in two respects. The delta- and theta-band increases reproduce in the current corpus (16 of 30 studies for delta; 20 of 35 for theta). The reported alpha-band inconsistency partly resolves once VR display hardware is accounted for: head-mounted-display and screen-based studies differ with large effects on alpha (Cliff's $\delta = 0.875$, $p = 0.004$) and beta ($\delta = 0.664$, $p = 0.015$).

A. Objective Markers and the Limits of Self-Report

VIMS has no diagnostic biomarker; its ground truth is self-report. It is defined by a symptom profile (nausea, disorientation, and oculomotor disturbance) composed of private experiences that cannot be externally verified [7], [17]; any candidate objective marker must therefore be calibrated against self-report rather than replacing it. Functional cognitive consequences of this state are documented in 4 corpus studies: working-memory decrements scaling with reported severity [6], inhibitory-control demand reflected in N2/P3 components [116], and degraded sensorimotor-rhythm neurofeedback performance under cybersickness [47], [48], indicating that the state extends beyond subjective discomfort and carries measurable functional costs. The EEG and fNIRS correlates established in Section IV-B are objective physiological measurements, but each was validated against self-reported severity and therefore presupposes the measure it would replace. Among the candidate objective markers, postural sway has a direct theoretical anchor in the postural-instability account, which predicts that pre-symptomatic sway covaries with subsequent severity [25], [26]; VR-specific reports document moderate coupling [9], [29], [130]. Pupillometry is a second candidate [131], [132], subject to arousal and cognitive-load confounds. Of the 92 corpus studies, 16 already co-record force-plate, balance-board, eye-tracker, or pupillometry alongside EEG; treating these signals as candidate outcome variables rather than covariates is a prerequisite for validating them as objective markers.

B. Methodological Challenges and Critical Perspective

Four classes of methodological issue emerge from the corpus and constrain cross-study synthesis in VIMS neuroimaging and neurostimulation: (1) classification reporting practices, (2) baseline selection, (3) protocol and hardware heterogeneity, and (4) conceptual and terminological variability.

Across the 38 classification or prediction studies in the corpus, three reporting practices limit the interpretability of headline performance numbers. Thirteen studies report a headline accuracy without sensitivity, specificity, AUC, or F1; under class imbalance, accuracy collapses toward the majority-class baseline and provides little information about the classifier's behavior at decision-relevant operating points [133], [134]. Nineteen studies disclose no sick-versus-non-sick threshold (neither a numeric cutoff nor a stated basis), and only 8 give an explicit numeric cutoff converting SSQ or FMS scores into a label; the target variable is therefore operator-defined and not comparable across studies. One study reports leave-one-subject-out validation; 21 use k-fold cross-validation without subject stratification, which permits subject-identity leakage from training into the test fold and inflates apparent generalization [135]. These three gaps are consistent with the high headline accuracies recurring in the corpus, including binary cross-subject accuracies above 95% [62], [63], [65], [67], [121] and pre-post resting-state contrasts approaching 99% [49]: where accuracy is the only reported metric, the threshold defining the label is undocumented, and

the validation scheme does not respect subject identity, the headline number does not support a generalization claim. A related ground-truth issue compounds these reporting gaps: the SSQ is retrospective and coarse, and continuous motor in-task ratings such as joystick ratings risk encoding the motor act of reporting rather than the sickness state itself, so the very signal that supervises classification is heterogeneous before any modeling choice is made.

A second design issue concerns the baseline against which in-exposure neural change is measured. Closing the eyes elicits a large posterior alpha increase relative to eyes-open [136]–[138], so a study that contrasts VR exposure against an eyes-closed pre-task recording confounds the alpha-change estimate with a state shift unrelated to the visual stimulus. The VR exposure condition is, by construction, eyes-open inside a head-mounted display viewing a structured scene; an eyes-closed pre-task block is therefore not matched to the exposure condition on visual input, oculomotor state, or vigilance. This baseline choice is a minority practice in the corpus (5 of 77 EEG studies, or 6 if eyes-open + eyes-closed composites are counted), but distorts the alpha-band estimates of the affected studies, which typically present those estimates as primary outcomes.

Baseline choice is one of several protocol parameters that vary widely across the corpus, and the broader heterogeneity carries methodological consequences. At the corpus level, VR display hardware substantially modulates the band-power response: head-mounted-display and screen-based studies differ on the direction of alpha change with a large effect, and the alpha-band inconsistency reported in the prior review [8] partly resolves once hardware is accounted for (see Section IV-F). Stimulus exposure duration is itself confounded with display: head-mounted-display studies use shorter exposures than non-head-mounted-display studies, so comparisons of band-power change between display classes necessarily mix the two effects. Recording quality varies similarly, ranging from research-grade systems to low-cost consumer-grade devices with fewer channels and lower signal quality [89], and the corpus is biased toward low-density montages: among the 72 EEG studies that disclose channel count, 58 (80.6%) use 32 or fewer channels and 14 (19.4%) use 10 or fewer. Low-density montages are appropriate for applied paradigms that target well-characterized localized signals; for exploratory characterization of VIMS neurophysiology they have insufficient spatial resolution for reliable source localization. Reporting quality compounds the hardware-protocol heterogeneity: of the 77 EEG corpus studies, 43 (55.8%) do not state the reference electrode and 59 (76.6%) do not state a notch-filter frequency, which limits both reproducibility and meta-analytic re-pooling. Hardware, stimulus, montage, and reporting parameters vary study by study; requiring several to match at once leaves little overlap. In the corpus, the largest subgroup that aligns on sickness measure (SSQ), display (head-mounted display), stimulus class (game), and modality (EEG) contains only 5 of the 92 corpus studies. Cross-study generalization is therefore difficult: the field has no replicable majority paradigm, and statements that pool results across the literature aggregate

category labels rather than a shared experimental design. Samples are small and demographically narrow (typically young university students), and 11 of the 92 studies recruited male-only samples [66], [96], which limits statistical power for individual-difference questions including the sex/gender effects discussed in Section V-C. The experiences and stimuli are also heterogeneous: passive 360-degree video [108] is not equivalent to active interactive navigation [27], and findings from head-mounted-display, screen, CAVE, and smartphone-VR systems [84], [139] are often discussed interchangeably.

Finally, the field exhibits conceptual variability in both the use of theory and the use of terminology. Most corpus studies (73 of 92 explicitly, 14 implicitly) cite an etiological theory of VIMS in their introduction or methods, but few return to that framework when interpreting their neurophysiological findings; a neural feature observed in a given paradigm could reflect sensory conflict, postural control, expectation-related vection, display optics, or some combination, and without an explicit theoretical commitment the literature cannot adjudicate between these possibilities. Terminology compounds the interpretive ambiguity: motion sickness, visually induced motion sickness, simulator sickness, and cybersickness are related but non-identical constructs tied to different exposure contexts (see Section II), yet they are frequently used interchangeably [14], [15], [17], [18]. A further consequence of these practices is the labeling of any statistically significant feature as a “neuromarker” without establishing its specificity, sensitivity, or reliability across contexts.

Taken together, these methodological and conceptual issues constrain cross-study synthesis and reproducibility.

C. Future Directions

The methodological challenges documented in Section V-B translate into six concrete directions for future work: standardized reporting and open data, causal mechanisms, generalization and ecological validation, sex- and gender-stratified design, objective behavioral markers, and longitudinal characterization of adaptation and recovery.

1) *Standardized Reporting and Open Data:* Standardized reporting represents a gap documented across the corpus. For classification and prediction studies, the minimum items follow from the gaps documented in Section V-B: a discriminative metric (AUC or balanced accuracy) with 95% confidence interval, sensitivity and specificity at a stated operating point, the numeric label threshold, a subject-respecting validation scheme, and comparison against a trivial and a non-neural feature baseline. For studies that quantify in-exposure neural change, a baseline matched to the exposure condition on visual input, oculomotor state, and vigilance is required for reproducibility, with the choice and its rationale reported; the available options are discussed in Section V-B. For EEG, the reference electrode and notch-filter frequency are required for reproducibility and meta-analytic re-pooling. For subjective sickness measurement, reporting the questionnaire instrument, its administration time-point relative to exposure end, and the numeric cutoff used to assign class labels enables

cross-study comparability; where continuous in-task ratings are used, the response modality and any associated motor artifact exclusion strategy are necessary for reproducibility. For stimulus and display, the device model, field-of-view, and frame rate should be reported alongside the stimulus class (passive or active, recorded or interactive); adoption of a small number of canonical reference paradigms, at minimum a passive visual-flow exposure [108] and an active first-person navigation task [27], would allow direct cross-laboratory comparison without the confound of idiosyncratic stimulus design.

These reporting standards are a precondition for, but not a substitute for, shared infrastructure: community-curated benchmark datasets with multimodal recordings (EEG, EOG, ECG, head motion, sway, eye tracking), well-defined subjective labels, and coverage of both passive and active paradigms. A small number of corpus papers have already released public datasets [55], [71]; releasing analysis code and pre-trained models alongside the data would further support replication and cumulative analysis.

2) *Causal Mechanisms*: The causal status of documented neural associations remains largely unexamined. The literature has documented associations between specific neural patterns and VIMS across multiple studies, but whether these associations are causal has not been established. Non-invasive brain stimulation offers a direct test: transcranial alternating-current stimulation can target specific oscillations, and corpus studies that apply it online during VR exposure [40], [82] provide a methodological template for asking whether modulating a candidate neuromarker changes sickness severity. Coupled with computational models of multisensory integration and predictive coding [33], interventional designs can begin to distinguish neural features that drive the VIMS experience from features that only accompany it.

3) *Generalization and Ecological Validation*: Generalization and ecological validation remain open challenges for models built for VIMS. In the 4 corpus studies that report both within- and cross-subject accuracy, cross-subject performance is lower in each case, and only 1 of 38 classification studies reports a leave-one-subject-out evaluation; transfer-learning and domain-adaptation approaches that explicitly model inter-subject and inter-session variability address a current gap if brain-directed methods are to operate without per-user calibration. Validation beyond the laboratory is a further open direction: at-home, naturalistic paradigms such as the in-home gaming setup of Moineau et al. [140] address the question of whether laboratory neuromarkers survive the noise of consumer-grade hardware and uncontrolled environments.

4) *Sex- and Gender-Stratified Design*: Sex and gender represent an underexplored factor in the corpus. Of the 92 corpus studies, 11 used male-only samples, 15 did not report sex composition, and only 3 (all from a single laboratory) reported higher cybersickness in women [46]–[48]. The broader literature on sex differences in VIMS susceptibility is mixed: Kelly et al. [141] argue that reported effects are modest in magnitude and plausibly mediated by interpupillary-distance fit, prior gaming experience, and

questionnaire framing; Kourtesis et al. [132] report that male–female differences disappear when prior gaming experience is included as a covariate; and Lawson and Bolkhovskiy [142] caution that the claim of greater female susceptibility is overstated in the simulation literature. Taken together, the 3 corpus reports and the broader literature are consistent with a modest, variable effect whose magnitude and moderators cannot be resolved at individual-study scale, and whose apparent direction may partly reflect gaming-experience differences and questionnaire framing rather than a stable sex difference in susceptibility. Reporting sex-stratified outcomes (band-power changes, connectivity, classification metrics), even when single-study samples are underpowered for inferential testing, would enable subsequent meta-analytic re-pooling. Logging interpupillary-distance setting and gaming experience as covariates would address both the sex confound and the hardware effects documented in Section IV-F. Adequately powered designs, such as pooled multi-site cohorts or longitudinal within-subject designs, are the precondition for resolving the magnitude and moderators of a sex effect on VIMS.

5) *Objective Behavioral Markers*: Objective behavioral markers represent an underdeveloped line of investigation. The candidate markers introduced in Section V-A are already collected by 16 of the 92 corpus studies, but typically as covariates rather than as candidate ground truth. Postural sway is the strongest theoretically anchored candidate: reporting a coupling estimate between pre-symptomatic sway dynamics and subsequent self-reported sickness, alongside the protocol metadata that this estimate depends on (stance, surface, sampling rate), would anchor the construct across laboratories. Pupillometry can supplement self-report subject to arousal and cognitive-load confounds [131], and requires controls on ambient luminance and on task demand. Heart-rate variability, electrodermal activity, and head-motion dynamics are further candidates whose corpus-level predictive value is not yet quantified [110], [119], [143]; multi-modal integration that fuses behavioral, autonomic, and neural signals offers a more holistic state estimate than any single modality. Treating these markers as outcome variables, rather than as covariates, is the precondition for shifting VIMS research toward objective ground truth.

6) *Longitudinal Characterization of Adaptation and Recovery*: Longitudinal designs represent a gap in the current corpus. The corpus is dominated by single-session experiments, leaving the neurodynamics by which users adapt to VR (“getting one’s VR legs”) largely uncharacterized, and the single corpus recovery study [115] is insufficient to support strong claims about residual neural state. Multi-session within-subject designs would enable tracking of how the EEG signatures identified in this review evolve with repeated exposure, and mapping the dissipation of post-exposure signatures onto behavior.

VI. CONCLUSION

This systematic review synthesized 92 studies on the neurophysiology of visually induced motion sickness in VR,

uniting EEG-based correlate and classification work with fNIRS hemodynamic evidence as candidate neuromarkers, alongside non-invasive neurostimulation as a complementary approach.

The corpus converges on a low-frequency EEG signature of VIMS (increased delta and theta power) that reproduces across diverse stimuli, hardware, and laboratories, and is consistent with the pattern reported by the prior review of Chang et al. [8]. Alpha-band findings have been inconsistent in the literature; the meta-analytic comparison reported here attributes a substantial portion of that inconsistency to VR display hardware, with head-mounted-display and screen-based studies systematically differing in the direction of alpha and beta responses. This hardware confound, compounded by differences in stimulus-exposure duration, limits the extent to which band-power results can be pooled across the literature.

The corpus also documents attentional and inhibitory-control costs during VIMS in event-related potential studies, regional hemodynamic correlates in fNIRS, and pre-exposure resting-state predictors of individual susceptibility. Early closed-loop systems (adaptive VR pipelines [10], [11] and direct brain-state interventions [40], [82]) demonstrate that neurophysiological detection of VIMS is sufficient in principle to drive real-time mitigation, though the evidence base for both threads remains small.

Alongside these findings, the review identifies methodological barriers that constrain generalization: sparse and heterogeneous reporting in the classification literature, baseline-selection practices that confound neural estimates, hardware and protocol heterogeneity that prevent direct cross-study replication, and terminological variability that complicates theoretical interpretation. The causal status of the observed neural associations, cross-subject and ecological generalization of detection models, the role of sex and gender, and the neurodynamics of adaptation and recovery over repeated exposure remain open questions in the current literature.

ACKNOWLEDGMENTS

This research was funded by the European Union ERANET CHIST-ERA 2020 (CHIST-ERA-20-BCI-003 - "GENESIS") and the French Research Agency (ANR-21-CHRA-0001-01).

REFERENCES

- [1] J. Chatain, V. Ramp, V. Gashaj, V. Fayolle, M. Kapur, R. W. Sumner, and S. Magnenat, "Grasping Derivatives: Teaching Mathematics through Embodied Interactions using Tablets and Virtual Reality," in *Proceedings of the 21st Annual ACM Interaction Design and Children Conference*, ser. IDC '22. New York, NY, USA: Association for Computing Machinery, Jun. 2022, pp. 98–108.
- [2] M. A. Gigante, "1 - Virtual Reality: Definitions, History and Applications," in *Virtual Reality Systems*, R. A. Earnshaw, M. A. Gigante, and H. Jones, Eds. Boston: Academic Press, Jan. 1993, pp. 3–14.
- [3] S. S. Kardong-Edgren, S. L. Farra, G. Alinier, and H. M. Young, "A Call to Unify Definitions of Virtual Reality," *Clinical Simulation in Nursing*, vol. 31, pp. 28–34, Jun. 2019.
- [4] P. Caserman, A. Garcia-Agundez, A. Gámez Zerban, and S. Göbel, "Cybersickness in current-generation virtual reality head-mounted displays: Systematic review and outlook," *Virtual Reality*, vol. 25, no. 4, pp. 1153–1170, Dec. 2021.
- [5] S. Weech, S. Kenny, and M. Barnett-Cowan, "Presence and Cybersickness in Virtual Reality Are Negatively Related: A Review," *Frontiers in Psychology*, vol. 10, 2019.
- [6] K. J. Minnaugh, E. G. Center, M. Suomalainen, I. Becerra, E. Lozano, R. Murrieta-Cid, T. Ojala, S. M. LaValle, and K. D. Federmeier, "Virtual Reality Sickness Reduces Attention During Immersive Experiences," *IEEE Transactions on Visualization and Computer Graphics*, vol. 29, no. 11, pp. 4394–4404, Nov. 2023.
- [7] R. S. Kennedy, N. E. Lane, K. S. Berbaum, and M. G. Lilienthal, "Simulator Sickness Questionnaire: An Enhanced Method for Quantifying Simulator Sickness," *International Journal of Aviation Psychology*, vol. 3, no. 3, p. 203, Jul. 1993.
- [8] E. Chang, M. Billinghurst, and B. Yoo, "Brain activity during cybersickness: A scoping review," *Virtual Reality*, vol. 27, no. 3, pp. 2073–2097, Sep. 2023.
- [9] M. S. Dennison, M. D'Zmura, A. V. Harrison, M. Lee, and A. J. Raglin, "Improving motion sickness severity classification through multi-modal data fusion," in *Artificial Intelligence and Machine Learning for Multi-Domain Operations Applications*, T. Pham, Ed. Baltimore, United States: SPIE, May 2019, p. 27.
- [10] U. Uyan and U. Celikcan, "CDMS: A real-time system for EEG-guided cybersickness mitigation through adaptive adjustment of VR content factors," *Displays*, vol. 83, p. 102704, Jul. 2024.
- [11] H. Yamamura, H. Baldauf, and K. Kunze, "HemodynamicVR - Adapting the User's Field Of View during Virtual Reality Locomotion Tasks to Reduce Cybersickness using Wearable Functional Near-Infrared Spectroscopy," in *Proceedings of the Augmented Humans International Conference 2021*, ser. AHs '21. New York, NY, USA: Association for Computing Machinery, Jul. 2021, pp. 223–227.
- [12] J. R. Lackner, "Motion sickness: More than nausea and vomiting," *Experimental Brain Research*, vol. 232, no. 8, pp. 2493–2510, Aug. 2014.
- [13] B. Keshavarz and J. F. Golding, "Motion sickness: Current concepts and management," *Current Opinion in Neurology*, vol. 35, no. 1, pp. 107–112, Feb. 2022.
- [14] R. S. Kennedy, J. Drexler, and R. C. Kennedy, "Research in visually induced motion sickness," *Applied Ergonomics*, vol. 41, no. 4, pp. 494–503, Jul. 2010.
- [15] Y.-H. Cha, J. F. Golding, B. Keshavarz, J. Furman, J.-S. Kim, J. A. Lopez-Escamez, M. Magnusson, B. J. Yates, and B. D. Lawson, "Motion sickness diagnostic criteria: Consensus Document of the Classification Committee of the Bárány Society," *Journal of Vestibular Research*, vol. 31, no. 5, pp. 327–344, 2021.
- [16] B. Keshavarz, H. Hecht, and B. D. Lawson, "Visually Induced Motion Sickness: Causes, Characteristics, and Countermeasures," in *Handbook of Virtual Environments: Design, Implementation, and Applications, Second Edition*, 2014, vol. 34, pp. 1–3.
- [17] K. M. Stanney, R. S. Kennedy, and J. M. Drexler, "Cybersickness is Not Simulator Sickness," *Proceedings of the Human Factors and Ergonomics Society Annual Meeting*, vol. 41, no. 2, pp. 1138–1142, Oct. 1997.
- [18] L. Rebenitsch and C. Owen, "Review on cybersickness in applications and visual displays," *Virtual Reality*, vol. 20, no. 2, pp. 101–125, Jun. 2016.
- [19] B. Keshavarz and H. Hecht, "Validating an efficient method to quantify motion sickness," *Human Factors*, vol. 53, no. 4, pp. 415–426, Aug. 2011.
- [20] J. T. Reason and J. J. Brand, *Motion Sickness*. London: Acad. Pr, 1975.
- [21] J. T. Reason, "Motion Sickness Adaptation: A Neural Mismatch Model," *Journal of the Royal Society of Medicine*, vol. 71, no. 11, pp. 819–829, Nov. 1978.
- [22] C. M. Oman, "Motion sickness: A synthesis and evaluation of the sensory conflict theory," *Canadian Journal of Physiology and Pharmacology*, vol. 68, no. 2, pp. 294–303, Feb. 1990.
- [23] W. Bles, J. E. Bos, B. De Graaf, E. Groen, and A. H. Wertheim, "Motion sickness: Only one provocative conflict?" *Brain Research Bulletin*, vol. 47, no. 5, pp. 481–487, Nov. 1998.
- [24] J. Prothero and D. Parker, "A unified approach to presence and motion sickness," in *Virtual and Adaptive Environments: Applications, Implications, and Human Performance Issues*. Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers, 2003, pp. 47–66.
- [25] G. E. Riccio and T. A. Stoffregen, "An ecological Theory of Motion Sickness and Postural Instability," *Ecological Psychology*, vol. 3, no. 3, pp. 195–240, Sep. 1991.

- [26] T. A. Stoffregen and L. Smart, "Postural instability precedes motion sickness," *Brain Research Bulletin*, vol. 47, no. 5, pp. 437–448, Nov. 1998.
- [27] C. A. T. Cortes, C.-T. Lin, T.-T. N. Do, and H.-T. Chen, "An EEG-based Experiment on VR Sickness and Postural Instability While Walking in Virtual Environments," Feb. 2023.
- [28] J. Teixeira, S. Mielliet, and S. Palmisano, "Unexpected Vection Exacerbates Cybersickness During HMD-Based Virtual Reality," *Frontiers in Virtual Reality*, vol. 3, Apr. 2022.
- [29] —, "Investigating the Relative Contributions of Unexpected Vection and Postural Instability to VR Cybersickness," *International Journal of Human-Computer Interaction*, pp. 1–17, Apr. 2024.
- [30] G. Kramida, "Resolving the Vergence-Accommodation Conflict in Head-Mounted Displays," *IEEE Transactions on Visualization and Computer Graphics*, vol. 22, no. 7, pp. 1912–1931, Jul. 2016.
- [31] S. M. Ebenholtz, "Motion sickness and oculomotor systems in virtual environments," *Presence: Teleoper. Virtual Environ.*, vol. 1, no. 3, pp. 302–305, Jan. 1992.
- [32] M. Treisman, "Motion Sickness: An Evolutionary Hypothesis," *Science*, vol. 197, no. 4302, pp. 493–495, Jul. 1977.
- [33] M. Nürnberg, C. Klingner, O. W. Witte, and S. Brodoehl, "Mismatch of Visual-Vestibular Information in Virtual Reality: Is Motion Sickness Part of the Brains Attempt to Reduce the Prediction Error?" *Frontiers in Human Neuroscience*, vol. 15, p. 757735, Oct. 2021.
- [34] E. Krokos and A. Varshney, "Quantifying VR cybersickness using EEG," *Virtual Reality*, vol. 26, no. 1, pp. 77–89, Mar. 2022.
- [35] F. Scholkmann, S. Kleiser, A. J. Metz, R. Zimmermann, J. Mata Pavia, U. Wolf, and M. Wolf, "A review on continuous wave functional near-infrared spectroscopy and imaging instrumentation and methodology," *NeuroImage*, vol. 85 Pt 1, pp. 6–27, Jan. 2014.
- [36] P. Pinti, I. Tachtsidis, A. Hamilton, J. Hirsch, C. Aichelburg, S. Gilbert, and P. W. Burgess, "The present and future use of functional near-infrared spectroscopy (fNIRS) for cognitive neuroscience," *Annals of the New York Academy of Sciences*, vol. 1464, no. 1, pp. 5–29, Mar. 2020.
- [37] S. S. Yeo, S. Y. Park, and S. H. Yun, "Investigating cortical activity during cybersickness by fNIRS," *Scientific Reports*, vol. 14, no. 1, p. 8093, Apr. 2024.
- [38] H. Yamamura, H. Baldauf, and K. Kunze, "Pleasant Locomotion – Towards Reducing Cybersickness using fNIRS during Walking Events in VR," in *Adjunct Publication of the 33rd Annual ACM Symposium on User Interface Software and Technology*. Virtual Event USA: ACM, Oct. 2020, pp. 56–58.
- [39] T. Tanimura, H. Takada, A. Sugiura, F. Kinoshita, and M. Takada, "Effects of The Low-Resolution 3D Video Clip on Cerebrum Blood Flow Dynamics," *Advances in Science, Technology and Engineering Systems Journal*, vol. 4, no. 2, pp. 380–386, 2019.
- [40] A. Benelli, F. Neri, A. Cinti, P. Pasqualetti, S. M. Romanella, A. Giannotta, D. De Monte, M. Mandalà, C. Smeralda, D. Prattichizzo, E. Santarnecchi, and S. Rossi, "Frequency-Dependent Reduction of Cybersickness in Virtual Reality by Transcranial Oscillatory Stimulation of the Vestibular Cortex," *Neurotherapeutics*, vol. 20, no. 6, pp. 1796–1807, Oct. 2023.
- [41] A. Abdalhadi, N. Koundal, R. Lou, M. S. Moosavi, F. Merienne, M. Z. Yusoff, and N. M. Saad, "Development of Virtual Environment to Enhance User Experience with the Help of Electroencephalography," in *Proceedings of the International Conference on E-Mobility—Volume 2*, R. Kannan and T. I. Mohamad, Eds. Singapore: Springer Nature, 2025, pp. 81–91.
- [42] M.-H. Ahn, J. H. Park, H. Jeon, H.-J. Lee, H.-J. Kim, and S. K. Hong, "Temporal Dynamics of Visually Induced Motion Perception and Neural Evidence of Alterations in the Motion Perception Process in an Immersive Virtual Reality Environment," *Frontiers in Neuroscience*, vol. 14, p. 600839, Nov. 2020.
- [43] P. Andrievskaia, S. Berti, J. Spaniol, and B. Keshavarz, "Exploring neurophysiological correlates of visually induced motion sickness using electroencephalography (EEG)," *Experimental Brain Research*, vol. 241, no. 10, pp. 2463–2473, Oct. 2023.
- [44] J. K. Argasiński, N. Lipp, and S. Mazurek, "Electroencephalographic (EEG) Correlates of Visually Induced Motion Sickness (VIMS) in the Virtual Reality (VR) Based Simulations," in *Human-Computer Interaction – INTERACT 2023*, J. Abdelnour Nocera, M. Kristín Lárusdóttir, H. Petrie, A. Piccinno, and M. Winckler, Eds. Cham: Springer Nature Switzerland, 2023, vol. 14145, pp. 59–67.
- [45] C. Balaji, S. Mishra, and S. K. Satapathy, "Neurodynamic Mapping for Cybersickness Prediction in 3D Game Immersion," in *2025 International Conference on Emerging Systems and Intelligent Computing (ESIC)*. Bhubaneswar, India: IEEE, Feb. 2025, pp. 1–6.
- [46] L. M. Berger, G. Wood, C. Neuper, and S. E. Kober, "Sex Differences in User Experience in a VR EEG Neurofeedback Paradigm," in *Games and Learning Alliance*, F. De Rosa, I. Marfisi Schottman, J. Baalsrud Hauge, F. Bellotti, P. Dondio, and M. Romero, Eds. Cham: Springer International Publishing, 2021, vol. 13134, pp. 111–120.
- [47] L. M. Berger, G. Wood, and S. E. Kober, "Influence of a placebo tDCS treatment on cybersickness and EEG-neurofeedback success," *Behavioural Brain Research*, vol. 465, p. 114917, May 2024.
- [48] —, "Manipulating cybersickness in virtual reality-based neurofeedback and its effects on training performance," *Journal of Neural Engineering*, vol. 22, no. 2, p. 026014, Apr. 2025.
- [49] L. Chai, C. Hua, Z. Zhou, X. Chen, and J. Tao, "An EEG Study of Virtual Reality Motion Sickness Based on MVMD Combined with Entropy Asymmetry," in *Proceedings of 2023 Chinese Intelligent Automation Conference*, Z. Deng, Ed. Singapore: Springer Nature Singapore, 2023, vol. 1082, pp. 372–377.
- [50] E. Chang, I. Hwang, H. Jeon, Y. Chun, H. T. Kim, and C. Park, "Effects of rest frames on cybersickness and oscillatory brain activity," in *2013 International Winter Workshop on Brain-Computer Interface (BCI)*. Gangwon Province, Korea (South): IEEE, Feb. 2013, pp. 62–64.
- [51] E. Chang, H. T. Kim, and B. Yoo, "Identifying physiological correlates of cybersickness using heartbeat-evoked potential analysis," *Virtual Reality*, vol. 26, no. 3, pp. 1193–1205, Sep. 2022.
- [52] Y.-C. Chen, J.-R. Duann, C.-L. Lin, S.-W. Chuang, T.-P. Jung, and C.-T. Lin, "Motion-Sickness Related Brain Areas and EEG Power Activates," in *Foundations of Augmented Cognition. Neuroergonomics and Operational Neuroscience*, D. D. Schmorow, I. V. Estabrooke, and M. Grootjen, Eds. Berlin, Heidelberg: Springer Berlin Heidelberg, 2009, vol. 5638, pp. 348–354.
- [53] Y.-C. Chen, J.-R. Duann, S.-W. Chuang, C.-L. Lin, L.-W. Ko, T.-P. Jung, and C.-T. Lin, "Spatial and temporal EEG dynamics of motion sickness," *NeuroImage*, vol. 49, no. 3, pp. 2862–2870, Feb. 2010.
- [54] F. Chiassi, Y. Weiss, T. Steinbrecher, C. Mai, and T. Kosch, "Mind the Visual Discomfort: Assessing Event-Related Potentials as Indicators for Visual Strain in Head-Mounted Displays," in *2024 IEEE International Symposium on Mixed and Augmented Reality (ISMAR)*. Bellevue, WA, USA: IEEE, Oct. 2024, pp. 150–159.
- [55] Y. Dagdemir, "Classification of Emotional and Immersive Outcomes in the Context of Virtual Reality Scene Interactions," *Diagnostics*, vol. 13, no. 22, p. 3437, Nov. 2023.
- [56] —, "Locomotion techniques with EEG signals in a virtual reality environment," *Displays*, vol. 80, p. 102538, Dec. 2023.
- [57] Ö. S. Emeksiz, E. Erzincan, Y. Yemez, and T. M. Sezgin, "A Convolutional Transformer Model for EEG-Driven Cybersickness Detection in VR Experience," in *2025 33rd Signal Processing and Communications Applications Conference (SIU)*. Sile, Istanbul, Türkiye: IEEE, Jun. 2025, pp. 1–4.
- [58] N. Feng, B. Zhou, Q. Zhang, C. Hua, and Y. Yuan, "A comprehensive exploration of motion sickness process analysis from EEG signal and virtual reality," *Computer Methods and Programs in Biomedicine*, vol. 264, p. 108714, Jun. 2025.
- [59] M. Gallagher, R. Dowsett, and E. R. Ferrè, "Vection in virtual reality modulates vestibular-evoked myogenic potentials," *European Journal of Neuroscience*, vol. 50, no. 10, pp. 3557–3565, Nov. 2019.
- [60] C. Groth, J.-P. Tauscher, N. Heesen, M. Hattenbach, S. Castillo, and M. Magnor, "Omnidirectional Galvanic Vestibular Stimulation in Virtual Reality," *IEEE Transactions on Visualization and Computer Graphics*, vol. 28, no. 5, pp. 2234–2244, May 2022.
- [61] C. Hua, L. Chai, Y. Yan, J. Liu, Q. Wang, R. Fu, and Z. Zhou, "Assessment of Virtual Reality Motion Sickness Severity Based on EEG via LSTM/BiLSTM," *IEEE Sensors Journal*, vol. 23, no. 20, pp. 24 839–24 848, Oct. 2023.
- [62] C. Hua, J. Tao, Z. Zhou, L. Chai, Y. Yan, J. Liu, and R. Fu, "EEG classification model for virtual reality motion sickness based on multi-scale CNN feature correlation," *Computer Methods and Programs in Biomedicine*, vol. 251, p. 108218, Jun. 2024.
- [63] C. Hua, L. Chai, Z. Zhou, J. Tao, Y. Yan, X. Chen, J. Liu, and R. Fu, "Detection of virtual reality motion sickness based on EEG using asymmetry of entropy and cross-frequency coupling," *Physiology & Behavior*, vol. 284, p. 114626, Oct. 2024.
- [64] C. Hua, Z. Zhou, J. Tao, Y. Yan, J. Liu, R. Fu, and L. Chai, "A Siam-LSTM Model for Multichannel EEG on VR Motion Sickness Recognition," *IEEE Sensors Journal*, vol. 24, no. 21, pp. 35 245–35 254, Nov. 2024.

- [65] C. Hua, Y. Chen, J. Tao, Z. Dai, W. Yang, D. Chen, J. Liu, and R. Fu, "Dual-pathway EEG model with channel attention for virtual reality motion sickness detection," *Journal of Neuroscience Methods*, vol. 418, p. 110425, Jun. 2025.
- [66] K.-M. Jang, M. Kwon, S. G. Nam, D. Kim, and H. K. Lim, "Estimating objective (EEG) and subjective (SSQ) cybersickness in people with susceptibility to motion sickness," *Applied Ergonomics*, vol. 102, p. 103731, Jul. 2022.
- [67] D. Jeong, S. Yoo, and J. Yun, "Cybersickness Analysis with EEG Using Deep Learning Algorithms," in *2019 IEEE Conference on Virtual Reality and 3D User Interfaces (VR)*. Osaka, Japan: IEEE, Mar. 2019, pp. 827–835.
- [68] Khaitami, A. D. Wibawa, S. Mardi, S. Nugroho, and A. Z. Khoirunnisaa, "EEG Visualization for Cybersickness Detection During Playing 3D Video Games," in *2019 International Seminar on Intelligent Technology and Its Applications (ISITIA)*. Surabaya, Indonesia: IEEE, Aug. 2019, pp. 325–330.
- [69] Y. Kim, H. Kim, H. Ko, and H. Kim, "Psychophysiological changes by navigation in a virtual reality," in *2001 Conference Proceedings of the 23rd Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, vol. 4, Oct. 2001, pp. 3773–3776 vol.4.
- [70] Y. Y. Kim, H. J. Kim, E. N. Kim, H. D. Ko, and H. T. Kim, "Characteristic changes in the physiological components of cybersickness," *Psychophysiology*, vol. 42, no. 5, pp. 616–625, 2005.
- [71] J. Kim, W. Kim, H. Oh, S. Lee, and S. Lee, "A Deep Cybersickness Predictor Based on Brain Signal Analysis for Virtual Reality Contents," in *2019 IEEE/CVF International Conference on Computer Vision (ICCV)*. Seoul, Korea (South): IEEE, Oct. 2019, pp. 10 579–10 588.
- [72] J.-Y. Kim, J.-B. Son, H.-S. Leem, and S.-H. Lee, "Psychophysiological Alteration After Virtual Reality Experiences Using Smartphone-Assisted Head Mount Displays: An EEG-Based Source Localization Study," *Applied Sciences*, vol. 9, no. 12, p. 2501, Jun. 2019.
- [73] L.-W. Ko, C.-S. Wei, T.-P. Jung, and C.-T. Lin, "Estimating the Level of Motion Sickness Based on EEG Spectra," in *Foundations of Augmented Cognition. Directing the Future of Adaptive Systems*, D. D. Schmorow and C. M. Fidopiastis, Eds. Berlin, Heidelberg: Springer Berlin Heidelberg, 2011, vol. 6780, pp. 169–176.
- [74] L.-W. Ko, H.-C. Lee, S.-F. Tsai, T.-C. Shih, Y.-T. Chuang, H.-L. Huang, S.-Y. Ho, and C.-T. Lin, "EEG-based motion sickness classification system with genetic feature selection," in *2013 IEEE Symposium on Computational Intelligence, Cognitive Algorithms, Mind, and Brain (CCMB)*. Singapore, Singapore: IEEE, Apr. 2013, pp. 158–164.
- [75] S. Lee, S. Kim, H. G. Kim, and Y. M. Ro, "Assessing Individual VR Sickness Through Deep Feature Fusion of VR Video and Physiological Response," *IEEE Transactions on Circuits and Systems for Video Technology*, vol. 32, no. 5, pp. 2895–2907, May 2022.
- [76] Y. Li, A. Liu, and L. Ding, "Machine learning assessment of visually induced motion sickness levels based on multiple biosignals," *Biomedical Signal Processing and Control*, vol. 49, pp. 202–211, Mar. 2019.
- [77] X. Li, C. Zhu, C. Xu, J. Zhu, Y. Li, and S. Wu, "VR motion sickness recognition by using EEG rhythm energy ratio based on wavelet packet transform," *Computer Methods and Programs in Biomedicine*, vol. 188, p. 105266, May 2020.
- [78] G. Li, F. M. Varela, A. Habib, Q. Zhang, M. McGill, S. Brewster, and F. Pollick, "Exploring the feasibility of mitigating VR-HMD-induced cybersickness using cathodal transcranial direct current stimulation," in *2020 IEEE International Conference on Artificial Intelligence and Virtual Reality (AIVR)*. Utrecht, Netherlands: IEEE, Dec. 2020, pp. 123–129.
- [79] G. Li, O. Onuoha, M. McGill, S. Brewster, C. P. Chen, and F. Pollick, "Comparing Autonomic Physiological and Electroencephalography Features for VR Sickness Detection Using Predictive Models," in *2021 IEEE Symposium Series on Computational Intelligence (SSCI)*. Orlando, FL, USA: IEEE, Dec. 2021, pp. 01–08.
- [80] G. Li, K. Pohlmann, M. McGill, C. P. Chen, S. Brewster, and F. Pollick, "Exploring Neural Biomarkers in Young Adults Resistant to VR Motion Sickness: A Pilot Study of EEG," in *2023 IEEE Conference Virtual Reality and 3D User Interfaces (VR)*. Shanghai, China: IEEE, Mar. 2023, pp. 328–335.
- [81] G. Li, Y.-K. Wang, M. McGill, K. Pöhlmann, S. Brewster, and F. Pollick, "Resting-state EEG in the Vestibular Region Can Predict Motion Sickness Induced by a Motion-Simulated in-car VR Platform," in *2023 IEEE Symposium Series on Computational Intelligence (SSCI)*, Dec. 2023, pp. 47–52.
- [82] G. Li and T. Zanto, "Reduced VR motion sickness by applying random-phase transcranial alternating current stimulation to the left parietal cortex," *Brain Stimulation*, vol. 17, no. 3, pp. 550–552, May 2024.
- [83] C.-Y. Liao, S.-K. Tai, R.-C. Chen, and H. Hendry, "Using EEG and Deep Learning to Predict Motion Sickness Under Wearing a Virtual Reality Device," *IEEE Access*, vol. 8, pp. 126 784–126 796, 2020.
- [84] H. K. Lim, K. Ji, Y. S. Woo, D.-u. Han, D.-H. Lee, S. G. Nam, and K.-M. Jang, "Test-retest reliability of the virtual reality sickness evaluation using electroencephalography (EEG)," *Neuroscience Letters*, vol. 743, p. 135589, Jan. 2021.
- [85] C.-T. Lin, S.-W. Chuang, Y.-C. Chen, L.-W. Ko, S.-F. Liang, and T.-P. Jung, "EEG Effects of Motion Sickness Induced in a Dynamic Virtual Reality Environment," in *2007 29th Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, Aug. 2007, pp. 3872–3875.
- [86] C.-T. Lin, S.-F. Tsai, H.-C. Lee, H.-L. Huang, S.-Y. Ho, and L.-W. Ko, "Motion sickness estimation system," in *The 2012 International Joint Conference on Neural Networks (IJCNN)*. Brisbane, Australia: IEEE, Jun. 2012, pp. 1–6.
- [87] Y.-T. Lin, Y.-Y. Chien, H.-H. Wang, F.-C. Lin, and Y.-P. Huang, "65-3: The Quantization of Cybersickness Level Using EEG and ECG for Virtual Reality Head-Mounted Display," *SID Symposium Digest of Technical Papers*, vol. 49, no. 1, pp. 862–865, May 2018.
- [88] R. Liu, E. Peli, and A. D. Hwang, "Measuring visually induced motion sickness using wearable devices," *Electronic Imaging*, vol. 29, no. 14, pp. 218–223, Jan. 2017.
- [89] R. Liu, M. Xu, Y. Zhang, E. Peli, and A. D. Hwang, "A Pilot Study on EEG-Based Evaluation of Visually Induced Motion Sickness," *The Journal of imaging science and technology*, vol. 64, no. 2, pp. 20 501–1–20 501–10, Mar. 2020.
- [90] M. Liu, B. Yang, M. Xu, P. Zan, L. Chen, and X. Xia, "Exploring quantitative assessment of cybersickness in virtual reality using EEG signals and a CNN-ECA-LSTM network," *Displays*, vol. 81, p. 102602, Jan. 2024.
- [91] M. Liu, B. Yang, P. Zan, L. Chen, B. Wang, and X. Xia, "Exploring the brain physiological activity and quantified assessment of VR cybersickness using EEG signals," *Displays*, vol. 85, p. 102879, Dec. 2024.
- [92] S. Luo, P. Ren, J. Wu, X. Wu, and X. Zhang, "Feature extraction method of EEG based on wavelet packet reconstruction and deep learning model of VR motion sickness feature classification and prediction," *PLOS ONE*, vol. 19, no. 7, p. e0305733, Jul. 2024.
- [93] M. Mawalid, A. Z. Khoirunnisa, M. H. Purnomo, and A. D. Wibawa, "Classification of EEG Signal for Detecting Cybersickness through Time Domain Feature Extraction using Naïve Bayes," in *2018 International Conference on Computer Engineering, Network and Intelligent Multimedia (CENIM)*, Nov. 2018, pp. 29–34.
- [94] M.-A. Moïnnereau and T. H. Falk, "Cybersickness Marker Prediction Using a Biosensors-Instrumented VR Headset: A Pilot Study," in *2024 IEEE 4th International Conference on Human-Machine Systems (ICHMS)*. Toronto, ON, Canada: IEEE, May 2024, pp. 1–6.
- [95] Y. Nakata, M. Otsuki, M. Sakai, M. Nagai, R. Yamamoto, and A. Aoyama, "Exploring Display Parameters Associated with Cybersickness Using Electroencephalography," in *2025 IEEE Conference on Virtual Reality and 3D User Interfaces Abstracts and Workshops (VRW)*. Saint Malo, France: IEEE, Mar. 2025, pp. 1390–1391.
- [96] S. Nam, K.-M. Jang, M. Kwon, H. K. Lim, and J. Jeong, "Electroencephalogram microstates and functional connectivity of cybersickness," *Frontiers in Human Neuroscience*, vol. 16, p. 857768, Aug. 2022.
- [97] W. Nunes Da Silva, T. M. Porcino, C. D. Castanho, and R. P. Jacobi, "Analysis of Cybersickness through Biosignals: An approach with Symbolic Machine Learning," in *Symposium on Virtual and Augmented Reality*. Manaus Brazil: ACM, Sep. 2024, pp. 11–20.
- [98] A. Ozkan, U. Uyan, and U. Celikcan, "Effects of speed, complexity and stereoscopic VR cues on cybersickness examined via EEG and self-reported measures," *Displays*, vol. 78, p. 102415, Jul. 2023.
- [99] E. S. Pane, A. Zuhriya Khoirunnisaa, A. D. Wibawa, and M. H. Purnomo, "Identifying Severity Level of Cybersickness from EEG signals using CN2 Rule Induction Algorithm," in *2018 International Conference on Intelligent Informatics and Biomedical Sciences (ICIIBMS)*, vol. 3, Oct. 2018, pp. 170–176.
- [100] G. N. Pradhan, R. C. Galvan-Garza, A. M. Perez, J. Stepanek, and M. J. Cevette, "Visual Vestibular Conflict Mitigation in Virtual Reality Using Galvanic Vestibular Stimulation," *Aerospace Medicine and Human Performance*, vol. 93, no. 5, pp. 406–414, May 2022.

- [101] R.-R. Qi, S.-F. Xiao, L.-L. Pan, Y.-Q. Mao, Y. Su, L.-J. Wang, and Y.-L. Cai, "Profiling of cybersickness and balance disturbance induced by virtual ship motion immersion combined with galvanic vestibular stimulation," *Applied Ergonomics*, vol. 92, p. 103312, Apr. 2021.
- [102] M. Recenti, C. Ricciardi, R. Aubonnet, I. Picone, D. Jacob, H. Å. R. Svansson, S. Agnarsdóttir, G. H. Karlsson, V. Baeringsdóttir, H. Petersen, and P. Gargiulo, "Toward Predicting Motion Sickness Using Virtual Reality and a Moving Platform Assessing Brain, Muscles, and Heart Signals," *Frontiers in Bioengineering and Biotechnology*, vol. 9, Apr. 2021.
- [103] O. Rosanne, D. Benesch, G. Kratzig, S. Paré, N. Bolt, and T. H. Falk, "To pre-process or not to pre-process? On the role of EEG enhancement for cybersickness characterization and the importance of amplitude modulation features," *Frontiers in Virtual Reality*, vol. 6, p. 1468971, Jan. 2025.
- [104] J. Sameri, H. Coenegracht, S. Van Damme, F. De Turck, and M. Torres Vega, "Physiology-driven cybersickness detection in virtual reality: A machine learning and explainable AI approach," *Virtual Reality*, vol. 28, no. 4, p. 174, Nov. 2024.
- [105] Z. Shen, X. Liu, W. Li, X. Li, and Q. Wang, "Classification of Visually Induced Motion Sickness Based on Phase-Locked Value Functional Connectivity Matrix and CNN-LSTM," *Sensors*, vol. 24, no. 12, p. 3936, Jun. 2024.
- [106] M. Sra, A. Jain, and P. Maes, "Adding Proprioceptive Feedback to Virtual Reality Experiences Using Galvanic Vestibular Stimulation," in *Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems*. Glasgow Scotland UK: ACM, May 2019, pp. 1–14.
- [107] G. R. F. Suwandi, S. N. Khotimah, F. Haryanto, and Suprijadi, "Electroencephalography Signal Analysis for Virtual Reality Sickness: Head-mounted Display and Screen-based," *International Journal on Advanced Science, Engineering and Information Technology*, vol. 13, no. 4, pp. 1449–1455, Aug. 2023.
- [108] G. R. F. Suwandi, S. N. Khotimah, F. Haryanto, and Suprijadi, "Exploring the Impact of Motion Parameter Variations in Virtual Reality Content on Visually Induced Motion Sickness: An Electroencephalography Signal Analysis Approach," *International Journal of Online and Biomedical Engineering (iJOE)*, vol. 20, no. 12, pp. 160–176, Sep. 2024.
- [109] N. Takeuchi, T. Mori, Y. Suzukamo, and S.-I. Izumi, "Modulation of Excitability in the Temporoparietal Junction Relieves Virtual Reality Sickness," *Cyberpsychology, Behavior, and Social Networking*, vol. 21, no. 6, pp. 381–387, Jun. 2018.
- [110] N. Tian and R. Boulic, "Who says you are so sick? An investigation on individual susceptibility to cybersickness triggers using EEG, EGG and ECG," *IEEE Transactions on Visualization and Computer Graphics*, pp. 1–11, 2024.
- [111] K. Watanabe, K. Onishi, M. Koeda, M. Tsujishita, and H. Noborio, "Study of VR Sickness Reduction Effect by Tunneling Effect According to the Gazing Point," in *Human-Computer Interaction*, M. Kurosu and A. Hashizume, Eds. Cham: Springer Nature Switzerland, 2025, vol. 15767, pp. 446–460.
- [112] S. Weech, J. Moon, and N. F. Troje, "Influence of bone-conducted vibration on simulator sickness in virtual reality," *PLOS ONE*, vol. 13, no. 3, p. e0194137, Mar. 2018.
- [113] S. Weech, T. Wall, and M. Barnett-Cowan, "Reduction of cybersickness during and immediately following noisy galvanic vestibular stimulation," *Experimental Brain Research*, vol. 238, no. 2, pp. 427–437, Feb. 2020.
- [114] C.-S. Wei, L.-W. Ko, S.-W. Chuang, T.-P. Jung, and C.-T. Lin, "EEG-based evaluation system for motion sickness estimation," in *2011 5th International IEEE/EMBS Conference on Neural Engineering*. Cancun: IEEE, Apr. 2011, pp. 100–103.
- [115] Y. S. Woo, K.-M. Jang, S. G. Nam, M. Kwon, and H. K. Lim, "Recovery time from VR sickness due to susceptibility: Objective and quantitative evaluation using electroencephalography," *Heliyon*, vol. 9, no. 4, p. e14792, Apr. 2023.
- [116] J. Wu, Q. Zhou, J. Li, X. Kong, and Y. Xiao, "Inhibition-related N2 and P3: Indicators of visually induced motion sickness (VIMS)," *International Journal of Industrial Ergonomics*, vol. 78, p. 102981, Jul. 2020.
- [117] W. Xu, H.-N. Liang, Y. Yu, D. Monteiro, K. Hasan, and C. Fleming, "Assessing the effects of a full-body motion-based exergame in virtual reality," in *Proceedings of the Seventh International Symposium of Chinese CHI*. Xiamen China: ACM, Jun. 2019, pp. 1–6.
- [118] M. Xu, B. Yang, M. Liu, Y. Huang, X. Xia, and J. Zhang, "SE-CNN Attention Structure for Quantitative EEG-Based Assessment of VR Motion Sickness," in *Proceedings of the 2023 9th International Conference on Computing and Artificial Intelligence*, ser. ICCAI '23. New York, NY, USA: Association for Computing Machinery, Aug. 2023, pp. 606–611.
- [119] A. H. X. Yang, N. K. Kasabov, and Y. O. Cakmak, "Prediction and detection of virtual reality induced cybersickness: A spiking neural network approach using spatiotemporal EEG brain data and heart rate variability," *Brain Informatics*, vol. 10, no. 1, p. 15, Dec. 2023.
- [120] S. S. Yeo, J. W. Kwon, and S. Y. Park, "EEG-based analysis of various sensory stimulation effects to reduce visually induced motion sickness in virtual reality," *Scientific Reports*, vol. 12, no. 1, p. 18043, Oct. 2022.
- [121] Z. Zhou, C. Hua, L. Chai, and J. Tao, "Application of EEG S-Transformation Combined with Dimensionless Metrics for Automatic Detection of Cybersickness," in *Proceedings of 2023 Chinese Intelligent Automation Conference*, Z. Deng, Ed. Singapore: Springer Nature Singapore, 2023, vol. 1082, pp. 634–640.
- [122] J. J. Newton and T. C. Thiagarajan, "EEG Frequency Bands in Psychiatric Disorders: A Review of Resting State Studies," *Frontiers in Human Neuroscience*, vol. 12, p. 521, 2018.
- [123] A. Z. Khoirunnisa, E. S. Pane, A. D. Wibawa, and M. H. Purnomo, "Channel Selection of EEG-Based Cybersickness Recognition during Playing Video Game Using Correlation Feature Selection (CFS)," in *2018 2nd International Conference on Biomedical Engineering (IBIOMED)*, Jul. 2018, pp. 48–53.
- [124] Z. Li, L. Zhao, J. Chang, W. Li, M. Yang, C. Li, R. Wang, and L. Ji, "EEG-based evaluation of motion sickness and reducing sensory conflict in a simulated autonomous driving environment," in *2022 44th Annual International Conference of the IEEE Engineering in Medicine & Biology Society (EMBC)*, Jul. 2022, pp. 4026–4030.
- [125] Y. Wei, Y. O. Okazaki, R. H. So, W. C. Chu, and K. Kitajo, "Motion sickness-susceptible participants exposed to coherent rotating dot patterns show excessive N2 amplitudes and impaired theta-band phase synchronization," *NeuroImage*, vol. 202, p. 116028, Nov. 2019.
- [126] Y.-H. Yu, P.-C. Lai, L.-W. Ko, C.-H. Chuang, B.-C. Kuo, and C.-T. Lin, "An EEG-based classification system of Passenger's motion sickness level by using feature extraction/selection technologies," in *The 2010 International Joint Conference on Neural Networks (IJCNN)*, Jul. 2010, pp. 1–6.
- [127] C.-S. Wei, L.-W. Ko, S.-W. Chuang, T.-P. Jung, and C.-T. Lin, "Genetic feature selection in EEG-based motion sickness estimation," in *The 2011 International Joint Conference on Neural Networks*, Jul. 2011, pp. 365–369.
- [128] C.-S. Wei, S.-W. Chuang, W.-R. Wang, L.-W. Ko, T.-P. Jung, and C.-T. Lin, "Implementation of a motion sickness evaluation system based on EEG spectrum analysis," in *2011 IEEE International Symposium of Circuits and Systems (ISCAS)*, May 2011, pp. 1081–1084.
- [129] J.-R. Park, D.-W. Lim, S.-Y. Lee, H.-W. Lee, M.-H. Choi, and S.-C. Chung, "Long-Term Study of Simulator Sickness: Differences in EEG Response Due to Individual Sensitivity," *International Journal of Neuroscience*, vol. 118, no. 6, pp. 857–865, Jun. 2008.
- [130] S. Palmisano, D. Apthorp, T. Seno, and P. J. Stapley, "Spontaneous postural sway predicts the strength of smooth vection," *Experimental Brain Research*, vol. 232, no. 4, pp. 1185–1191, Apr. 2014.
- [131] P. Kourtesis, J. Linnell, R. Amir, F. Argelaguet, and S. E. MacPherson, "Cybersickness in Virtual Reality Questionnaire (CSQ-VR): A Validation and Comparison against SSQ and VRSQ," *Virtual Worlds*, vol. 2, no. 1, pp. 16–35, Mar. 2023.
- [132] P. Kourtesis, R. Amir, J. Linnell, F. Argelaguet, and S. E. MacPherson, "Cybersickness, Cognition, & Motor Skills: The Effects of Music, Gender, and Gaming Experience," *IEEE Transactions on Visualization and Computer Graphics*, vol. 29, no. 5, pp. 2326–2336, May 2023.
- [133] T. Saito and M. Rehmsmeier, "The Precision-Recall Plot Is More Informative than the ROC Plot When Evaluating Binary Classifiers on Imbalanced Datasets," *PLOS ONE*, vol. 10, no. 3, p. e0118432, Mar. 2015.
- [134] E. Combrisson and K. Jerbi, "Exceeding chance level by chance: The caveat of theoretical chance levels in brain signal classification and statistical assessment of decoding accuracy," *Journal of Neuroscience Methods*, vol. 250, pp. 126–136, Jul. 2015.
- [135] G. Varoquaux, "Cross-validation failure: Small sample sizes lead to large error bars," *NeuroImage*, vol. 180, no. Pt A, pp. 68–77, Oct. 2018.
- [136] H. Berger, "Über das Elektrenkephalogramm des Menschen," *Archiv für Psychiatrie und Nervenkrankheiten*, vol. 87, no. 1, pp. 527–570, Dec. 1929.

- [137] W. Klimesch, P. Sauseng, and S. Hanslmayr, "EEG alpha oscillations: The inhibition-timing hypothesis," *Brain Research Reviews*, vol. 53, no. 1, pp. 63–88, Jan. 2007.
- [138] R. J. Barry, A. R. Clarke, S. J. Johnstone, C. A. Magee, and J. A. Rushby, "EEG differences between eyes-closed and eyes-open resting conditions," *Clinical Neurophysiology*, vol. 118, no. 12, pp. 2765–2773, Dec. 2007.
- [139] C. Zhang, S. Li, Y. Li, S. E. Li, and B. Nie, "Analysis of Motion Sickness Associated Brain Activity Using fNIRS: A Driving Simulator Study," *IEEE Access*, vol. 8, pp. 207 415–207 425, 2020.
- [140] M.-A. Moineau, A. A. Oliveira, and T. H. Falk, "Instrumenting a virtual reality headset for at-home gamer experience monitoring and behavioural assessment," *Frontiers in Virtual Reality*, vol. 3, Oct. 2022.
- [141] J. W. Kelly, S. B. Gilbert, M. C. Dorneich, and K. A. Costabile, "Gender differences in cybersickness: Clarifying confusion and identifying paths forward," in *2023 IEEE Conference on Virtual Reality and 3D User Interfaces Abstracts and Workshops (VRW)*. Shanghai, China: IEEE, Mar. 2023, pp. 283–288.
- [142] B. Lawson and J. Bolkhovsky, "Attention Military/Commercial Simulation Developers, Users, & Trainers: Visually-induced or Motion-induced Sickness is not Necessarily More Severe for Women," in *14th International Conference on Applied Human Factors and Ergonomics (AHFE 2023)*, 2023.
- [143] R. Islam, Y. Lee, M. Jaloli, I. Muhammad, D. Zhu, P. Rad, Y. Huang, and J. Quarles, "Automatic Detection and Prediction of Cybersickness Severity using Deep Neural Networks from user's Physiological Signals," in *2020 IEEE International Symposium on Mixed and Augmented Reality (ISMAR)*, Nov. 2020, pp. 400–411.

Gaël Van der Lee received the B.S. degree in Cognitive Science, with a specialization in Machine Learning and Neural Computation, from the University of California, San Diego, CA, USA, in 2018, and the M.Sc. degree in Artificial Intelligence and Visual Computing from École Polytechnique, Institut Polytechnique de Paris, France, in 2022. He received the Ph.D. degree in Signal Processing and Control Science from the University of Lille, France, in 2025, for work on electroencephalographic markers of self-motion perception and cybersickness in virtual reality, carried out at the Lille Joint Research Center for Informatics, Signal Processing and Control Science (CRISTAL, UMR CNRS 9189) within the European project CHIST-ERA GENESIS. Prior to his doctoral work he was Lead Machine Learning Scientist at Lofty AI, San Francisco, CA, USA. His research interests include electroencephalography, signal processing, and machine learning applied to the estimation of user states in immersive environments, with a focus on vection, acceleration perception, and cybersickness.

François Cabestaing (Senior Member, IEEE) received the B.S. degree in Mathematics in 1987 from USTL, the M.Sc. degree in Engineering and Control Science in 1987 from ISEN, and the PhD degree in Signal Processing and Control Science in 1991 from the University of Lille, France. He holds an Habilitation à Diriger des Recherches obtained in 2003 from ULille. He has been an associate professor at ULille since 1992 and became full professor in 2006 in the same university. He is a researcher at the Lille Joint Research Center for Informatics, Signal Processing and Control Science (CRISTAL, UMR CNRS 9189), where he heads the "BCI" (Brain-Computer Interfaces) research team. He studies physiological signal processing techniques, mainly electroencephalogram, and classification techniques, particularly to estimate the mental states of brain-machine interface users. He contributed to the creation of the Collectif pour la Recherche Transdisciplinaire sur les Interfaces Cerveau-Ordinateur (CORTICO), a learned society of which he was president from 2017 to 2024.

Hakim Si-Mohammed received the M.Sc. and Engineering degrees from the Higher National School of Computer Engineering (ESI), Algiers, Algeria, in 2016, and the Ph.D. degree in Computer Science from Inria and INSA Rennes, France, in 2019. He is currently an Associate Professor with the University of Lille, France, and a researcher at the CRISTAL Laboratory (UMR CNRS 9189). He serves as Deputy Director of the CNRS Research Federation on Visual Sciences and Cultures, and Head of the BCI and Neuroprosthetics Axis at the Institut Cerveau, Société et Technologie. His research interests include Brain-Computer Interfaces and immersive technologies. He led and coordinated the European project CHIST-ERA GENESIS (2021–2025) on cybersickness, and serves as Secretary General of CORTICO, the French Society for Research in Brain-Computer Interfaces.