

1 pyRTX: a Python package for high precision 2 computation of non gravitational forces on spacecraft

3 **Gael Cascioli** ^{1,2*}, **Ariele Zurria** ^{3*}, and **Erwan Mazarico** ²

4 ¹ University of Maryland Baltimore County, United States  ² NASA Goddard Space Flight Center,
5 United States ³ Sapienza University, Italy  Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#))

In partnership with



AMERICAN
ASTRONOMICAL
SOCIETY

This article and software are linked
with research article DOI
[10.3847/xxxxx](https://doi.org/10.3847/xxxxx) <- [update this](#)
[with the DOI from AAS once you](#)
[know it.](#), published in the The
Planetary Science Journal.

6 Summary

7 With the constant improvement of radiometric tracking systems, inaccuracies in the non-
8 gravitational force modeling have become one of the limiting factors to precise orbit deter-
9 mination, and the scientific products that it enables. The main factor impacting the limited
10 accuracy of non-gravitational force models is the complex 3D shape of the spacecraft. While
11 fast, reliable, analytical models are available for simple shapes (spheres, cubes, etc), no such
12 model is generally available for a complex shape. This software package aims to address this
13 limitation by leveraging ray-tracing to compute the complex interaction between the forcing
14 environment (radiation, atmosphere) and the three dimensional shape of the spacecraft.

Statement of need

Several scientific investigations require high-precision reconstruction of spacecraft trajectories.
Among these, one of the most demanding is the determination of the gravity field of Solar
System bodies (planets, moons). This task is accomplished by solving the so-called orbit
determination (OD) problem ([Milani & Gronchi, 2009](#); [Tapley et al., 2004](#)). The solution of
the OD in the adjustment of a dynamical model (a set of differential equations) describing the
spacecraft motion. Systematic errors in the dynamical model will almost inevitably lead to
systematic errors in the solution. In the recent years significant improvements in radiometric
tracking system, have led to more and more precise measurements of the spacecraft position
and velocity (the input to the OD), thus requiring increasingly more accurate dynamical
models ([Asmar et al., 2005](#); [Cappuccio et al., 2020](#); [Mazarico et al., 2023](#)). One of the
major limitations of current dynamical modelling of deep-space probes consists in the complex
interaction between the spacecraft shape and the atmosphere, and with radiative forces (solar
radiation pressure, albedo, thermal infrared radiation). We developed the pyRTX software
package to address this limitation. Leveraging the ray-tracing technique, originally developed
in computer graphics, instead of relying on simplified macro-models (flat plates, cylinders, etc)
pyRTX is able to use the actual 3D shape of the spacecraft (provided as, for example, an .obj
file), to compute several non-gravitational accelerations.

pyRTX addresses the gap in open source solutions for a comprehensive modelling of several,
important, non-gravitational forces. Being built around the SPICE astrodynamics library ([Annex
et al., 2020](#)), pyRTX is intended to be a plug-in tool that can be used in existing OD codes,
and applications.

Functionality

In this section we describe the main functionalities of the pyRTX software. All of these functionalities are discussed in the example library included in the code distribution. Our companion paper (Zurria et al., 2026) discusses an actual application of the pyRTX library for ameliorating the OD of NASA's Lunar Reconnaissance Orbiter.

- *Solar Radiation Pressure Modeling*: pyRTX computes the acceleration due to solar photons by casting rays from a pixel plane representing the incoming solar flux. The engine inherently accounts for self-shadowing (where spacecraft components block light from reaching others) and multiple reflections. Users can specify optical properties for each mesh face, allowing the software to simulate both specular and diffuse (Lambertian) reflections.
- *Planetary Radiation Pressure*: The software models the effects of radiation reflected (albedo) and emitted (thermal infrared) by planetary bodies. This includes the ability to use complex planetary shape models (e.g., topography from digital elevation models, DEMs) and spatially variable maps for albedo, emissivity, and surface temperature.
- *Eclipse and Shadow Function Analysis*: pyRTX can compute precise shadow functions during eclipse transitions. It supports advanced modeling features such as solar limb darkening, where the variation in intensity across the solar disk is integrated into the flux calculation.
- *Atmospheric Drag*: For low-altitude missions, the software calculates the effective aerodynamic cross-section of the spacecraft, providing inputs for atmospheric drag modeling.
- *Lookup Table (LUT) Generation*: To enable efficient integration with OD software, pyRTX can pre-compute accelerations over a grid of incident directions. This feature supports spacecraft with articulating components (e.g., solar arrays), allowing users to generate comprehensive LUTs that map acceleration vectors to specific spacecraft orientations and strongly reduce computation time.

Acknowledgements

Work by G.C. was supported by NASA under award No. 80GSFC24M0006.

References

- Annex, A., Pearson, B., Seignovert, B., Carcich, B., Eichhorn, H., Mapel, J., Von Forstner, J., McAuliffe, J., Del Rio, J., Berry, K., Aye, K.-M., Stefko, M., De Val-Borro, M., Kulumani, S., & Murakami, S. (2020). SpiceyPy: A pythonic wrapper for the SPICE toolkit. *Journal of Open Source Software*, 5(46), 2050. <https://doi.org/10.21105/joss.02050>
- Asmar, S. W., Armstrong, J. W., Iess, L., & Tortora, P. (2005). Spacecraft Doppler tracking: Noise budget and accuracy achievable in precision radio science observations. *Radio Science*, 40(2), n/a–n/a. <https://doi.org/10.1029/2004RS003101>
- Cappuccio, P., Notaro, V., Ruscio, A. di, Iess, L., Genova, A., Durante, D., Stefano, I. di, Asmar, S. W., Ciarcia, S., & Simone, L. (2020). Report on First Inflight Data of BepiColombo's Mercury Orbiter Radio Science Experiment. *IEEE Transactions on Aerospace and Electronic Systems*, 56(6). <https://doi.org/10.1109/TAES.2020.3008577>
- Mazarico, E., Buccino, D., Castillo-Rogez, J., Dombard, A. J., Genova, A., Hussmann, H., Kiefer, W. S., Lunine, J. I., McKinnon, W. B., Nimmo, F., Park, R. S., Roberts, J. H., Srinivasan, D. K., Steinbrügge, G., Tortora, P., & Withers, P. (2023). The Europa

- 89 <https://doi.org/10.0000/pending>