

pyRTX: a Python package for high precision computation of non gravitational forces on deep space probes

Gael Cascioli^{1,2*}, Ariele Zurria^{3*}, and Erwan Mazarico²

¹ University of Maryland Baltimore County, United States ² NASA Goddard Space Flight Center, United States ³ Sapienza University, Italy * Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

Summary

With the constant improvement of radiometric tracking systems, inaccuracies in non-gravitational force modeling have become one of the limiting factors in precise orbit determination for deep space missions and the scientific products that depend on it. A major source of modeling error arises from the complex 3D shape of spacecraft. While fast and reliable analytical models exist for simple geometries (e.g. spheres or flat plates), they are generally not available for more realistic spacecraft representations. We present pyRTX, a software package that addresses this limitation by leveraging ray-tracing techniques to model the interaction between the space environment (radiation and atmosphere) and detailed spacecraft geometries. By operating directly on accurate 3D mesh models, pyRTX enables high-fidelity computation of non-gravitational accelerations. The software is specifically geared towards planetary orbit determination where inaccuracies of the model often cannot be overcome with continuous tracking as it is routinely done for Earth-orbiting spacecraft.

Statement of need

Accurate reconstruction of spacecraft trajectories is essential for several scientific applications, including the determination of the gravity field of Solar System bodies. These tasks are accomplished by solving the orbit determination (OD) problem (Milani & Gronchi, 2009; Tapley et al., 2004). The solution is retrieved by minimizing the difference between observed data and predictions. This is accomplished through an iterative adjustment of a dynamical model (a set of differential equations) describing the spacecraft motion. Systematic errors in the dynamical model directly translate into biases in the estimated trajectory and derived geophysical parameters. In recent years, improvements in radiometric tracking systems have significantly increased measurement precision (Asmar et al., 2005; Cappuccio et al., 2020; Cappuccio et al., 2025; Mazarico et al., 2023), placing tighter requirements on the accuracy of dynamical models.

One of the main limitation in the dynamical modeling of deep space probes, especially in the inner Solar System, is the simplified treatment of non-gravitational forces, such as radiation pressure or atmospheric drag. This occurs as photons or atmospheric particles interact with the probe's surfaces, leading to a momentum exchange and, thus, an acceleration. Existing approaches typically represent the spacecraft shape with simplified macro-models (e.g. flat plates or cylinders). While fast and efficient, these methods cannot fully capture effects such as self-shadowing (when parts of the spacecraft are shielded from the incoming flux by other structural elements) and multiple reflections (reflected particles which impact subsequent surfaces) [Mazarico et al. 2014; Li et al. 2014].

41 The pyRTX software package addresses this gap by implementing a ray-tracing-based approach
 42 for computing non-gravitational accelerations using realistic spacecraft geometries. Built
 43 around the NAIF SPICE library (Acton, 1996) and its Python interface (Annex et al., 2020),
 44 pyRTX is designed to integrate with existing OD pipelines and provide improved dynamical
 45 modeling.

46 State of the field

47 Non-gravitational force modeling in orbit determination has traditionally relied on the simplified
 48 “plate model”, which consists in the discretization of the probe as a set of plates or 2D
 49 elementary shapes. This approach enables efficient computation but lacks the capability to
 50 fully capture the complex interaction between particles and complex spacecraft geometries.
 51 Ray-tracing techniques, widely used in computer graphics, have been explored to address this
 52 task [Darugna et al. 2018; Kenneally & Schaub 2020; Li et al. 2018]. However, to the best of
 53 our knowledge, there is currently no open-source software package that computes solar and
 54 planetary radiation pressure and atmospheric drag with a ray-tracing approach, allowing direct
 55 integration with standard astrodynamics libraries. pyRTX fills this gap by providing a flexible
 56 and open-source tool for high-fidelity non-gravitational force modeling, specifically designed
 57 for orbit determination applications.

58 Software design

59 pyRTX is designed as a modular and extensible library. The software operates on detailed 3D
 60 spacecraft models and computes non-gravitational accelerations by simulating the interaction
 61 between incoming fluxes and the spacecraft geometry. Built around the NAIF SPICE toolkit,
 62 it ensures compatibility with standard astrodynamics pipelines. At its core, pyRTX uses a ray-
 63 tracing engine [Reference a pyEmbree?] in which rays are cast from a discretized representation
 64 of the incoming flux toward the spacecraft. This approach has the advantage to account for
 65 multiple physical effects:

- 66 ■ *3D spacecraft modeling*: The software operates directly on high-resolution 3D mesh
 67 models (.obj files), allowing accurate representation of spacecraft geometry. This naturally
 68 accounts for self-shadowing and multiple reflections. Optical properties can be assigned
 69 for individual mesh elements.
- 70 ■ *Radiation pressure computation*: Solar radiation pressure and planetary albedo and
 71 thermal infrared emission are computed by casting rays from a pixel plane representing
 72 the incoming flux towards the spacecraft [Ziebart et al. 2004]. Momentum transfer is
 73 evaluated through ray-surface interactions, capturing both specular and diffuse reflection
 74 components.
- 75 ■ *Eclipse and shadow modeling*: pyRTX computes shadow functions with high fidelity,
 76 including partial illumination conditions and solar limb darkening, improving the modeling
 77 of eclipse transitions.
- 78 ■ *Atmospheric drag computation*: The effective aerodynamic cross-section is derived via
 79 ray tracing, with the possibility to define an atmospheric density model. This allows
 80 flexible and accurate drag modeling without relying on simplified geometric assumptions.
- 81 ■ *Planetary surface modeling*: The software supports complex planetary shape models
 82 (e.g. digital elevation models) and spatially variable surface properties such as albedo,
 83 emissivity, and temperature. This capability refines not only the computation of planetary
 84 radiation fluxes but also the eclipse transitions by accounting for local topography.
- 85 ■ *Lookup table generation*: The software can precompute accelerations over grids of
 86 incident directions and spacecraft orientations. This enables fast interpolation during

the computation of non-gravitational accelerations, strongly reducing the computational times.

Research impact statement

pyRTX enables a new level of accuracy in non-gravitational force modeling, addressing a major challenge in the orbit determination of deep space missions. By leveraging ray-tracing techniques and realistic spacecraft geometries, the software reduces modeling errors that can otherwise bias scientific results. By providing an open-source and extensible framework, pyRTX supports the scientific community across different missions and institutions.

The package has already been applied to the orbit determination of NASA's Lunar Reconnaissance Orbiter (Zurria et al., 2026), demonstrating its relevance and impact on the data analysis of a present mission. The software is expected to support future missions where non-gravitational effects are dominant and critical to the science objectives, including those to Mercury, Venus, Earth, the Moon, and Mars.

Acknowledgements

Work by G.C. was supported by NASA under award No. 80GSFC24M0006.

AI Usage Disclosure

Generative AI tools were used to support the preparation of the manuscript and the documentation of the the software. The development of the code, including its algorithms, implementation, and testing was carried out without AI assistance. All content produced with the support of AI has been carefully reviewed and verified by the authors to ensure its validity.

References

- Acton, C. H. (1996). Ancillary data services of NASA's navigation and ancillary information facility. *Planetary and Space Science*, 44(1), 65–70. [https://doi.org/10.1016/0032-0633\(95\)00107-7](https://doi.org/10.1016/0032-0633(95)00107-7)
- Annex, A., Pearson, B., Seignovert, B., Carcich, B., Eichhorn, H., Mapel, J., Von Forstner, J., McAuliffe, J., Del Rio, J., Berry, K., Aye, K.-M., Stefko, M., De Val-Borro, M., Kulumani, S., & Murakami, S. (2020). SpiceyPy: A pythonic wrapper for the SPICE toolkit. *Journal of Open Source Software*, 5(46), 2050. <https://doi.org/10.21105/joss.02050>
- Asmar, S. W., Armstrong, J. W., Iess, L., & Tortora, P. (2005). Spacecraft Doppler tracking: Noise budget and accuracy achievable in precision radio science observations. *Radio Science*, 40(2), n/a–n/a. <https://doi.org/10.1029/2004RS003101>
- Cappuccio, P., Notaro, V., Ruscio, A. di, Iess, L., Genova, A., Durante, D., Stefano, I. di, Asmar, S. W., Ciarcia, S., & Simone, L. (2020). Report on First Inflight Data of BepiColombo's Mercury Orbiter Radio Science Experiment. *IEEE Transactions on Aerospace and Electronic Systems*, 56(6). <https://doi.org/10.1109/TAES.2020.3008577>
- Cappuccio, P., Sesta, A., Di Benedetto, M., Durante, D., De Filippis, U., Di Stefano, I., Iess, L., Mackenzie, R., & Godard, B. (2025). Analysis of radio science data from the KaT instrument of the 3GM experiment during JUICE's early cruise phase. *Aerospace*, 12(1), 56. <https://doi.org/10.3390/aerospace12010056>
- Mazarico, E., Buccino, D., Castillo-Rogez, J., Dombard, A. J., Genova, A., Hussmann, H., Kiefer, W. S., Lunine, J. I., McKinnon, W. B., Nimmo, F., Park, R. S., Roberts, J.

- 128 H., Srinivasan, D. K., Steinbrügge, G., Tortora, P., & Withers, P. (2023). The Europa
129 Clipper Gravity and Radio Science Investigation. *Space Science Reviews*, 219(4), 30.
130 <https://doi.org/10.1007/s11214-023-00972-0>
- 131 Milani, A., & Gronchi, G. (2009). *Theory of Orbit Determination*. Cambridge University Press.
132 <https://doi.org/10.1017/CBO9781139175371>
- 133 Tapley, B. D., Schutz, B. E., & Born, G. H. (2004). *Statistical Orbit Determination*. Elsevier.
134 <https://doi.org/10.1016/B978-0-12-683630-1.X5019-X>
- 135 Zurria, A., Cascioli, G., & Mazarico, E. (2026). Refining non-gravitational force modelling
136 of the Lunar Reconnaissance Orbiter with ray tracing. *The Planetary Science Journal*.
137 <https://doi.org/10.0000/pending>

DRAFT