Indian Institute of Technology Roorkee MAN-001(Mathematics-1): B. Tech. I Year

Autumn Semester: 2018-19

Assignment Sheet-3: Differential Calculus (Limit, Continuity and Differentiability)

(a)
$$\lim_{(x,y)\to(-1,2)} \frac{xy^3}{x+y}$$
 (b) $\lim_{(x,y)\to(0,0)} \frac{x^3y}{x^6+y^2}$ (c) $\lim_{(x,y)\to(0,1)} \tan^{-1}\left(\frac{y}{x}\right)$

1. Find the following limits, if they exist

(a)
$$\lim_{(x,y)\to(-1,2)} \frac{xy^3}{x+y}$$
 (b) $\lim_{(x,y)\to(0,0)} \frac{x^3y}{x^6+y^2}$ (c) $\lim_{(x,y)\to(0,1)} \tan^{-1}\left(\frac{y}{x}\right)$ (d) $\lim_{(x,y)\to(0,0)} \frac{xy}{\sqrt{x^2+y^2}}$ (e) $\lim_{(x,y)\to(0,0)} \frac{\sin^2(x+y)}{|x|+|y|}$ (f) $\lim_{(x,y)\to(0,0)} \frac{2x^2y}{x^4+y^2}$ (g) $\lim_{(x,y)\to(1,1)} f(x,y)$ where $f(x,y) = \begin{cases} 1, & \text{if } x+y \geq 2\\ -1, & \text{if } x+y < 2 \end{cases}$ (h) $\lim_{(x,y,z)\to(0,0,0)} \frac{xy+yz+zx}{\sqrt{x^2+y^2+z^2}}$ (i) $\lim_{(x,y)\to(0,0)} \frac{2x^2y}{x^2+y^2}$

(g)
$$\lim_{(x,y)\to(1,1)} f(x,y)$$
 where $f(x,y) = \begin{cases} 1, & \text{if } x+y \ge 2\\ -1, & \text{if } x+y < 2 \end{cases}$

(h)
$$\lim_{(x,y,z)\to(0,0,0)} \frac{xy+yz+zx}{\sqrt{x^2+y^2+z^2}}$$
 (i) $\lim_{(x,y)\to(0,0)} \frac{2x^2y}{x^2+y^2}$

- 2. (a) Consider the function $f(x,y) = \frac{x+y}{x-y}$ for $(x,y) \in \mathbb{R}^2$ with $x-y \neq 0$. Show that $\lim_{x\to 0} [\lim_{y\to 0} f(x,y)] = 1$, but $\lim_{y\to 0} [\lim_{x\to 0} f(x,y)] = -1$. What can you say about the existence of $\lim_{(x,y)\to(0,0)} f(x,y)$?
 - (b) Let f(x,y) = 0 if y = 0, and $f(x,y) = x \sin\left(\frac{1}{y}\right)$, if $y \neq 0$. Compute $\lim_{\substack{(x,y)\to(0,0)\\ (x,y)=(0,0)}} f(x,y)$ and iterated limits $\lim_{\substack{x\to 0\\ y\to 0}} [\lim_{y\to 0} f(x,y)]$ and $\lim_{y\to 0} [\lim_{x\to 0} f(x,y)]$ if they exist.
 - (c) Let $f(x,y) = \frac{x^2y^2}{x^2y^2 + (x-y)^2}$ if $x^2y^2 + (x-y)^2 \neq 0$. Show that $\lim_{x \to 0} [\lim_{y \to 0} f(x,y)]$ and $\lim_{y \to 0} [\lim_{x \to 0} f(x,y)] = 0$. But $\lim_{(x,y) \to (0,0)} f(x,y)$ does not exist.

3. Let

$$f(x,y) = \begin{cases} 0 & \text{if } x \text{ is rational} \\ 1 & \text{if } x \text{ is irrational} \end{cases}.$$

Show that for any point (a,b), $\lim_{(x,y)\to(a,b)} f(x,y)$ does not exist.

4. Examine the continuity of the function f(x, y) at (0, 0) in each of the following cases:

(a)
$$f(x,y) = \begin{cases} \frac{\sin^{-1}(x+2y)}{\tan^{-1}(2x+4y)}, & (x,y) \neq (0,0) \\ \frac{1}{2}, & (x,y) = (0,0) \end{cases}$$

(b) $f(x,y) = \begin{cases} x \sin\frac{1}{x} + y \sin\frac{1}{y}, & xy \neq 0 \\ 0, & xy = 0 \end{cases}$
(c) $f(x,y) = \begin{cases} xy \log(x^2 + y^2), & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$
(d) $f(x,y) = \begin{cases} \frac{x^2y^2}{x^3 + y^3}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$
(e) $f(x,y) = \sqrt{|xy|}$.

- 5. For the function $f(x,y) = \begin{cases} \frac{y(x^2 y^2)}{x^2 + y^2}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$. Compute $f_x(0,y), f_y(x,0), f_x(0,0)$ and $f_y(0,0)$, if they exist.
- 6. Show that for the function $f(x,y) = \begin{cases} -xy, & |y| \ge |x| \\ xy, & |y| > |x| \end{cases}$, $f_{xy}(0,0)$ and $f_{yx}(0,0)$ both exist and are unequal.
- 7. Prove that |x| + |y| is continuous, but not differentiable at (0,0).
- 8. Prove that $f(x,y) = \begin{cases} \frac{(x+y)\{\sqrt{(x^2+y^2)}+xy\}}{\sqrt{(x^2+y^2)}}, & \text{when } x^2+y^2 \neq 0 \\ 0, & \text{when } x=y=0, \end{cases}$ is differentiable at (0,0). Hence, deduce that $f_x(0,0) = f_y(0,0) = 1$.
- 9. Show that the function $f(x,y) = \begin{cases} x^3 \sin\frac{1}{x^2} + y^3 \sin\frac{1}{y^2}, & \text{when } xy \neq 0 \\ x^3 \sin\frac{1}{x^2}, & \text{when } x \neq 0 \text{ and } y = 0 \\ y^3 \sin\frac{1}{y^2}, & \text{when } x = 0 \text{ and } y \neq 0 \\ 0, & \text{when } x = y = 0 \end{cases}$ is differentiable at (0,0), whereas none of f_x, f_y is continuous at (0,0).
- 10. Show that the function $f(x,y) = \begin{cases} (x^2 + y^2) \cos\left(\frac{1}{\sqrt{x^2 + y^2}}\right), & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$ is differentiable at (0,0) and that f_x, f_y are not continuous at (0,0).

Answers:

- 1. (a) -8 (b) does not exist (c) does not exist (d) 0 (e) 0 (f) does not exist (g) does not exist (h) 0 (i) 0
- 2. (a) does not exist (b) 0, does not exist, 0
- 4. (a) continuous (b) continuous (c) continuous (d) discontinuous (e) continuous
- 5. 0, 1, 0, -1.