

# Systematic survey occurrence data template instructions

## Overview

Use this template to record occurrence data; that is the presence or absence of an organism at a particular site locality at a point in time.

Templates have been provided to facilitate integration of your data into the Biodiversity Data Repository database. Not all types of data have been catered for in the available templates at this stage; therefore, if you are unable to find a suitable template, please contact [bdr-support@gaiaresources.com.au](mailto:bdr-support@gaiaresources.com.au) to make us aware of your data needs.

## NEED TO KNOW:

For data validation, you will need your data file to:

- be the correct **file format**,
- have **matching template fields** to the template downloaded (do not add, remove, or change the order of fields),
- have populated the relevant fields using the correct data type (for example dates for date fields),
- have values in **mandatory fields**,
- comply with data **value constraints** for example the geographic coordinates are consistent with a [geodeticDatum](#) type of the four available options and latitude and longitude values fall within Australia (see [Further Information](#) for details), and
- align with existing controlled [vocabularies](#) wherever possible (this is mandatory for geodeticDatum), but new terms may be submitted for consideration and will not cause a validation error.

## File format

The systematic survey occurrence data template is a **UTF-8** encoded csv (not Microsoft Excel Spreadsheets). Be sure to save this file with your data as a .csv (UTF-8) otherwise it will not pass the csv validation step upon upload. **Do not include additional columns or empty rows.**

## Template fields

The template contains the field names in the top row. Table 1 will assist you in transferring your data to the template with the information:

- **Field name** in the template (and an external link to the [Darwin Core standard](#) for that field where relevant);
- **Description** of the field;
- **Required** whether the field is **mandatory** or **optional**;

- **Format** (datatype) required for the data values for example text (string), number (integer, float), or date;
- **Example** of an entry for that field; and
- **Vocabulary** [links](#) within this document (for example pick list values) where relevant. The fields that have suggested values are highlighted in Table 1 and the options for those fields are listed in Table 2 in alphabetical order of field name.

Table 1: Systematic survey occurrence data template fields with descriptions, conditions, datatype format, and examples.

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
recordID	The unique identifier allocated to this record by the data provider e.g. when data is provided by an aggregator. It connects the record to the collection and storage of the specimens when these parts of the workflow may have been done by different parties.	Optional	String	89T22FSJMJ079c5cf
<a href="#">locality</a>	The specific description of the place.	Optional	String	Cowaramup Bay Road, Margaret River
<b><a href="#">decimalLatitude</a></b>	The geographic latitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location. Positive values are north of the Equator, negative values are south of it. Valid coordinate ranges for the BDR system apply - see <a href="#">Further Information</a> section.	Mandatory	Float	-33.812314
<b><a href="#">decimalLongitude</a></b>	The geographic longitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location. Positive values are east of the Greenwich Meridian, negative values are west of it. Valid coordinate ranges for the BDR system apply - see <a href="#">Further Information</a> section.	Mandatory	Float	115.231512

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
<b><u>geodeticDatum</u></b>	The acronym for the ellipsoid, geodetic datum, or spatial reference system (SRS) upon which the geographic (non-projected) coordinates given in decimalLatitude and decimalLongitude as based.	Mandatory	String	WGS84 <a href="#">(Vocabulary link)</a>
<a href="#">coordinateUncertaintyInMeters</a>	The horizontal distance (in metres) from the given decimalLatitude and decimalLongitude describing the smallest circle containing the whole of the Location. Leave the value empty if the uncertainty is unknown, cannot be estimated, or is not applicable (because there are no coordinates). Zero is not a valid value for this term.	Optional	Float	50
<a href="#">dataGeneralizations</a>	Actions taken to make the shared data less specific or complete than in its original form, due to restrictions around identifying locations of particular species. Suggests that alternative data of higher quality may be available on request.	Optional	String	Coordinates rounded to the nearest 10 km for conservation concern
<b><u>eventDate</u></b>	The date (DD/MM/YYYY or (YYYY-MM-DD are accepted) or date-time with zone (in <a href="#">ISO 8601</a> format for example2022-05-20T0623+08) during which a species occurrence was observed. For occurrences, this is the date-time when the event was recorded. Not suitable for a time in a geological context.	Mandatory	Date	2019-09-24

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
<a href="#">samplingProtocol</a>	<p>The sampling protocol is the method used to sample the locality to determine the presence (or absence) of the taxon referred to in this record at the indicated time. This may be a collecting method or a method to observe an organism without collection.</p> <p>Recommended best practice is to describe a species occurrence with no more than one sampling protocol. In the case of a summary, in which a specific protocol can not be attributed to specific species occurrences, the recommended best practice is to repeat the property for each IRI that denotes a different sampling protocol that applies to the occurrence.</p>	Optional	String	Human observation
<a href="#">basisOfRecord</a>	The specific nature of the data record.	Optional	String	Preserved Specimen <a href="#">(Vocabulary link)</a>
<a href="#">recordedBy</a>	A person, group, or organisation responsible for recording the original Occurrence.	Optional	String	J. Doe
<a href="#">recordNumber</a>	An identifier given to the Occurrence at the time it was recorded. Often serves as a link between field notes and an Occurrence record, such as a specimen collector's number.	Optional	String	OPP 7101
<a href="#">occurrenceStatus</a>	A statement about the presence or absence of a Taxon at a Location.	Optional	String	Present <a href="#">(Vocabulary link)</a>

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
<a href="#">habitat</a>	A category or description of the habitat in which the event occurred.	Optional	String	Closed forest of <i>Melaleuca lanceolata</i> . White, grey or brown sand, sandy loam.
<a href="#">establishmentMeans</a>	Statement about whether an organism or organisms have been introduced to a given place and time through the direct or indirect activity of modern humans.	Optional	String	Native ( <a href="#">Vocabulary link</a> )
<a href="#">organismRemarks</a>	Comments or notes about the Organism instance.	Optional	String	good condition
<a href="#">individualCount</a>	The number of individuals present at the time of the Occurrence. 0 = none, no value = the specific number was not recorded.	Optional	Integer	1
<a href="#">lifeStage</a>	The age class or life stage of the Organism(s) at the time the Occurrence was recorded.	Optional	String	Mature ( <a href="#">Vocabulary link</a> )
<a href="#">sex</a>	The sex of the biological individual(s) represented in the Occurrence.	Optional	String	Unspecified ( <a href="#">Vocabulary link</a> )
<a href="#">reproductiveCondition</a>	The reproductive condition of the biological individual(s) represented in the Occurrence.	Optional	String	No breeding evident
<a href="#">occurrenceID</a>	An identifier for the Occurrence (as opposed to a particular digital record of the occurrence). There are no constraints on how the identifier is named, except that it should be unique.	Optional	String	S09-20190924-02

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
<a href="#">ownerInstitutionCode</a>	The name (or acronym) in use by the institution having ownership of the object(s) or information referred to in the record. This institution is the custodian of the occurrenceID and/or catalogNumber that are appended to this record.	Optional	String	WAM
<a href="#">collectionCode</a>	The name, acronym, code, or initialism identifying the collection or data set from which the record was derived. It is associated with the catalogNumber.	Optional	String	HERB
<a href="#">catalogNumber</a>	An identifier (preferably unique) for the record within the data set or collection.	Optional	String	145732, 145732a, 2008.1334, R-4313
<a href="#">institutionCode</a>	The name (or acronym) in use by the institution having custody of the object(s) or information referred to in the record. This institution is a stakeholder in the creation of this record.	Optional	String	MUS
<a href="#">otherCatalogNumbers</a>	A list (concatenated and separated with a space vertical bar space (   )) of previous or alternate fully qualified catalog numbers or other human-used identifiers for the same Occurrence, whether in the current or any other data set or collection.	Optional	String	NPS YELLO6778   MBG 33424
<a href="#">preparations</a>	A list (concatenated and separated with a space vertical bar space (   )) of preparations and preservation methods for a specimen.	Optional	String	alcohol ( <a href="#">Vocabulary link</a> )

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
preparedDate	The date (DD/MM/YYYY or (YYYY-MM-DD are accepted) or date-time with zone (in <a href="#">ISO 8601</a> format for example 2022-05-20T0623+08) representing the date or date-time the specimen was prepared.	Optional	Date	2019-09-24
<a href="#">associatedSequences</a>	A list (concatenated and separated with a space vertical bar space (   )) of identifiers (publication, global unique identifier, URI) of genetic sequence information associated with the Occurrence.	Optional	String	<a href="https://www.ncbi.nlm.nih.gov/nuccore/MT766606.1">https://www.ncbi.nlm.nih.gov/nuccore/MT766606.1</a>
sequencingMethod	The method used to obtain sequence data for example DNA, RNA, or protein from the sample.	Optional	String	Sanger-dideoxy-sequencing
<a href="#">verbatimIdentification</a>	A string representing the taxonomic identification as it appeared in the original record. This term is meant to allow the capture of an unaltered original identification/determination, including identification qualifiers, hybrid formulas, uncertainties, etc. This term is meant to be used in addition to scientificName (and identificationQualifier etc.), not instead of it.	Optional	String	<i>Caladenia ?excelsa</i>
<a href="#">dateIdentified</a>	The date (DD/MM/YYYY or (YYYY-MM-DD are accepted) or date-time with zone (in <a href="#">ISO 8601</a> format for example 2022-05-20T0623+08) on which the subject was determined as representing the Taxon.	Optional	Date	2019-09-24



Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
<a href="#">identifiedBy</a>	A list (concatenated and separated with a space vertical bar space (   )) of names of people, groups, or organisations who assigned the Taxon to the subject.	Optional	String	J. Doe   S. Smith
identificationMethod	Method used to associate the organism with the scientificName label.	Optional	String	DNA
<b><a href="#">scientificName</a></b>	The full scientific name, with authorship and date information if known. When forming part of an Identification, this should be the name in lowest level taxonomic rank that can be determined. This term should not contain identification qualifications, which should instead be supplied in the identificationQualifier column. NOTE: Phrase names such as <b>Rhagodia sp. Hamersley (M.Trudgen 17794)</b> are permitted in the scientificName field where those are in use.	Mandatory	String	<i>Caladenia excelsa</i>
<a href="#">identificationQualifier</a>	A brief phrase or a standard term ("cf.", "aff.") to express the determiner's doubts about the Identification.	Optional	String	Species incerta <a href="#">(Vocabulary link)</a>
<a href="#">identificationRemarks</a>	Comments or notes about the Identification.	Optional	String	DNA evidence may indicate a new species. Further analysis required.
<a href="#">acceptedNameUsage</a>	The full name, with authorship and date information if known, of the currently valid (zoological) or accepted (botanical) taxon.	Optional	String	<i>Calaendia excelsa</i> Hopper & A.P.Br.

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
<b>kingdom</b>	The full scientific name of the kingdom in which the taxon is classified.	Mandatory	String	Plantae <a href="#">(Vocabulary link)</a>
<a href="#">taxonRank</a>	The taxonomic rank of the most specific name in the scientificName.	Optional	String	Species <a href="#">(Vocabulary link)</a>
threatStatus	The conservation status (or code) assigned to an organism that is recognised in conjunction with a specific jurisdiction.	Conditionally mandatory with conservationJurisdiction	String	EN <a href="#">(Vocabulary link)</a>
conservationJurisdiction	The jurisdiction under which an organism is recognised to have a specific conservation status applied.	Conditionally mandatory with threatStatus	String	EPBC, WA <a href="#">(Vocabulary link)</a>
threatStatusCheck Protocol	The method used to determine if the organism is listed under the relevant jurisdictional threatened species list.	Optional	String	Species name check of the Department of Climate Change, Energy, the Environment and Water's Species Profile and Threat Database <a href="http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl">http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl</a>
threatStatusDateDetermined	The date on which this record of this organism was assigned to the nominated threatStatus and conservationJurisdiction	Optional	Date	dd/mm/yyyy

Field name	Description	Mandatory / Optional	Datatype Format	Examples / Vocabulary
threatStatusDeterminedBy	The person and/organisation responsible for appending the threatStatus and conservationJurisdiction to this organism's occurrence record.	Optional	String	J. Bloggs, Department of Environment

## Vocabulary lists

Apart from geodeticDatum, the data validation does not require adherence to the below vocabularies for each of the fields indicated as having vocabularies. These vocabularies are provided as a means of assistance in developing consistent language within the database. New terms can be added to more appropriately describe your data that goes beyond the current list. Table 2 provides some suggested values from existing sources such as: [Biodiversity Information Standard \(TDWG\)](#), [EPSG.io Coordinate systems worldwide](#), the [Global Biodiversity Information Facility](#), and [Open Nomenclature in the biodiversity era](#).

Table 2: Suggested values for controlled vocabulary fields in the template. Each term has a preferred label with a definition to aid understanding of its meaning. For some terms, alternative labels are provided that mean the same sort of thing. Note: **geodeticDatum value must come from one of the four options in this table.**

Template field name	Preferred label	Definition	Alternate label
basisOfRecord	FossilSpecimen	A preserved specimen that is a fossil.	Fossil Specimen
	HumanObservation	An output of a human observation process.	Human Observation
	LivingSpecimen	A specimen that is alive.	Living Specimen
	MachineObservation	An output of a machine observation process.	Machine Observation
	MaterialSample	A physical result of a sampling (or subsampling) event. In biological collections, the material sample is typically collected, and either preserved or destructively processed.	Material Sample
	Occurrence	An existence of an Organism (sensu <a href="http://rs.tdwg.org/dwc/terms/Organism">http://rs.tdwg.org/dwc/terms/Organism</a> ) at a particular place at a particular time.	
	PreservedSpecimen	A specimen that has been preserved.	Preserved Specimen

establishmentMeans	introduced	Establishment of a taxon by human agency into an area that is not part of its natural range.	alien, exotic, non-native, nonindigenous
	introducedAssistedColonisation	Establishment of a taxon specifically with the intention of creating a self-sustaining wild population in an area that is not part of the taxon's natural range.	assisted colonisation
	native	A taxon occurring within its natural range.	native (indigenous)
	nativeReintroduced	A taxon re-established by direct introduction by humans into an area that was once part of its natural range, but from where it had become extinct.	native: reintroduced
	uncertain	The origin of the occurrence of the taxon in an area is obscure.	unknown, cryptogenic
	vagrant	The temporary occurrence of a taxon far outside its natural or migratory range.	casual
geodeticDatum	AGD84	Australian Geodetic Datum 1984	EPSG:4203
	GDA2020	Geocentric Datum of Australia 2020	EPSG:7844
	GDA94	Geocentric Datum of Australia 1994	EPSG:4283
	WGS84	World Geodetic System 1984, used in GPS	EPSG:4326
identificationQualifier	animalia cetera	It groups all the unidentified specimens that are not listed as separate taxa. The term cetera (abbreviated c. or cet.) may be applied to a given high-rank taxon, meaning that identification at a lower taxonomic level has not been attempted (see also stetit) but explicitly not including subordinate taxa that may have been identified.	a.c., A.C.
	confer	"Compare with". Specimens should be compared to reference material, since most of the diagnostic characters correspond to a given species but some are unclear. Also used in the sense of affinis and species incerta (these usages are discouraged).	cf., cfr., conf., sp. cf.

	ex grege	"Of the group including". The specimen has some affinity to a known species or it belongs to a species group or species complex; see also <i>affinis</i> and <i>species proxima</i> .	ex gr., gr.
	familia genus species	The specimen has not been attributed to any known species nor family; see also <i>species</i> .	fam. gen. sp.
	genus et species nova	The specimen is considered to belong to a new species and a new genus; for more details, see <i>species nova</i> .	gen. et sp. nov., gen. nov., sp.nov., nov. gen. et sp.
	genus novum	The specimen is considered to belong to a new species and a new genus; for more details, see <i>species nova</i>	gen. nov., g. nov., gen. n., g. n., nov. gen.
	genus species	The specimen has not been related to any known species nor genus; see also <i>species</i> .	gen. sp., g. sp.
	species	The specimen has not been identified, nor it has been related to any known species; the uncertainty is potentially provisional: it could be due to the lack of suitable dichotomous keys, or to the occurrence of a species not previously described. Also used in the sense of <i>species indeterminabilis</i> and <i>stetit</i> (these usages are discouraged).	sp.
	species (pl.),	More than one species belonging to the same genus (or higher-rank taxon) are included.	spp., sp. pl.
	species affinis	"Has affinity with". The specimen has some affinity to a known species but it is not identical to it; it generally implies distinction more than a possible identity, in contrast with the qualifier <i>confer</i> ; see also <i>species Proxima</i> and <i>ex grege</i> . It is often used in combination with the ON qualifier <i>species nova</i> . Also used in the sense of <i>confer</i> (this usage is discouraged).	aff., sp. aff.

	species incerta	The identification is uncertain; it usually indicates a higher reliability with respect to confer. The sign "sp. inc." is also used in the sense of species, species indeterminabilis and species inquirenda (these usages are discouraged).	?, sp. Inc
	species indeterminabilis	The specimen is indeterminable beyond a certain taxonomic level due to the deterioration or lack of diagnostic characters. Also used in the sense of species and stetit (these usages are discouraged).	indet., ind., sp. indet., sp. ind.
	species nova	The specimen is considered to belong to a new, previously undescribed species. (1) When describing a new species, the use of the qualifier is required by the ICZN (1999) to explicitly indicate the taxa name as intentionally new. (2) Used as ON qualifier to refer to a new, still unnamed species before the formal publication of the description.	sp. nov., spec. nov., sp. n., nov. sp., nov. spec., n. sp.
	species proxima	The specimen is near to a known species but it is not identical to it; see also affinis and ex grege.	prox., sp. prox., nr., sp. nr.
	stetit	Identification at a lower taxonomic level has not been attempted, even if allowed by the sample conditions. It may also be used when more records with different ON qualifiers need to be merged at a safe taxonomic level.	stet., the intentional absence of qualifiers
	subspecies	The only infraspecific rank regulated by the ICZN (1999). As ON qualifier, it indicates that the specimen probably belongs to a subspecies but it has not been related to any known one; see also species.	ssp., subsp.
kingdom	Animalia	Kingdom Animalia	
	Fungi	Kingdom (taxonRank: Regnum) Fungi	
	Plantae	Kingdom (taxonRank: Regnum) Plantae	Plantae Haeckel
lifeStage	adult	An adult is a plant, animal, or person who has reached full growth or alternatively is capable of reproduction.	imago

	embryo	An embryo is a multicellular diploid eukaryote in its earliest stage of development, from the time of first cell division until birth, hatching, or germination.	egg, seed
	gamete	A gamete is a cell that fuses with another gamete during fertilization in organisms that reproduce sexually. In species that produce two morphologically distinct types of gametes, and in which each individual produces only one type, a female is any individual that produces the larger type of gamete — called an ovum (or egg) — and a male produces the smaller tadpole-like type — called a sperm. This is an example of anisogamy or heterogamy, the condition wherein females and males produce gametes of different sizes. In contrast, isogamy is the state of gametes from both sexes being the same size and shape, and given arbitrary designators for mating type. Gametes carry half the genetic information of an individual, one chromosome of each type.	ovum, sperm, pollen
	gametophyte	In plants and algae that undergo alternation of generations, a gametophyte is the multicellular structure, or phase, that is haploid, containing a single set of chromosomes. The gametophyte produces male or female gametes (or both), by a process of cell division called mitosis. In mosses, liverworts and hornworts (bryophytes), the gametophyte is the commonly known phase of the plant. An early developmental stage in the gametophyte of mosses (immediately following germination of the meiospore) is called the protonema. In most other land plants the gametophyte is very small (as in ferns and their relatives) or even reduced as in flowering plants (angiosperms), where the female gametophyte (ovule) is known as a megagametophyte and the male gametophyte (pollen) is called a microgametophyte.	gamont, protonema, pollen, ovule



	juvenile	A juvenile is an individual organism that has not yet reached its adult form, sexual maturity or size. Juveniles sometimes look very different from the adult form, particularly in terms of their colour. In many organisms the juvenile has a different name from the adult.	seedling, eft, calf, hatchling, infant, foal, kitten, kit, chick, nymph, fawn, whelp, pup, elver, fry
	larva	A larva (Latin; plural larvae) is a young (juvenile) form of animal with indirect development, going through or undergoing metamorphosis (for example, insects, amphibians, or cnidarians). The larva can look completely different from the adult form, for example, a caterpillar differs from a butterfly. Larvae often have special (larval) organs which do not occur in the adult form. The larvae of some species can become pubescent and not further develop into the adult form (for example, in some newts). This is a type of neoteny. It is a misunderstanding that the larval form always reflects the group's evolutionary history. It could be the case, but often the larval stage has evolved secondarily, as in insects. In these cases the larval form might differ more from the group's common origin than the adult form. The early life stages of most fish species are considerably different from juveniles and adults of their species and are called larvae.	larvae, tadpole, polliwog, pollywog, polliwig, polewig, polwig, planula, nauplius, zoea, nymph, caterpillar, grub, maggot, wriggler, trochophore, veliger, glochidium, ammocoete, leptocephalus, bipinnaria, caterpillar, grub, maggot
	pupa	A pupa is the life stage of some insects undergoing transformation between immature and mature stages. The pupal stage is found only in holometabolous insects, those that undergo a complete metamorphosis, with four life stages: egg (-> embryo), larva, pupa, and imago (-> adult).	puppe

	spore	A spore is a reproductive structure that is adapted for dispersal and surviving for extended periods of time in unfavorable conditions. Spores form part of the life cycles of many bacteria, plants, algae, fungi and some protozoans. A chief difference between spores and seeds as dispersal units is that spores have very little stored food resources compared with seeds. Spores are usually haploid and unicellular and are produced by meiosis in the sporangium by the sporophyte. Once conditions are favorable, the spore can develop into a new organism using mitotic division, producing a multicellular gametophyte, which eventually goes on to produce gametes. Many ferns, especially those adapted to dry conditions, produce diploid spores. In this case spores are the units of asexual reproduction, because a single spore develops into a new organism. By contrast, gametes are the units of sexual reproduction, as two gametes need to fuse to create a new organism.	-
	sporophyte	All land plants, and some algae, have life cycles in which a haploid gametophyte generation alternates with a diploid sporophyte, the generation of a plant or alga that has a double set of chromosomes. A multicellular sporophyte generation or phase is present in the life cycle of all land plants and in some green algae. For common flowering plants (Angiosperms), the sporophyte generation comprises almost their whole life cycle (that is whole green plant, roots etc), except phases of small reproductive structures (pollen and ovule).	agamont

	zygote	A zygote (or zygocyte) describes the first stage of a new unique organism when it consists of just a single cell. The term is also used more loosely to refer to the group of cells formed by the first few cell divisions, although this is properly referred to as a blastomere. A zygote is usually produced by a fertilisation event between two haploid cells - an ovum from a female and a sperm cell from a male - which combine to form the single diploid cell. Thus the zygote contains DNA originating from both mother and father and this provides all the genetic information necessary to form a new individual	blastomere
occurrenceStatus	absent	The occurrence was not present at the location and time of the observation.	-
	present	The occurrence was present at the location and time of the observation.	-
preparations	alcohol	Alcohol	-
	deepFrozen	Deep frozen	Deep frozen
	dried	Dried	-
	driedAndPressed	Dried and pressed	Dried and pressed
	formalin	Formalin	-
	freezeDried	Freeze-dried	Freeze-dried
	glycerin	Glycerin	-
	gumArabic	Gum arabic	Gum arabic
	microscopicPreparation	Microscopic preparation	Microscopic preparation
	mounted	Mounted	-
	noTreatment	No treatment	No treatment
	other	Other	unspecified

	pinned	Pinned	-
	refrigerated	Refrigerated	-
sex	female	Female (♀) is the sex of an organism, or a part of an organism, which produces mobile ova (egg cells).	F, ♀
	hermaphrodite	One organism having both male and female sexual characteristics and organs; at birth an unambiguous assignment of male or female cannot be made	Zwitter
	male	Male (♂) refers to the sex of an organism, or part of an organism, which produces small mobile gametes, called spermatozoa.	M, ♂
	undetermined	If the sex of an organism can't be determined for some reason.	Undet., unknown
taxonRank	class	class	-
	cultivar	The epithet is usually output in single quotes and may contain multiple words, see ICBN §28. Examples: <i>Taxus baccata</i> 'Variegata', <i>Juniperus ×pfitzeriana</i> 'Wilhelm Pfitzer'; <i>Magnolia</i> 'Elizabeth' (= a hybrid, no species epithet).	-
	cultivarGroup	cultivar group	grex
	family	family	-
	form	form	forma
	genus	genus	-
	informal	informal	-
	infragenericname	Used for any other unspecific rank below genera and above species.	-
	infraorder	infraorder	-
	infraspecificname	Used for any other unspecific rank below genera and above species.	-
	infrasubspecificname	Used for any other unspecific rank below subspecies.	-

	kingdom	kingdom	regnum
	order	order	alliance
	phylum	phylum	division
	section	Section within a genus. In Zoology a section sometimes refers to a group above family level, this is NOT meant	
	series	Series within a genus	
	species	species	
	speciesAggregate	A loosely defined group of species. Zoology: 'Aggregate - a group of species, other than a subgenus, within a genus. An aggregate may be denoted by a group name interpolated in parentheses.' -- The Berlin/MoreTax model notes: '[these] aren't taxonomic ranks but circumscriptions because on the one hand they are necessary for the concatenation of the fullname and on the other hand they are necessary for distinguishing the aggregate or species group from the microspecies.' Compare subspecific aggregate for a group of subspecies within a species!	aggregate, species group, species complex
	subfamily	subfamily	
	subform	subform	subforma
	subgenus	subgenus	
	suborder	suborder	
	subsection	Subsection within a genus	
	subseries	Subseries within a genus	
	subspecies	subspecies	

	subspecificAggregate	A loosely defined group of subspecies. Zoology:'Aggregate - a group of subspecies within a species. An aggregate may be denoted by a group name interpolated in parentheses.'	
	subtribe	subtribe	
	subvariety	subvariety	subvarietas
	superfamily	superfamily	
	supragenericname	Used for any other unspecific rank above genera.	
	tribe	tribe	
	unranked	unranked	
	variety	variety	varietas

Table 2b: Suggested values for conditionally mandatory values for the threatStatus and conservationJurisdiction in the template. State and Territory conservationJurisdictions spelt out in words are also valid. For some threatStatus terms, alternative labels are provided that are also valid for that conservationJurisdiction.

<b>conservationJurisdiction</b>	<b>threatStatus</b>	<b>threatStatus alternative labels</b>
ACT	extinct	
	extinct in the Wild	
	critically endangered	
	endangered	
	vulnerable	
	regionally conservation dependent	
EPBC	EX	extinct
	XW	extinct in the wild, EW
	CE	critically endangered, CR
	E	Endangered, EN
	V	Vulnerable, VU
	CD	conservation dependent
	JAMBA	
	CAMBA	
	KAMBA	
	CITES	
NSW	EX	Extinct
	CE	Critically Endangered
	En	Endangered
	V	Vulnerable
NT	Ex	Extinct
	Ew	Critically endangered (possibly extinct)
	CE	Critically endangered
	EN (extinct in NT)	Endangered (extinct in NT)
	EN (extinct in wild in NT)	Endangered (extinct in wild in NT)
	EN	Endangered
	Vu (extinct in NT)	Vulnerable (extinct in NT)
	Vu	Vulnerable
	NT	Near Threatened
	LC (extinct in NT)	Least concern (extinct in NT)
	LC	Least concern

	DD	Data Deficient
	NE	Not Evaluated
QLD	EX	Extinct wildlife
	PE	Extinct in the wild wildlife
	CR	Critically endangered wildlife
	E	Endangered wildlife
	V	Vulnerable wildlife
	NT	Near threatened wildlife
	SL	Special least concern wildlife
	C	Least concern wildlife
	I	International wildlife
SA	E	Endangered
	V	Vulnerable
	R	Rare
	ssp.	indicates that at least one subspecies for this species has been given a conservation rating
	sp.	indicates that a rating has been applied to the species level. The status has not been assessed at the subspecies level
TAS	x	Extinct
	e	Endangered
	v	Vulnerable
	r	Rare
VIC	Extinct	
	Extinct in the Wild	
	Critically Endangered	
	Endangered (Extinct in Victoria)	
	Endangered	
	Vulnerable	
	Conservation Dependent	
	Restricted	
WA	T	
	CR	critically endangered species
	EN	endangered species
	VU	vulnerable species
	EX	extinct species
	EW	extinct in the wild
	Specially protected species	
	MI	migratory species



	CD	species of special conservation interest (conservation dependent fauna)
	OS	other specifically protected fauna
	P1	priority 1: poorly-known species
	P2	priority 2: poorly-known species
	P3	priority 3: poorly-known species
	P4	priority 4: rare, near threatened and other species in need of monitoring

## Further information

The valid latitude and longitude coordinates range in the BDR system is based on information from Geoscience Australia. The **Australian Continent** coordinate extremities can be found on this site:

<https://www.ga.gov.au/scientific-topics/national-location-information/dimensions/continental-extremities>

The coordinates range for the seven **Australian External and Offshore territories** can be found on these sites:

<https://d28rz98at9flks.cloudfront.net/70562/AustraliaAndExternalTerritories.pdf>

<https://www.ga.gov.au/scientific-topics/national-location-information/dimensions/remote-offshore-territories>)

In summary:

- **Australian continent**  
Latitude range: -10.689167 and -43.644444  
Longitude range: 153.637222 and 113.155000
- **Heard and McDonalds Island**  
Latitude range: -52.902770 and -53.195018  
Longitude range: 73.872715 and 72.577376
- **Lord Howe Island**  
Latitude range: -31.486129 and -31.787767  
Longitude range: 159.280368 and 159.036807
- **Macquarie Island**  
Latitude range: -54.355874 and -55.123198  
Longitude range: 158.998625 and 158.674929
- **Norfolk Island**  
Latitude range: -29.136568 and -28.994170  
Longitude range: 167.998035 and 167.913770
- **Ashmore and Cartier Islands**  
Latitude range: -12.184700 and -12.547300  
Longitude range: 123.581854 and 122.927010
- **Christmas Island**  
Latitude range: -10.412390 and -10.570559  
Longitude range: 105.712810 and 105.533149
- **Cocos Islands**  
Latitude range: -11.822133 and -12.211000  
Longitude range: 96.930763 and 96.815497
- **Coral Sea Islands**  
Latitude range: -15.721024 and -29.982747  
Longitude range: 159.140729 and 147.839456
- **Australian Antarctic Territory**  
Latitude range: -64.928558 to -90.000026  
Longitude range: 167.724334 to 45.000000

For assistance, please contact: [bdr-support@gaiaresources.com.au](mailto:bdr-support@gaiaresources.com.au)