

Common Name: LITTLE RIVER BLACK-EYED SUSAN

Scientific Name: Rudbeckia heliopsidis Torrey & Gray

Other Commonly Used Names: sun-facing coneflower

Previously Used Scientific Names: none

Family: Asteraceae/Compositae (aster)

**Rarity Ranks:** G2/S1

**State Legal Status:** Threatened

Federal Legal Status: none

Federal Wetland Status: none

**Description:** Perennial **herb** with hairy, many-branched stems, 2 - 4 feet (60 - 120 cm) tall. **Leaves** 23/8 - 8 inches (6 - 20 cm) long and 3/4 - 2 inches (2 - 5 cm) wide, alternate, oval, with rounded bases and pointed tips, smooth to roughly hairy surfaces, and toothed margins (occasionally without teeth); basal leaves have long leaf stalks, stem leaves have shorter stalks. **Flower heads** consist of a raised, rounded, purplish-brown disk, 3/8 - 5/8 inch (1 - 1.5 cm) wide;

6 - 12 yellow ray flowers,  $\frac{1}{2}$  -  $\frac{1}{4}$  inch (1.5 - 3 cm) long; a whorl of gland-dotted bracts surrounding the base of the head; and thin, very hairy scales (**chaff**), less than  $\frac{1}{16}$  inch (0.15 mm) long, attached near the base of each disk flower. **Fruits**  $\frac{1}{8}$  inch (3 mm) long, dry, seedlike.

**Similar Species:** Black-eyed Susan (*Rudbeckia hirta*) is a common, widespread species with very hairy stems, leaves, and bracts. Its deep yellow ray flowers are tinged with orange on the lower half and are ½ - 1¾ inches (1.5 - 4.5 cm) long. Orange coneflower (*Rudbeckia fulgida*) also resembles Little River black-eyed Susan; it grows in wet pine savannas or in deep sands along streams. Its chaff is smooth or only slightly hairy.

**Related Rare Species:** Large-flowered coneflower (*Rudbeckia grandiflora*, Special Concern) occurs on limestone glades in Catoosa County. It has thick, hairy, lance-shaped basal leaves up to 10 inches (25 cm) long and 4 inches (10 cm) wide. The ray flowers are yellow, drooping, and up to 2 inches (5 cm) long; the disk is dark brown, flat-topped, and raised as much as 1 inch (2.5 cm) above the rays.

**Habitat:** Limestone or sandstone outcrops and nearby streamsides, Coosa Valley prairies, and roadsides and rights-of-way through these habitats; requires openings created by fire, stream scouring, or other clearing.

**Life History:** Little River black-eyed Susan is a perennial herb that reproduces sexually as well as vegetatively by producing new plants at the tips of spreading underground stems. There is no information about pollination of Little River black-eyed Susan; most members of this genus are pollinated by a variety of bees, butterflies, and flies and require cross-pollination to set seed. Seeds are dispersed by gravity, flowing water, and small animals.

**Survey Recommendations:** Surveys are best conducted during flowering (July–October).

Range: Georgia, Alabama, South Carolina, North Carolina, and Virginia.

**Threats:** Conversion of habitat to pine plantations or developments; use of herbicides in rights-of-way; disruption of natural flooding cycles in streams; fire suppression in prairies; overgrowth of aggressive weedy competitors.

Georgia Conservation Status: Six populations are currently known, none on conservation land.

**Conservation and Management Recommendations:** Use prescribed fire or hand-clearing to remove competing vegetation. Avoid clearcutting and other mechanical disturbances to the ground cover. Avoid use of herbicides in highway rights-of-way. Avoid damming streams.

## **Selected References:**

Chafin, L.G. 2007. Field guide to the rare plants of Georgia. State Botanical Garden of Georgia and University of Georgia Press, Athens.

Cronquist, A. 1980. Vascular flora of the southeastern United States, Vol. 1, Asteraceae. University of North Carolina Press, Chapel Hill.

FNA. 2006. Flora of North America, Vol. 21, Magnoliophyta: Asteridae, Part 8: Asteraceae, Part 3. Oxford University Press, New York.

NatureServe. 2008. NatureServe Explorer. Arlington, Virginia. <a href="http://www.natureserve.org/explorer">http://www.natureserve.org/explorer</a>

Radford, A.E., H.E. Ahles, and C.R. Bell. 1968. Manual of the vascular flora of the Carolinas. University of North Carolina Press, Chapel Hill.

Sorrie, B.A. 1995. Status survey for sun-facing sunflower (*Rudbeckia heliopsidis*) status survey. North Carolina Natural Heritage Program, Division of Parks and Recreation. Raleigh.

Weakley, A.S. 2008. Flora of the Carolinas, Virginia, Georgia, northern Florida, and surrounding areas. University of North Carolina Herbarium, Chapel Hill. http://www.herbarium.unc.edu/flora.htm

**Author of Species Account:** Linda G. Chafin

## **Date Compiled or Updated:**

L. Chafin, Aug. 2008: original account K. Owers, Feb. 2010: added pictures



