



Common Name: HEARTLEAF BRICKELLIA

Scientific Name: *Brickellia cordifolia* Elliott

Other Commonly Used Names: Flyr's brickell-bush

Previously Used Scientific Names: *Coleosanthus cordifolius* (Elliott) Kuntze

Family: Asteraceae/Compositae (aster)

Rarity Ranks: G2G3/S2

State Legal Status: Threatened

Federal Legal Status: none

Federal Wetland Status: none

Description: Perennial **herb** with slightly ribbed **stems**, 1½ - 5 feet (50 - 150 cm) tall, often fallen over during flowering. **Leaves** 2 - 4 inches (5 - 10 cm) long, with toothed margins, pointed tips, and rounded or squared-off bases; opposite at midstem, sometimes alternate near the top of the stem; lower surfaces of leaves slightly hairy and dotted with glands. **Flower heads** up to ¾ inch (2 cm) across, with long, purplish bristles and 35 - 45 purplish-pink **disc flowers**; no **ray flowers**, although long, pink style branches may resemble ray flowers. **Fruit** less than ¼ inch (5 mm) long, seed-like, surrounded by barbed, purplish bristles. **Bracts** surrounding the base of the flower heads are strongly ribbed.

Similar Species: Eastern false-boneset (*Brickellia eupatorioides*) leaves are narrow with wedge-shaped bases and flower heads with white bristles. Ageratum (*Ageratina aromatica*), with white flowers, and mistflower (*Conoclinium coelestinum*), with blue flowers, have similar leaves but smaller flower heads. Young sprouts of American beauty-berry (*Callicarpa americana*), with opposite, toothed leaves, resemble immature heartleaf brickellia, but beauty-berry leaves are wedge-shaped at the base.

Related Rare Species: None in Georgia.

Habitat: Moist pine-oak-hickory woods and flats; dry woods, often with southern red oak and loblolly pine; sandy, well drained riverbanks; upper ravine slopes with spruce pine, southern magnolia, and white oak; often in basic soils. Usually in sunny openings or along edges of roads and trails; may persist in shady, overgrown woods and disturbed areas.

Life History: Little is known about the life history of heartleaf brickellia. It is a perennial herb that dies back to the ground in the winter, producing new growth in the spring and flowering in late summer and fall. Its flowers are visited by a variety of insect pollinators, including butterflies, bees, and flies. The seeds are dispersed by wind, gravity, and animals; bristles surrounding the seeds probably aid in seed dispersal by both wind and animals.

Survey Recommendations: Surveys are best conducted during flowering (late August–October).

Range: Georgia, southeast Alabama, and north Florida.

Threats: Conversion of habitat to pine plantations, pasture, and developments, fire suppression in upland woods, logging and other mechanical clearing in ravines and on slopes.

Georgia Conservation Status: About 17 populations have been seen in the last 20 years; 15 of these occur on Fort Benning, the others on private land.

Conservation and Management Recommendations: Apply prescribed fire every 3 - 5 years during the growing season in pine-oak woods, allowing fire to burn into upper slopes of ravines. Avoid clearcutting and conversion of upland woods to pine plantations.

Selected References:

Chafin, L.G. 2007. Field guide to the rare plants of Georgia. State Botanical Garden of Georgia and University of Georgia Press, Athens.

Chafin, L.G. 2000. Field guide to the rare plants of Florida. Florida Natural Areas Inventory, Tallahassee.

Cronquist, A. 1980. Vascular flora of the southeastern United States, Vol. 1, Asteraceae. University of North Carolina Press, Chapel Hill.

FNA. 2006. Flora of North America. Vol. 22, Magnoliophyta: Asteridae, Part 8: Asteraceae, Part 3. Oxford University Press, New York.

Kral, R. 1983. A report on some rare, threatened, or endangered forest-related vascular plants of the South. Technical Publication R8-TP2. United States Forest Service, Atlanta.

NatureServe. 2007. NatureServe Explorer. Arlington, Virginia.
<http://www.natureserve.org/explorer>

Weakley, A.S. 2007. Flora of the Carolinas, Virginia, Georgia, and surrounding areas. University of North Carolina Herbarium, Chapel Hill.

Author of species account: Linda G. Chafin

Date Compiled or Updated:

L. Chafin, Feb. 2008: original account

K. Owers, Jan. 2010: updated status and ranks, added pictures




©Hugh and Carol Nourse

Inflorescence

HEARTLEAF BRICKELLIA
Brickellia cordifolia



J.C. Putnam H.





Inflorescence



©Hugh and Carol Nourse

Inflorescence