



Common Name: LEAN CRAYFISH

Scientific Name: *Cambarus (Depressicambarus) strigosus* Hobbs

Other Commonly Used Names: none

Previously Used Scientific Names: none

Family: Cambaridae

Rarity Ranks: G2/S2

State Legal Status: Threatened

Federal Legal Status: none

Description: The overall color of the lean crayfish is bluish-olive with the margins of the rostrum creamy orange. The areola is very narrow to non-existent and the abdomen appears much narrower than the cephalothorax. The claws may be robust. This species reaches a maximum total body length of about 75 mm (3 inches).

Similar Species: The variable crayfish (*Cambarus latimanus*) may occur in close proximity to this species, but it is a dull brownish color and there is space within the areola.

Habitat: Complex burrows adjacent to streams or in low areas where the water table is near the surface of the ground.

Diet: Crayfishes are considered opportunistic omnivores that will consume virtually any living or dead organic matter that they find or can capture. Night video of burrowing

crayfishes indicates they may also be active predators of invertebrates that venture close to their burrow opening.

Life History: Burrowing crayfishes inhabit a system of tunnels that may be very complex with several openings to the surface. Openings to the tunnels are often marked by piles of dirt or mud pellets (chimneys). Depending on the soil type and moisture content, these chimneys can reach heights of 15 cm (6 inches) or more. These crayfishes are typically confined to their burrows, but a male must leave its burrow to search for females during the reproductive season. As mentioned above, they may also forage near the opening of their burrow. Active burrows with fresh soil are seen from late spring to late fall, particularly after rain events. During the dry part of the summer, burrow openings may be plugged to help conserve moisture in the burrow. Reproduction probably occurs during the spring and fall, but males in reproductive condition may be found at any time during the year. It is very rare to find more than one adult crayfish in the same burrow. When a female crayfish releases her eggs, she attaches them to her swimmerets and is said to be “in berry.” Upon hatching, the juvenile crayfish are attached to the mother by a thread. After the juveniles molt for the second time, they are free of the mother, but stay close and will hold on to her for some time. Multiple juveniles are occasionally found in a single burrow. Eventually they move off on their own. Crayfishes molt 6 or 7 times during their first year of life and most are probably able to reproduce by the end of that year. They molt once or twice a year for the remainder of their lives. Although it is difficult to study burrowing crayfishes, some researchers believe they may live as long as 10 years. Male lean crayfish in reproductive condition have been collected in May and October and females with eggs have been collected in April. Clutch sizes for these females ranged from 31-39 eggs, and their egg diameters ranged from 2.0-2.1 mm (about 1/8 inch).

Survey Recommendations: Burrowing crayfishes may be collected by direct excavation of their burrows, by trapping, and during night surveys. Excavating burrows is time consuming and can be very difficult. It also results in destruction of the animals' burrow. Traps made with PVC pipes or mist nets can be effective. Burrowing crayfishes are sometimes captured around the openings of their burrows on damp nights. Active burrows are found from about mid-March to mid-November if the water table is within about 2 feet of the surface of the ground.

Range: The lean crayfish is currently known from about 10 locations in the Broad River and Little River systems (Savannah River drainage) in northeastern Georgia. It is known from Wilkes, Oglethorpe, and Elbert Counties, all of which are part of the Piedmont physiographic province.

Threats: The small range of this species makes it vulnerable to land disturbing activities around streams and wetlands.

Georgia Conservation Status: A large population is known from Nancy Hart State Park.

Conservation and Management Recommendations: Areas with burrows should be protected from land disturbing activities. Additional surveys and life history studies are needed to better define the range of the lean crayfish and help predict its response to environmental change. Environmental education programs should include information about burrowing crayfishes and encourage protection of burrows.

Selected References:

Hobbs, H. H., Jr. 1981. The crayfishes of Georgia. *Smithsonian Contributions to Zoology* 318:1-549.

Hobbs, H. H., Jr. 1989. An illustrated checklist of the American crayfishes (Decapoda: Astacidae, Cambaridae, and Parastacidae). *Smithsonian Contributions to Zoology* 480:1-236.


Skelton, C.E. 2008. Crayfish surveys for Elba Express Pipeline Project. Unpublished report to ENSR. 19 pp.

Skelton, C.E., S. Cammack, and E. Van De Genachte. 2002. Surveys of rare burrowing crayfish species. Unpublished report to the Georgia Department of Natural Resources. 13 pp.

Taylor, C. A., G. A. Schuster, J. E. Cooper, R. J. DiStefano, A. G. Eversole, P. Hamr, H. H. Hobbs III, H. W. Robison, C. E. Skelton, and R. F. Thoma. 2007. A reassessment of the conservation status of crayfishes of the United States and Canada after 10+ years of increased awareness. *Fisheries* 32(8): 72-389.

Author of Species Account: Christopher E. Skelton

Date Compiled or Updated: June 2008



Watersheds (Huc 10) with known occurrences. Streams, county lines, and major river basin boundaries are also shown. Map generated from GADNR (Nongame Conservation Section) data on December 18, 2008.