

POLYLOGARITHMS AND ALGEBRAIC K -THEORY SEMINAR

The goal of this seminar, organized by Andrew Blumberg and Soren Galatius, is to understand the preprint [KRS26a], by Kupers, Rudenko, and Sierra. Time permitting, we also hope to discuss the sequel [KRS26b], which gives interesting applications to number fields.

This tentative schedule may be updated as we go (for instance, adding references). Current version: September 4, 2026.

Week 1 *Introduction.*

Broad overview of what we'll try to cover. Some history and motivation.

Week 2 *Derived indecomposables 1. (Optional historical perspective.)*

Quillen defined $D_q(B/A, M)$ for a commutative ring A , a commutative A -algebra B , and a B -module M . Sketch his definition using simplicial resolutions, explain a few basic properties [Qui, §2–4]. Then discuss non-unital commutative k -algebras I , and their unitalization $I^+ = I \oplus k$. Discuss exactness properties of the functor $I \mapsto D_*(I^+/k, k)$, which is the analogy in Quillen's setting to the E_k -homology functors we'll discuss in the next lecture. If there is time, explain in what sense this is the derived functor of $I \mapsto I/I^2$.

Then discuss the case when k is a field of characteristic zero, as in [Qui, §11]. By Theorem 11.7 of op.cit., the linear dual $D^*(k/I^+, k)$ is canonically identified with the primitives of a co-commutative Hopf algebra, and hence acquires the structure of a graded Lie algebra.

Week 3 *Derived indecomposables 2.*

Briefly recall what a (non-unital) E_k -algebra is, in spaces and other symmetric monoidal categories. Explain “indecomposables” as left adjoint functor to “trivial algebra”, following [GKRW25a, §12–13], [Fra13], and/or [Heu24, §6]. Denoting the (derived) indecomposables of a non-unital E_k -algebra A by $Q_{\mathbb{L}}^{E_k}(A)$, define E_k -homology as $H_*^{E_k}(A) = \tilde{H}_*(Q_{\mathbb{L}}^{E_k})$. Explain the relationship to k -fold bar constructions.

Discuss special features in the case of an E_{∞} algebra in $\mathcal{D}(\mathbb{Q})$, in particular that homology of the 1-fold bar construction has the structure of a commutative Hopf algebra, and that the indecomposables of this Hopf algebra (in the sense of Milnor–Moore) can be identified with E_{∞} -homology, up to a shift. This explains the Lie coalgebra structure on E_{∞} -homology.

If time permits, the explicit 1-categorical model B^{Com} from [KRS26a, Definition C.13] could be discussed.

Week 4 *The Quillen spectral sequence and the Rognes spectral sequence.*

Given an E_1 algebra M in spaces, for instance a topological monoid, and an E_1 map $M \rightarrow \mathbb{N}$, we can regard M as an E_1 algebra in based spaces and the derived indecomposables form an $\mathbb{N}$ -graded based space, whose homology groups are the bigraded homology groups $H_{*,*}^{E_1}(M)$. In this situation there is also the usual bar construction $BM = |[p] \mapsto M^p|$, which is just a plain space. Briefly explain (if this wasn't done in the previous talk) that the E_1 -map $M \rightarrow \mathbb{N}$ gives rise to a filtration on BM whose associated graded is the suspension of $Q_{\mathbb{L}}^{E_1} M$, so there results a spectral sequence $H_{s,t-1}^{E_1}(M) \Rightarrow H_{s+t}(BM)$ in this situation, see for instance [GKRW25a, §13.8]. Explain or state that with field coefficients, this is

a spectral sequence of coalgebras, converging to the “cup coproduct” on homology (linear dual to cup product on cohomology).

Then explain the corresponding statement when M is an E_∞ algebra, with group completion $\Omega^\infty M^{\text{gp}}$ for a spectrum M^{gp} . The spectrum homology $H_*(M^{\text{gp}})$ is the abutment of a spectral sequence whose first page is $H_{*,*}^{E_\infty}(M)$. Over a field of characteristic 0, this spectral sequence can be identified, up to a shift, with the Hopf algebra indecomposables of the spectral sequence for the underlying E_1 algebra. For general Hopf algebra reasons, the indecomposables have a canonical Lie coalgebra structure (this goes back to at least Milnor–Moore’s paper on Hopf algebras, the linear dual statement is that the primitives in a cocommutative Hopf algebra inherit a Lie algebra structure).

This could also be a good time to state the formula in [KRS26a, §9.4] for the d^1 -differential in the Rognes spectral sequence, which uses the Lie cobracket.

We will be especially interested in the case $M = \coprod BGL_n(F)$ whose group completion is the K -theory spectrum $K(F)$. Summarize some known results about rational K -theory, including Borel’s result when F is the ring of integers in a number field or a localization of such a ring. It would be nice to also say something about Borel and Yang’s complete calculation of $H^*(BGL_n(F); \mathbb{Q})$ in [BY94, section 7], but there might not be time for it in this lecture.

Week 5 *The Goncharov Lie coalgebra.*

The first goal of this lecture is to give the definition of $\mathcal{G}(F) = \bigoplus_n \mathcal{G}_n(F)$, in terms of E_∞ homology of $M = \coprod_n BGL_n(F)$. State the vanishing result in [GKRW25b, Corollary 6.12], motivating the definition of $\mathcal{G}_n(F)$ as the lowest interesting value of $H_{n,*}^{E_\infty}(M)$.

The rest of this lecture could be spent on a hands-on exploration of $\mathcal{G}_n(F)$ for $n \in \{1, 2\}$, both of which can be deduced from the definition without a lot of theory. See [GKRW25b, §9.2] regarding $\mathcal{G}_2(F) = H_{2,3}^{E_\infty}(M; \mathbb{Q})$ in terms of the Bloch group $B_2(F)$. (This group $B_2(F)$ is denoted $\mathfrak{p}(F)$ and called the “pre-Bloch” group in [GKRW25b], but we will follow [KRS26a] and call it simply the “Bloch group”.) The mathematical content in the case $n = 2$ largely goes back to [Sus90]. The connection between $B_2(F)$ and $\mathcal{G}_2(F)$ is explained in a different way in [KRS26a, §7.2], using a presentation of $\mathcal{G}_2(F)$ as a rational vector space which we will discuss later in the seminar.

Week 6 *Polylogarithms and multiple polylogarithms 1.*

Begin by introducing these as holomorphic functions

$$\text{Li}_{n_1, \dots, n_k} : \mathbb{D}^k \rightarrow \mathbb{C},$$

where $\mathbb{D} \subset \mathbb{C}$ is the open unit disk and $\mathbb{D}^k = \mathbb{D} \times \dots \times \mathbb{D}$. Perhaps the rest of this lecture could be about the special case $k = 1$, the so-called “classical polylogarithms”. In particular, $\text{Li}_1(z) = -\log(1 - z)$ and $\text{Li}_2(z)$ is called the dilogarithm and was introduced by Euler.

The function $\text{Li}_n(z)$ can be holomorphically continued to a *multi-valued* function on $\mathbb{C} \setminus \{0, 1\}$: each determination of $\text{Li}_n(z) \in \mathbb{C}$ depends on a choice of homotopy class of path in $\mathbb{C} \setminus \{0, 1\}$ from z to a point in the open interval $(0, 1)$, but it is well understood how the value depends on the path (“monodromy”, see for instance [BD94, §1.1] or [Hai94, §2]). We will not make much actual use of these multivalued “functions”, but there are several ways to replace them by single-valued functions. For instance, the real

part of $\text{Li}_1(z)$ is $-\log|1-z|$, which is a single-valued real-analytic function $\mathbb{C} \setminus \{1\} \rightarrow \mathbb{R}$.

Explain the Bloch–Wigner function $D_2(z) = \text{Im}(\text{Li}_2(z)) + \arg(1-z) \log|z|$, a single-valued function $\mathbb{C} \setminus \{0, 1\} \rightarrow \mathbb{R}$ extracted from Li_2 , and state the functional equation it satisfies. Define, or recall from a previous lecture, the Bloch group $B_2(\mathbb{C})$ and explain how D_2 gives rise to a function $B_2(\mathbb{C}) \rightarrow \mathbb{R}$. Possible reference [Zag07, sections 2–4].

Week 7 *Multiple polylogarithms 2.*

Returning to the general case, define the “iterated integral” $I(a_0; a_1, \dots, a_k; a_{k+1})$ from e.g. [Gon01, eq.(20)] and [Dup21, eq.(38)], and state how the multi-valued functions $\text{Li}_{n_1, \dots, n_k}$ arise as special cases of the iterated integral. In this lecture, we’ll try to explain how to lift these complex numbers to elements of fancier objects. The goal is, firstly, to explain the meaning of the Lie coalgebra denoted $\mathcal{L}^{\text{Hod}}$, and secondly, to explain the meaning of $I^{\text{Hod}}(a_0; a_1, \dots, a_k; a_{k+1}) \in \mathcal{L}^{\text{Hod}}$. These will allow us to parse the explanation around [KRS26a, Theorem D.b].

Start with generalities about Lie coalgebras and commutative Hopf algebras over a field k of characteristic zero, for instance starting from Milnor–Moore (dualized): any Lie coalgebra L over k is canonically the indecomposables of a commutative Hopf algebra $H = \mathcal{U}(L)$, its universal enveloping coalgebra. The category $\mathcal{M} = \text{Comod}_H$ of comodules over H which are finite-dimensional over k is an abelian category, and $\otimes_k$ can be upgraded to a symmetric monoidal structure $\otimes_{\mathcal{M}}$ on Comod_H , such that the forgetful functor $r : \text{Comod}_H \rightarrow \text{Vect}_k$ is symmetric monoidal. In algebraic geometry language, Comod_H is by definition the category of representations of the group scheme $\text{Spec}(H)$ over $\text{Spec}(k)$. In practice, there are many situations where one *first* defines the category $\mathcal{M}$ and then uses it to extract the Hopf algebra H , equivalently the Lie coalgebra L , for which $\mathcal{M} = \text{Comod}_H$. The reconstruction of H from $(\mathcal{M}, \otimes_{\mathcal{M}}, r)$ is often summarized as “by the Tannakian formalism”, but is actually very concrete, and importantly, it lets one easily *name elements* of H : the Hopf algebra H is generated as a k -vector space by $(M, v, \phi) \in H$ where $M \in \mathcal{M}$, $v : k \rightarrow r(M)$, and $\phi : r(M) \rightarrow k$, subject to $(M, v, \phi) \in H$ being linear in both v and ϕ , as well as the equivalence relation $(M, v, \phi \circ r(f)) \sim (N, r(f) \circ v, \phi)$ for each morphism $f : M \rightarrow N$. See for instance [Gon01, Theorem 3.2] or [Dup24, §8.3].

Then briefly explain the category MHS of “mixed Hodge structures”, which is an example. State its definition, in terms of two filtrations, and basic properties. It’s somewhat surprising that it’s an abelian category! Its role in life is to fix into a diagram

$$\begin{array}{ccc} \text{Var}_F & \xrightarrow{\mathcal{H}^j} & \text{MHS} \\ \downarrow & & \downarrow r \\ \text{Top} & \xrightarrow{H^j(-;k)} & \text{Vect}_k, \end{array}$$

where $F \subset \mathbb{C}$ is a subfield (such as $F = \mathbb{C}$), the left vertical arrow sends X to $X(\mathbb{C})$ in the euclidean topology, and $k \in \{\mathbb{Q}, \mathbb{R}, \mathbb{C}\}$. The category MHS depends on k and we’ll mostly take $k = \mathbb{Q}$.

Explain the special “Tate” objects $\mathbb{Q}(n) \in \text{MHS}$, one for each $n \in \mathbb{Z}$, and the full subcategory MHTS $\subset \text{MHS}$ on those objects which are iterated extensions of Tate objects. Then, MHS is comodules over some Hopf algebra H^{MHS} , and the subcategory MHTS is comodules over a sub Hopf algebra

$H^{\text{MHTS}} \subset H^{\text{MHS}}$, which by definition is $H^{\text{MHTS}} = \mathcal{U}(\mathcal{L}^{\text{Hod}})$. We can name explicit elements of this graded Hopf algebra by the general theory: in degree m , they look like (M, v, ϕ) , with $M \in \text{MHTS}$, $v : \mathbb{Q}(0) \rightarrow \text{Gr}_0^W(M)$, and $\phi : \text{Gr}_{-2m}^W(M) \rightarrow \mathbb{Q}(m)$.

Finally, fundamental groups of certain varieties give a good source of objects of $\text{MHTS} \subset \text{MHS}$. If π is a finitely generated group, when $\mathbb{Q}[\pi]/I^{m+1}$, the group ring modulo the $(m+1)$ st power of the augmentation ideal, is a finite-dimensional vector space. It is an iterated extension of the vector spaces I^k/I^{k+1} , so there exists noncanonically an isomorphism $\mathbb{Q}[\pi]/I^{m+1} \cong \bigoplus_{k=0}^m I^k/I^{k+1}$ in rational vector spaces. Moreover, there is a canonical quotient map $S^k(I/I^2) \rightarrow I^k/I^{k+1}$ from the k th symmetric power of $I/I^2 = \pi^{\text{ab}}$, and this quotient map is an isomorphism if $H_2(\pi) = 0$. If $\pi = \pi_1(X, a)$ is the fundamental group of a variety X , then there is a formula for $\mathbb{Q}[\pi]/I^{m+1}$ as $H^m(X^m, D)$, for a certain subvariety $D \subset X^m$, see for instance [Gon01, Theorem 4.1] or the main result of [Loo24]. This formula then gives a canonical lift of $\mathbb{Q}[\pi]/I^{m+1}$ to MHS, and in interesting examples the splitting $\mathbb{Q}[\pi]/I^{m+1} \cong \bigoplus_{k=0}^m I^k/I^{k+1}$ does not exist in MHS. In many cases, $I/I^2 = H_1(X; \mathbb{Q})$ is isomorphic to a sum of copies of $\mathbb{Q}(1)$, and then $\mathbb{Q}[\pi]/I^{m+1} \in \text{MHTS} = \text{Comod}_{\mathcal{U}(\mathcal{L}^{\text{Hod}})}$. This discussion generalizes to $\pi = \pi_{\leq 1}(X)(x, y)$, the set of morphisms in the fundamental groupoid of X ; in this case, $\mathbb{Q}\pi/I^{m+1} \in \text{MHTS}$ is denoted $\mathcal{P}_M^m(X; x, y)$ in [Gon01, §4.4].

The iterated integral now lifts to an element $I^{\text{Hod}}(a_0; a_1, \dots, a_k; a_{k+1}) \in \mathcal{L}^{\text{Hod}}$ as in [Gon01, Definition 5.5].

Week 8 Steinberg modules 1.

This talk should give a hands-on perspective on $\text{St}(V)$, $\text{St}^2(V)$, and $\text{St}^\infty(V)$, with emphasis on how one names elements of these vector spaces.

Define the Tits building $T(V)$, state the Solomon–Tits theorem, define $\text{St}(V) = \tilde{H}_{n-2}(T(V); \mathbb{Q})$, and define apartment classes $[v_1, \dots, v_n] \in \text{St}(V)$ for any ordered basis $(v_1, \dots, v_n)$ of V , and state the presentation in [KRS26a, §2.3.2]. Mention the product $\text{St}(V) \otimes \text{St}(W) \rightarrow \text{St}(V \oplus W)$, under which $[v_1, \dots, v_n]$ is the product of the canonical element $[v_j] \in \text{St}(P_j)$, where $P_j = Fv_j \subset V$.

Then state the two perspectives on $\text{St}^2(V)$: as a functor, it is the bar construction of St ; and objectwise, it is just $\text{St}^2(V) = \text{St}(V) \otimes \text{St}(V)$. The comparison of these two definitions is an important result of Miller–Patz–Wilson, and leads to a comparison map $\text{St}(V) \otimes \text{St}(V) \hookrightarrow (B^{\text{As}}\text{St})_n(V)$ called the associative symbol in [KRS26a, §2.6.3]. (Remark on notation in [KRS26a]: they use “Bar” for an ∞ -categorical notion of bar-constructing and B^{As} for the usual 1-categorical model with tensor powers assembled into a simplicial object.) Define the elements $I[v_1, \dots, v_n] \in \text{St}^2(V)$ and state [GKRW25b, Theorem C], which imply that any one of these elements generate $H_0(\text{GL}(V); \text{St}^2(V)) \cong \mathbb{Q}$.

Either concurrently with the discussion of St^2 or separately, state the two (equivalent) perspectives on $\text{St}^\infty(V)$: as a quotient of $\text{St}^2(V)$ by the sum over all non-trivial decompositions $V = V_1 \oplus V_2$ of the images of the product map $\text{St}^2(V_1) \otimes \text{St}^2(V_2) \rightarrow \text{St}^2(V)$, and as the commutative Koszul dual $B^{\text{Com}}(\text{St})$. Define the Steinberg correlators $C[u_0 : \dots : u_n] \in \text{St}^\infty(V)$ for ordered affine bases $(u_0, \dots, u_n)$ of V . (If possible, it would be nice to do St^2 and St^∞ in parallel, which would perhaps require some prerequisites about associative and commutative Koszul duality. I.e., the 1-categorical models B^{As} and B^{Com} ; there’s some discussion in the appendices of [KRS26a], and perhaps the book of Loday and Vallette is also useful. On the other

hand, an extensive detour into Koszul duality theory could perhaps be a distraction in this seminar.)

Week 9 *Steinberg modules 2.*

This talk should connect the hands-on algebraic perspective from the previous talk with the derived indecomposables and E_k -homology from previous talks. Sources could include [KRS26a] and [GKRW25a]. A goal could be to state [KRS26a, Theorem 4.3] and explain some ingredients in its proof.

This talk will make use of the flexibility of studying E_k -algebras in different underlying symmetric monoidal categories $\mathcal{C}$, we'll especially be interested in $\mathcal{C}$ being a category of functors out of finite-dimensional F -vector spaces and their isomorphisms, equipped with the Day convolution monoidal structure (called the “Levi tensor product” and denoted $\boxtimes$ in [KRS26a]). Following [KRS26a, §4.2–4.4], explain or state how the “split buildings” $\tilde{D}^k(V)$ show up, using the iterated-bar-construction model for E_k -indecomposables. Mention the “non-split buildings” $D^k(V)$, essentially due to Rognes. The forgetful map $\tilde{D}^k(V) \rightarrow D^k(V)$ is not an equivalence, but state Nesterenko–Suslin’s comparison theorem. The result of Miller–Patz–Wilson and its generalization in [KRS26a, §4.4] identifies $D^k(V)$ with the bar construction of $D^{k-1}(V)$ when $k \geq 2$, and in particular an equivalence between St^2 and the bar construction of St .

Finally, explain the description of the Lie cobracket on $\bigoplus \mathcal{G}_n(F) = \bigoplus H_1(\mathrm{GL}_n(F); \mathrm{St}^\infty)$, following [KRS26a, section 4.5]. In principle it would be nice to also discuss [KRS26a, §5.3.1] and the proof of their Theorem C.b.

Week 10 *Steinberg modules 3.*

This talk will focus on $\mathrm{St}^\infty(V)$.

Explain the resolution of $\mathrm{St}^\infty(V)$ given by [KRS26a, Proposition 5.12]. Following [KRS26a, §5.2], use this resolution to deduce the presentation of $\mathcal{G}_n(F)$ as a vector space, generated by the elements $\mathrm{Cor}^{\mathcal{G}}(a_0 : \dots : a_n)$. One form of the presentation is in [KRS26a, Proposition 5.15], which is slightly reformulated in [KRS26a, §7.1].

Mention the realization constructions from [KRS26a, Theorem 7.29]. Imprecisely, this tells us that “relations between multiple polylogarithms” is related to the relations in $\mathcal{G}_n(F)$, and hence to algebraic K -theory of F via the Rognes spectral sequence.

Finally, define the depth filtration on $\mathcal{G}_n(F)$ and explain how depth-one corresponds to (one-variable) polylogarithms.

Week 11 *Number fields 1*

When F is an algebraic extension of $\mathbb{Q}$, the abelian category of mixed Tate motives has an a priori meaning. This special case has many interesting features, and a separate paper [KRS26b] is devoted to it.

The detailed construction of an abelian category of mixed Tate motives is quite involved, but the conclusion can be summarized fairly concisely, for instance following [Dup24, §4, §6]. As explained in [Dup24, §3], it comes with a “realization” functor to the category MHTS from a previous lecture for each embedding $F \rightarrow \mathbb{C}$.

State or sketch the realization functor R^{MTM} from [KRS26a, Theorem D.a], and sketch how it is used in [KRS26b] to prove that the category of mixed Tate motives over F is generated by motivic multiple polylogarithms (i.e., “Goncharov’s universality conjecture”).

Time permitting, it would be nice to say something about how [KRS26b, Corollary B], that any period of a mixed Tate motive over F can be expressed in terms of multiple polylogarithms evaluated at elements in F . [KRS26b, Corollary C] establishes a version of Zagier’s conjecture (special values of ζ -functions expressed in terms of special values of multiple polylogarithms). An earlier survey regarding Zagier’s conjecture can be found in [Dup21, §1.1–1.5].

REFERENCES

- [BD94] A. Beilinson and P. Deligne, *Interprétation motivique de la conjecture de Zagier reliant polylogarithmes et régulateurs*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Part 2, Amer. Math. Soc., Providence, RI, 1994, pp. 97–121.
- [BY94] Armand Borel and Jun Yang, *The rank conjecture for number fields*, Math. Res. Lett. **1** (1994), no. 6, 689–699.
- [Dup21] Clément Dupont, *Progrès récents sur la conjecture de Zagier et le programme de Goncharov [d’après Goncharov, Rudenko, Gangl, ...]*, no. 430, 2021, Séminaire Bourbaki. Vol. 2019/2021. Exposés 1166–1180, pp. Exp. No. 1176, 295–343.
- [Dup24] Clément Dupont, *An introduction to mixed Tate motives*, 2024.
- [Fra13] John Francis, *The tangent complex and Hochschild cohomology of $\mathcal{E}_n$ -rings*, Compositio Mathematica **149** (2013), no. 3, 430–480.
- [GKRW25a] S. Galatius, A. Kupers, and O. Randal-Williams, *Cellular E_k -algebras*, Astérisque (2025), no. 460, x+299.
- [GKRW25b] Søren Galatius, Alexander Kupers, and Oscar Randal-Williams, *E_∞ -cells and general linear groups of infinite fields*, Duke Math. J. **174** (2025), no. 14, 2927–3046.
- [Gon01] A. B. Goncharov, *Multiple polylogarithms and mixed Tate motives*, 2001.
- [Hai94] Richard M. Hain, *Classical polylogarithms*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Part 2, Amer. Math. Soc., Providence, RI, 1994, pp. 3–42.
- [Heu24] Gijs Heuts, *Koszul duality and a conjecture of Francis–Gaitsgory*, 2024.
- [KRS26a] Alexander Kupers, Daniil Rudenko, and Ismael Sierra, *The Goncharov Lie coalgebra of a field*, 2026.
- [KRS26b] ———, *Mixed Tate motives over number fields*, 2026.
- [Loo24] Eduard Looijenga, *On the motivic description of truncated fundamental group rings*, 2024.
- [Qui] Daniel Quillen, *Homology of commutative rings*, preprint, 1–81.
- [Sus90] A. A. Suslin, *K_3 of a field, and the Bloch group*, Trudy Mat. Inst. Steklov. **183** (1990), 180–199, 229, Translated in Proc. Steklov Inst. Math. 1991, no. 4, 217–239, Galois theory, rings, algebraic groups and their applications (Russian).
- [Zag07] Don Zagier, *The dilogarithm function*, Frontiers in number theory, physics, and geometry. II, Springer, Berlin, 2007, pp. 3–65.