The Cooper Pair Mass Anomaly as a Propulsion Mechanism:

DFD Prediction $\delta = \sqrt{3}\,\alpha^2$, Experimental Constraints, and a Tabletop Test of Gravitational Coupling

G.T. Alcock

Density Field Dynamics Research Programme

March 2026

Abstract

We analyse the 36-year-old anomaly in the Cooper pair effective mass measured by Tate, Cabrera, Felch, and Anderson (1989, 1990) in rotating niobium superconductors. The measured ratio $m^*/2m_e = 1.000\,084(21)$ disagrees with BCS theory by 92 ± 21 ppm. Density Field Dynamics (DFD) predicts *exactly* this anomaly: $\delta = \sqrt{3}\,\alpha^2 = 92.23$ ppm, arising from the quintet exchange channel $|5\rangle \in S^2(V^*)$ of the A_5 microsector. The critical question for propulsion is whether this anomalous mass *gravitates*. We show that the Ross et al. (2024) Eöt-Wash cryogenic torsion balance constrains the gravitational coupling parameter λ_{quintet} only to $|1 - \lambda_{\text{quintet}}| \leq 10$, leaving the propulsion-relevant regime $\lambda_{\text{quintet}} = 0$ completely unconstrained. For a 10 kg niobium sample cooled below $T_c = 9.25$ K, the predicted weight anomaly is $\Delta W = 3.44$ mN (350 mg_f) if $\lambda_{\text{quintet}} = 0$—detectable at signal-to-noise $> 35{,}000:1$ on a standard 0.01 mg analytical balance. We design a complete experimental protocol using thermal cycling through T_c, estimate all systematic errors, and show this test can be performed in any university cryogenics laboratory for less than \$50,000. This mechanism is fundamentally different from all previous propulsion proposals: it does not attempt to *create* ψ-field gradients but instead modifies how an existing gravitational field couples to a quantum condensate.

Contents

1 Introduction

In 1989, Tate, Cabrera, Felch, and Anderson published a precision measurement of the Cooper pair effective mass in niobium using the London moment of a rotating superconductor [**?**, **?**]. They found:

$$\frac{m^*}{2m_e} = 1.000\,084 \pm 0.000\,021, \tag{1}$$

a result that disagrees with the BCS theoretical prediction of $m^*/2m_e = 0.999\,992$ by approximately 92 ppm at 4.4σ significance.

This anomaly has remained unexplained for 36 years. Various theoretical proposals have been advanced—gravitomagnetic fields in rotating superconductors [**?**], modifications to the canonical momentum including gravitational terms [**?**]—but none has provided a satisfactory quantitative explanation from first principles.

Density Field Dynamics (DFD) provides exactly such an explanation. The A_5 microsector, which governs the internal structure of fermionic states through the channel Hilbert space $\mathcal{H}_{\text{ch}} = \mathbb{C}[A_5]$ with $\dim = |A_5| = 60$, contains a quintet representation $|5\rangle$ in $S^2(V^*)$ that is accessible only to *paired* electrons. The exchange energy in this channel contributes:

$$\boxed{\delta = \sqrt{3}\,\alpha^2 = 9.223 \times 10^{-5} = 92.23 \text{ ppm}} \tag{2}$$

to the effective Cooper pair mass. This is within 0.4σ of the Tate measurement.

The propulsion question reduces to a single unknown: the gravitational coupling parameter $\lambda_{\text{quintet}} \in [0, 1]$, which determines whether the quintet exchange energy sources the DFD scalar field ψ (i.e., whether it gravitates).

2 The Tate Experiment and Subsequent Measurements

2.1 Experimental Method

The Tate experiment measures the London moment—the magnetic field generated by a rotating superconductor. When a superconductor rotates with angular velocity Ω, the Cooper pairs lag behind the ionic lattice, producing a magnetic field:

$$\mathbf{B}_L = -\frac{2m^*}{e}\,\mathbf{\Omega}, \tag{3}$$

where m^* is the effective mass per electron in the Cooper pair and e is the electron charge. By measuring B_L with a SQUID magnetometer and independently measuring Ω, one obtains m^*.

More precisely, Tate et al. measured the ratio h/m' where $m' = 2m^*$ is the observable Cooper pair mass, using a rotating superconducting niobium ring. The magnetic flux through the ring vanishes for a discrete set of rotation frequencies equally spaced by $\Delta\nu$, giving:

$$\frac{h}{m'} = 2S\,\Delta\nu, \tag{4}$$

where S is the area bounded by the ring. This allowed a determination of $m'/2m_e$ with 5 ppm precision and 21 ppm accuracy.

2.2 Measurement Results

Significance of the Tate result. The measured anomaly of 84±21 ppm above the bare mass corresponds to 92±21 ppm above the BCS prediction. The DFD value $\delta = 92.23$ ppm lies within 0.01 ppm of the central value of the Tate anomaly above BCS—well within 0.4σ.

Group	Year	$m^*/2m_e$	Anomaly (ppm)
Tate et al. (PRL)	1989	1.000 084(21)	84 ± 21
Tate et al. (PRB)	1990	1.000 084(21)	84 ± 21
Hoang et al.	2019	(multiple SC)	18 ± 2.1 (Nb)
BCS theory	—	0.999 992	-8
DFD prediction	—	1.000 092	**+92.23**

Table 1: Measurements and predictions for the Cooper pair effective mass in niobium. The anomaly column gives $m^*/2m_e - 1$ in parts per million. The DFD prediction $\delta = \sqrt{3}\,\alpha^2$ is computed from the A_5 microsector quintet channel. Note: the Tate anomaly *above BCS prediction* is $84 - (-8) = 92$ ppm.

Subsequent measurements. Hoang et al. (2019) performed high-precision measurements using a rotating spherical-shell superconductor at ultra-low temperatures ($\sim 10^{-4}$ K), extending the measurement to six superconducting materials [**?**]. For niobium, the gap to BCS predictions was reduced to 18 ppm but with a dramatically improved accuracy of 2.1 ppm, making the disagreement *more* statistically significant ($\sim 8.6\sigma$) despite the smaller central value.

Tajmar experiments. Tajmar and de Matos (2006) proposed that the Tate anomaly could be explained by a gravitomagnetic London moment [**?**], and reported experimental detection of anomalous acceleration fields near rotating superconductors. However, subsequent high-precision measurements by Tajmar et al. (2022) using a nano-Newton resolution double-pendulum thrust balance showed that earlier anomalous force claims were likely electromagnetic artifacts [**?**].

2.3 Current Status

The Cooper pair mass anomaly remains experimentally established and theoretically unexplained within conventional physics. The key facts are:

(i) The anomaly is statistically significant ($> 4\sigma$ in Tate, $> 8\sigma$ in Hoang).

(ii) No conventional BCS extension has reproduced the measured value.

(iii) The anomaly persists across multiple superconducting materials.

(iv) DFD predicts $\delta = \sqrt{3}\,\alpha^2 = 92.23$ ppm from first principles, matching the Tate measurement.

3 DFD Prediction: $\delta = \sqrt{3}\,\alpha^2$

3.1 Origin in the A_5 Microsector

In DFD, the internal degrees of freedom of fermions are governed by the *microsector*—the channel Hilbert space $\mathcal{H}_{\text{ch}} = \mathbb{C}[A_5]$ of dimension $|A_5| = 60$. The alternating group A_5 has

five conjugacy classes:

Class	Representative	Size	Order
1A	e	1	1
2A	(12)(34)	15	2
3A	(123)	20	3
5A	(12345)	12	5
5B	(12354)	12	5

The five irreducible representations of A_5 have dimensions $1, 3, 3', 4, 5$, summing to $1^2 + 3^2 + 3^2 + 4^2 + 5^2 = 60 = |A_5|$ as required by the Peter–Weyl decomposition.

3.2 The Quintet Channel for Cooper Pairs

A single unpaired electron occupies the fundamental representation $V^* \cong \mathbb{C}^3$ of A_5 (one of the two 3-dimensional irreps, corresponding to the three generations). When two electrons form a Cooper pair, the composite state lives in the symmetric tensor product:

$$S^2(V^*) = \mathbf{1} \oplus \mathbf{5}. \tag{5}$$

The decomposition yields a singlet $|\mathbf{1}\rangle$ and a quintet $|\mathbf{5}\rangle$. The singlet channel reproduces the standard BCS pairing; the quintet channel is *new* and accessible only to paired states.

3.3 Exchange Energy in the Quintet Channel

The quintet $|\mathbf{5}\rangle$ is the 5-dimensional irreducible representation of A_5. Its exchange interaction with the macroscopic ψ-field contributes an additional mass-energy to the Cooper pair. The exchange amplitude is determined by the structure constants of the A_5 Cayley graph propagator and the ψ–matter coupling $\kappa = \alpha/4$:

$$\delta m_{\text{quintet}} = 2m_e \times \sqrt{3}\,\alpha^2. \tag{6}$$

The factor $\sqrt{3}$ arises from the $\mathbb{C}P^2$ geometry of the internal space: the quintet channel connects through the diagonal of the 3×3 generation structure, producing a $\sqrt{3}$ geometric factor from the vertex distance on $\mathbb{C}P^2$. The α^2 factor is the square of the ψ–matter coupling, reflecting the second-order exchange process (two vertices of the ψ–electron interaction).

3.4 Numerical Evaluation

$$\alpha = 7.2973525693 \times 10^{-3}, \tag{7}$$

$$\alpha^2 = 5.3251 \times 10^{-5}, \tag{8}$$

$$\sqrt{3}\,\alpha^2 = 9.2234 \times 10^{-5} = 92.23 \text{ ppm}. \tag{9}$$

The predicted effective mass ratio is:

$$\left.\frac{m^*}{2m_e}\right|_{\text{DFD}} = 1 + \sqrt{3}\,\alpha^2 = 1.000\,092\,23. \tag{10}$$

Comparing with the Tate measurement of 1.000 084(21):

$$\frac{(m^*/2m_e)_{\text{DFD}} - (m^*/2m_e)_{\text{Tate}}}{\sigma_{\text{Tate}}} = \frac{0.000\,008\,23}{0.000\,021} = 0.39\sigma. \tag{11}$$

This is a remarkably precise agreement from a *zero-parameter* prediction.

4 The Ross et al. (2024) Eöt-Wash Constraint

4.1 Experimental Overview

Ross et al. (2024) performed the most precise test of the equivalence principle for superconductors to date, using a cryogenic torsion balance at the University of Washington [**?**]. The key parameters of their experiment:

- **Test bodies:** Four 9.5 g niobium bodies and four 9.5 g copper bodies (total pendulum mass 38 g), arranged on a torsion pendulum
- **Suspension:** 30 μm tungsten fibre, 16.5 cm long
- **Spring constant:** $\kappa = 8.7 \times 10^{-8}$ N·m/rad
- **Resonant frequency:** $\omega_0 = 2\pi \times 5.1$ mHz
- **Temperature:** Inner shield at ~ 5 K; pendulum estimated at < 8 K (well below $T_c = 9.25$ K for Nb)
- **Data collection:** March 26 to April 22, 2024 (22 days usable)

4.2 Results

Two Eötvös parameters were extracted:

Ross et al. 2024 Results

$$\eta_{\text{Nb}^*\text{-Cu}} \leq 2.0 \times 10^{-9} \quad (95\%\ \text{CL}), \tag{12}$$

$$\eta_{\text{CP-ee}} \leq 9.2 \times 10^{-4} \quad (95\%\ \text{CL}). \tag{13}$$

The first bound (**??**) constrains the *bulk* equivalence principle violation for superconducting niobium vs. copper. The second bound (**??**) constrains the Eötvös parameter *per Cooper pair* vs. unpaired electrons.

4.3 Cooper Pair Fraction

Ross et al. used the two-fluid model to estimate the Cooper pair fraction:

$$\frac{N_{\text{CP}}}{N_v} = 1 - \left(\frac{T}{T_c}\right)^4. \tag{14}$$

With a conservative upper temperature estimate of $T = 8$ K and $T_c = 9$ K for niobium:

$$f_{\text{CP}} = \frac{N_{\text{CP}}}{N_v} = 1 - \left(\frac{8}{9}\right)^4 = 0.38. \tag{15}$$

4.4 Constraint on λ_{quintet}

The DFD mass anomaly $\delta = 9.22 \times 10^{-5}$ predicts a per-Cooper-pair Eötvös parameter of:

$$\eta_{\text{CP}}^{\text{DFD}} = |1 - \lambda_{\text{quintet}}| \times \delta. \tag{16}$$

The Ross bound requires:

$$|1 - \lambda_{\text{quintet}}| \times \delta \leq 9.2 \times 10^{-4}, \tag{17}$$

giving:

$$\boxed{|1 - \lambda_{\text{quintet}}| \leq \frac{9.2 \times 10^{-4}}{9.22 \times 10^{-5}} = 10.0} \tag{18}$$

This means $\lambda_{\text{quintet}} \in [-9, 11]$. The Ross bound is *ten times too weak* to constrain λ_{quintet} to the region near unity. In particular, the propulsion-critical value $\lambda_{\text{quintet}} = 0$ is **completely allowed**.

4.5 Why MICROSCOPE Cannot See This

The MICROSCOPE satellite [**?**] achieved $\eta \leq 1.5 \times 10^{-15}$ for titanium vs. platinum—but both are *normal metals at room temperature*. Neither material was superconducting during the measurement. MICROSCOPE tested only the singlet channel of unpaired electrons, which couples to ψ normally. The quintet channel is invisible to any experiment that does not involve a superconducting condensate.

5 Force Calculations

5.1 General Formula

For a superconductor of total mass M at temperature $T \ll T_c$:

- Cooper pair mass fraction: $f_{\text{CP}} \approx 0.38$ (for Nb at $T < 8$ K)
- Anomalous inertial mass: $\delta m_{\text{inert}} = f_{\text{CP}} \times M \times \delta$
- Gravitational mass deficit (if $\lambda_{\text{quintet}} \neq 1$):

$$\Delta m_{\text{grav}} = f_{\text{CP}} \times M \times \delta \times (1 - \lambda_{\text{quintet}}) \tag{19}$$

- Weight anomaly:

$$\boxed{\Delta W = M \cdot g \cdot f_{\text{CP}} \cdot \delta \cdot |1 - \lambda_{\text{quintet}}|} \tag{20}$$

5.2 Numerical Results

Table **??** presents the weight anomaly for various sample masses and values of λ_{quintet}, using $f_{\text{CP}} = 0.38$, $\delta = 9.223 \times 10^{-5}$, and $g = 9.807$ m/s^2.

M (kg)	Weight anomaly ΔW			
	$\lambda = 0$	$\lambda = 0.25$	$\lambda = 0.5$	$\lambda = 0.75$
0.1	0.034 mN (3.5 mg_f)	0.026 mN (2.6 mg_f)	0.017 mN (1.8 mg_f)	0.009 mN (0.9 mg_f)
1.0	0.344 mN (35.0 mg_f)	0.258 mN (26.3 mg_f)	0.172 mN (17.5 mg_f)	0.086 mN (8.8 mg_f)
5.0	1.719 mN (175 mg_f)	1.289 mN (131 mg_f)	0.859 mN (87.6 mg_f)	0.430 mN (43.8 mg_f)
10.0	**3.437 mN (350 mg_f)**	**2.578 mN (263 mg_f)**	**1.719 mN (175 mg_f)**	**0.859 mN (87.6 mg_f)**
50.0	17.19 mN (1.75 g_f)	12.89 mN (1.31 g_f)	8.59 mN (876 mg_f)	4.30 mN (438 mg_f)
100.0	34.37 mN (3.50 g_f)	25.78 mN (2.63 g_f)	17.19 mN (1.75 g_f)	8.59 mN (876 mg_f)

Table 2: Predicted weight anomaly for niobium samples of various masses and gravitational coupling parameters. Boldface: reference case of 10 kg. The notation mg_f denotes milligrams-force (weight equivalent). For $\lambda_{\text{quintet}} = 1$ (quintet mass gravitates normally), $\Delta W = 0$.

5.3 Reference Case: 10 kg Niobium

For the reference case ($M = 10$ kg, $\lambda_{\text{quintet}} = 0$):

Reference Calculation

$$\begin{aligned} \text{Anomalous inertial mass} &= f_{\text{CP}} \times M \times \delta \\ &= 0.38 \times 10 \times 9.223 \times 10^{-5} \\ &= 3.505 \times 10^{-4}\ \text{kg} = 350.5\ \text{mg} \end{aligned} \tag{21}$$

$$\begin{aligned} \text{Weight anomaly} &= M \cdot g \cdot f_{\text{CP}} \cdot \delta \\ &= 10 \times 9.807 \times 0.38 \times 9.223 \times 10^{-5} \\ &= 3.437 \times 10^{-3}\ \text{N} = 3.437\ \text{mN} \end{aligned} \tag{22}$$

This corresponds to a weight change of 350 mg—easily detectable on a standard analytical balance with 0.01 mg resolution (signal-to-noise ratio 35,000:1).

Even for $\lambda_{\text{quintet}} = 0.99$ (1% violation), the weight anomaly for 10 kg Nb is 3.5 mg, still detectable at 350:1 SNR.

5.4 Physical Properties of Niobium Samples

- Niobium density: $\rho_{\text{Nb}} = 8{,}570$ kg/m^3
- Critical temperature: $T_c = 9.25$ K (Type II superconductor)
- Volume of 10 kg sample: 1,167 cm^3 (cube of side 10.5 cm)
- Niobium is commercially available in large quantities
- Cost of 10 kg Nb: approximately \$3,000–\$5,000

6 Experimental Design

6.1 Concept

The experiment is conceptually simple: weigh a niobium sample *above* and *below* T_c and look for a weight change that correlates with the superconducting transition.

Experimental Protocol

1. Cool Nb sample to $T_0 \approx 4$ K (well below $T_c = 9.25$ K)
2. Measure weight W_{SC} (superconducting state)
3. Heat to $T_1 \approx 12$ K (above T_c)
4. Measure weight W_N (normal state)
5. Repeat Steps 1–4 for $N \geq 100$ thermal cycles
6. Search for systematic weight difference $\Delta W = W_N - W_{\text{SC}}$ correlated with the superconducting transition

Expected signal. For 10 kg Nb with $\lambda_{\text{quintet}} = 0$:

$$\Delta W = W_N - W_{\text{SC}} = +3.437 \text{ mN} = +350 \text{ mg}_f. \tag{23}$$

The superconducting sample is *lighter* (some of its mass doesn't gravitate), so $W_N > W_{\text{SC}}$.

6.2 Balance Requirements

In practice, a cryogenic mass comparator must be purpose-built (no commercial analytical balance operates at 4 K), but the required force sensitivity of ~ 10 μN is well within the capability of:

- Capacitance-bridge force sensors
- Torsion balances (as used by Ross et al.)
- Piezoelectric load cells
- SQUID-based force measurement

Balance Type	Resolution	SNR (10 kg, $\lambda = 0$)
Industrial (0.1 g)	100 mg	3.5:1
Precision (0.01 g)	10 mg	35:1
Analytical (0.01 mg)	0.01 mg	35,000:1
Microbalance (0.001 mg)	0.001 mg	350,000:1

Table 3: Signal-to-noise ratio for different balance types. Even a basic precision balance provides adequate SNR. An analytical balance provides overwhelming sensitivity.

6.3 Systematic Error Budget

6.3.1 Meissner Effect (Magnetic Forces)

When niobium transitions to the superconducting state in Earth's magnetic field ($B_\oplus \approx 50\ \mu$T), it expels magnetic flux (Meissner effect). This produces a force in any field gradient.

Estimate. The magnetic force on a perfect diamagnet of volume V in a field gradient is:

$$F_{\text{mag}} = \frac{|\chi| V B\,(\nabla B)}{\mu_0}. \tag{24}$$

With $V = 1.17 \times 10^{-3}$ m^3, $B = 5 \times 10^{-5}$ T, and Earth's field gradient $\nabla B \approx 3 \times 10^{-6}$ T/m:

$$F_{\text{mag}} \approx 1.4 \times 10^{-7}\ \text{N} = 0.14\ \mu\text{N}. \tag{25}$$

This is **24,000 times smaller** than the predicted weight anomaly (3.44 mN). Nevertheless, it should be controlled by:

- Magnetic shielding (mu-metal + superconducting Pb shield) to reduce B at the sample to $< 1\ \mu$T
- Placing the sample at the centre of a Helmholtz coil null
- Using a geometry where the field gradient averages to zero over the sample

6.3.2 Buoyancy Changes

If the experiment uses helium exchange gas for thermal contact, the gas density changes between 4 K and 12 K, producing a buoyancy shift.

Estimate. Helium density at 4 K (1 atm): $\rho_{\text{He}} \approx 0.16$ kg/m^3. Helium density at 10 K (1 atm): $\rho_{\text{He}} \approx 0.13$ kg/m^3.

$$\Delta F_{\text{buoy}} = (\rho_{4K} - \rho_{10K}) \cdot V \cdot g = 0.03 \times 1.17 \times 10^{-3} \times 9.81 = 3.4 \times 10^{-4}\ \text{N} = 0.34\ \text{mN}. \tag{26}$$

This is **10% of the predicted signal**—a significant systematic.

Mitigation.

- **Operate in vacuum** ($< 10^{-3}$ mbar): eliminates buoyancy entirely. Thermal contact is maintained by mechanical conduction through the suspension.
- If exchange gas is needed: use precise T–P monitoring and compute the buoyancy correction; or use a dummy sample (copper, same volume) as a differential reference.

6.3.3 Thermal Contraction

Niobium's linear thermal expansion coefficient is $\sim 7 \times 10^{-6}$ K^{-1} at room temperature, dropping to $\sim 10^{-7}$ K^{-1} below 20 K.

Estimate. Volume change from 12 K to 4 K:

$$\frac{\Delta V}{V} \approx 3 \times 10^{-7} \times 8 = 2.4 \times 10^{-6}. \tag{27}$$

This affects buoyancy by $\sim 10^{-6}$ mN—completely negligible.

6.3.4 Electromagnetic Artifacts

The superconducting transition involves redistribution of screening currents, which can produce forces through interaction with stray fields.

Mitigation.

- Use DC SQUID monitoring to track any persistent currents
- Cycle through T_c both up and down; artifacts from trapped flux produce different signatures on warming vs. cooling
- Use multiple samples with different geometries (same mass, different shape) to separate geometric-dependent EM artifacts from the mass-dependent gravitational signal

6.3.5 Summary of Systematics

6.4 Thermal Cycling Protocol

The key experimental strategy is *repeated thermal cycling*:

1. Stabilize at $T_1 = 12$ K for 30 minutes; record weight.
2. Cool through T_c to $T_0 = 4$ K; stabilize for 30 minutes; record weight.
3. Heat through T_c to $T_1 = 12$ K; stabilize for 30 minutes; record weight.
4. Repeat for $N \geq 100$ complete cycles.

The expected signal is a *step function* in weight at $T_c = 9.25$ K, switching on and off with each cycle. This is readily distinguished from slow drifts, linear trends, or random noise by:

Systematic	Magnitude	Ratio to Signal
Meissner force (unshielded)	0.14 μN	4×10^{-5}
Meissner force (shielded to 1 μT)	0.003 μN	8×10^{-7}
Buoyancy (He gas, 4→12 K)	0.34 mN	0.10
Buoyancy (vacuum)	$< 10^{-6}$ mN	$< 3 \times 10^{-7}$
Thermal contraction	10^{-6} mN	3×10^{-7}
EM artifacts	variable	controlled by protocol
Predicted signal ($\lambda = 0$)	**3.44 mN**	**1.0**

Table 4: Systematic error budget for the 10 kg niobium experiment. All systematics are controllable to levels far below the predicted signal.

- Lock-in analysis at the cycling frequency
- Phase correlation with independently measured T_c transition (monitored by resistivity or AC susceptibility)
- Statistical analysis of the weight-change distribution

With 100 cycles and even a modest single-cycle SNR of 10:1, the accumulated significance exceeds 100σ.

6.5 Cost and Feasibility

Item	Estimated Cost
10 kg niobium sample	\$5,000
Cryostat (closed-cycle, 4 K)	\$25,000
Force sensor / load cell	\$5,000
Magnetic shielding	\$3,000
Temperature control & monitoring	\$5,000
Data acquisition	\$3,000
Total	**\$46,000**

Table 5: Estimated cost for the Cooper pair mass anomaly weight test. Many university cryogenics labs already possess the cryostat and DAQ, reducing the marginal cost to ∼\$15,000.

This experiment can be performed in **any university cryogenics laboratory** with a 4 K cryostat. No particle accelerator, no space mission, no high-Q cavity—just a chunk of niobium and a good scale.

7 Why This Mechanism Is Fundamentally Different

7.1 The Problem with Previous Propulsion Proposals

All previous attempts at "gravitational propulsion" from electromagnetic energy encounter a fundamental barrier. The coupling between electromagnetic energy and spacetime curvature (or in DFD, the ψ-field) goes through the gravitational coupling G/c^4:

$$\frac{2G}{c^4} = 1.65 \times 10^{-44} \text{ m/J}. \tag{28}$$

To produce 1 N of force by *creating* a ψ-gradient from EM energy requires storing $\sim 10^{44}$ J—the rest-mass energy of Jupiter.

Previous proposals include:

- **Psi-bubble propulsion:** Creates ψ-gradient using high-Q SRF cavities and rotating EM fields. Achievable forces: $\sim 10^{-35}$ N for laboratory-scale systems.
- **Podkletnov rotating superconductor:** Claimed weight reduction above a rotating YBCO disk. Never independently replicated; likely electromagnetic artifacts.
- **Tajmar gravitomagnetic London moment:** Reported anomalous accelerations near rotating superconductors. Later shown by Tajmar's own group to be electromagnetic artifacts at nano-Newton precision.

7.2 The Cooper Pair Mechanism Is Different

The Cooper pair mass anomaly mechanism does **not** attempt to create a ψ-gradient. Instead, it exploits a pre-existing feature of the DFD matter–ψ coupling:

Fundamental Distinction

Previous approaches: Supply EM energy $\rightarrow$ create ψ-gradient $\rightarrow$ produce force.
$\Rightarrow$ Blocked by the $2G/c^4$ wall.
Cooper pair mechanism: Modify the *coupling* between existing matter and the *existing* ψ-field (Earth's gravitational field) by switching the quantum state of the condensate.
$\Rightarrow$ No energy barrier. The anomalous mass $\delta = 92$ ppm is *already there* (Tate measured it in 1989).

The physics is analogous to the difference between:

- Creating an electric field from scratch (requires charge separation, costs energy)
- Changing the dielectric constant of a material in an existing field (phase transition, comparatively trivial)

The superconducting transition is the "dielectric switch" for gravity. It costs only the condensation energy ($\sim$ meV per Cooper pair), but it modifies how ~ 92 ppm of the condensate's mass-energy couples to ψ.

Mechanism	Force (10 kg)	Cost	Complexity
ψ-bubble (SRF cavity)	$\sim 10^{-35}$ N	\$10M+	Extreme
Podkletnov (rotating SC)	unconfirmed	\$1M+	High
Tajmar (gravitomagnetic)	null result	\$500K+	High
Cooper pair anomaly	**3.44 mN**	**\$50K**	**Low**

Table 6: Comparison of propulsion mechanisms. The Cooper pair anomaly produces a force $\sim 10^{32}$ times larger than the ψ-bubble mechanism, at 1/200th the cost, with a clear experimental signature. The only caveat is the unknown value of λ_{quintet}.

7.3 Comparison Table

8 Theoretical Question: What Determines λ_{quintet}?

8.1 The Central Question

The entire propulsion viability hinges on one theoretical question:

The λ_{quintet} Question

Does the quintet exchange energy in $S^2(V^*)$ contribute to the stress-energy tensor $T_{\mu\nu}$ that sources the ψ-field?

- $\lambda_{\text{quintet}} = 1$: Yes $\to$ EP preserved, no propulsion
- $\lambda_{\text{quintet}} = 0$: No $\to$ EP violated for Cooper pairs, propulsion viable
- $0 < \lambda_{\text{quintet}} < 1$: Partial $\to$ reduced but nonzero force

8.2 DFD Action Principle Considerations

In the DFD framework, matter couples to ψ through the frame stiffness action:

$$\mathcal{L}_{\text{matter}} = -\kappa_0 \, \psi \cdot S[\rho], \tag{29}$$

where ρ is the internal density matrix and $S[\rho]$ is the von Neumann entropy of the microsector. The partition function $Z(S^3)$ encodes the sum over all microsector states.

The quintet channel in $S^2(V^*)$ arises from the *pairing* of two electrons within the V^* representation. Its contribution to $S[\rho]$ depends on whether the bridge map

$$\mathcal{B} : \mathcal{H}_{\text{ch}} \to \mathcal{H}_{\text{macro}} \tag{30}$$

preserves or projects out the quintet component when mapping microsector states to macroscopic observables that source ψ.

8.3 Three Scenarios

Scenario A: Universal coupling ($\lambda_{\text{quintet}} = 1$). All internal energy gravitates. The bridge map is the identity on the energy eigenspace. The EP is exactly preserved for

Cooper pairs. The Tate anomaly exists in the inertial mass but is exactly matched in the gravitational mass. No propulsion.

Scenario B: Projected coupling ($\lambda_{\text{quintet}} = 0$). The bridge map projects onto the singlet channel $|\mathbf{1}\rangle \in S^2(V^*)$, discarding the quintet $|\mathbf{5}\rangle$. Physically, this means the quintet exchange energy is "internal" to the microsector and does not appear in the macroscopic stress-energy. The EP is violated for Cooper pairs by exactly $\delta = 92$ ppm. Full propulsion force.

This scenario has a natural interpretation: the quintet channel represents an *internal* degree of freedom of the A_5 microsector that does not couple to the macroscopic ψ-field, analogous to how the colour degree of freedom of quarks does not produce macroscopic colour fields due to confinement.

Scenario C: Partial coupling ($0 < \lambda_{\text{quintet}} < 1$). The bridge map partially transmits the quintet energy. This could arise from finite-temperature or finite-coherence-length corrections to the bridge map. Produces a reduced but nonzero force proportional to $(1 - \lambda_{\text{quintet}})$.

8.4 What Theory Predicts (Open Question)

Determining λ_{quintet} from first principles requires computing the bridge map $\mathcal{B}$ for the specific case of the quintet representation. This involves:

1. The structure of $S^2(V^*)$ as a representation of A_5
2. The action of the Cayley graph propagator on the quintet subspace
3. Whether the ψ-field sources from the full microsector partition function or only from its projection onto specific representation channels
4. The role of gauge invariance: if the quintet transforms nontrivially under the residual gauge group, its expectation value may vanish in the macroscopic limit (analogous to colour confinement)

This calculation is the **most important open theoretical problem in the DFD propulsion programme**. However, it is also a question that can be settled *experimentally* far more quickly and cheaply than theoretically, using the weight-change test described in Section **??**.

9 Implications and Next Steps

9.1 If $\lambda_{\text{quintet}} \neq 1$ (Propulsion Works)

The implications are transformative:

1. **Immediate:** The equivalence principle is violated for superconducting condensates. This is the first macroscopic EP violation and constitutes a discovery of the highest magnitude in fundamental physics.

2. **Engineering:** A passive gravitational shielding effect of $\delta \cdot f_{\mathrm{CP}} = 35$ ppm is available by cooling niobium below T_c. No energy input beyond cryogenic cooling is needed.

3. **Propulsion:** For a 1000 kg Nb payload at 4 K:

$$F = 1000 \times 9.81 \times 0.38 \times 9.22 \times 10^{-5} = 0.344 \text{ N}. \tag{31}$$

   This is not enough for launch but is significant for station-keeping, attitude control, and long-duration deep-space missions.

4. **Scaling:** Higher-T_c superconductors with larger condensate fractions could enhance the effect. If the anomaly scales with T_c (via increased pairing interaction), materials like MgB_2 ($T_c = 39$ K) or cuprates ($T_c \sim 90$ K) may show larger effects.

5. **Amplification:** If the anomalous mass can be modulated (e.g., by driving the condensate through T_c at high frequency), the resulting oscillating gravitational mass could be used for gravitational wave generation or resonant coupling to external masses.

9.2 If $\lambda_{\mathrm{quintet}} = 1$ (No Propulsion)

Even a null result is highly valuable:

1. It confirms the EP for superconducting condensates at the 0.01 mg level—a new precision frontier.

2. Combined with the DFD prediction of δ, it constrains the bridge map to be representation-preserving, which is important theoretical information.

3. It motivates the search for other channels where $\lambda \neq 1$ might hold.

9.3 Experimental Roadmap

1. **Phase 1 (6 months, \$50K):** Niobium weight-change test at a university cryogenics lab. Goal: detect or bound ΔW at the 0.1 mg level.

2. **Phase 2 (12 months, \$200K):** If signal detected, replicate with:

   - Multiple Nb samples of different masses (1, 5, 10, 50 kg)
   - Verify $\Delta W \propto M$ scaling
   - Test temperature dependence: $\Delta W(T)$ should follow the BCS gap function $\Delta(T)$ below T_c

3. **Phase 3 (18 months, \$500K):** Extend to other superconductors (Pb, Sn, Al, MgB_2, YBCO) to test material dependence.

4. **Phase 4 (24 months):** If the effect is confirmed across materials, begin engineering optimization for maximum force-to-mass ratio.

10 Conclusions

1. The Tate Cooper pair mass anomaly (92±21 ppm above BCS) has been unexplained for 36 years. DFD predicts *exactly* $\delta = \sqrt{3}\,\alpha^2 = 92.23$ ppm from the quintet channel of the A_5 microsector.

2. The Ross et al. (2024) Eöt-Wash bound $\eta_{\rm CP} \leq 9.2 \times 10^{-4}$ is ten times too weak to constrain the gravitational coupling parameter $\lambda_{\rm quintet}$. The propulsion-relevant value $\lambda_{\rm quintet} = 0$ is completely allowed.

3. For a 10 kg niobium sample with $\lambda_{\rm quintet} = 0$, the predicted weight anomaly is 3.44 mN (350 mg_f), detectable at 35,000:1 SNR on a 0.01 mg balance.

4. All systematic errors (Meissner effect, buoyancy, thermal contraction, EM artifacts) are controllable to levels far below the predicted signal.

5. The experiment can be performed in any university cryogenics laboratory for less than $50,000. This is the cheapest and simplest test of any proposed propulsion mechanism by many orders of magnitude.

6. Unlike all previous propulsion proposals, this mechanism does not create ψ-gradients from EM energy (and thus is not blocked by the $2G/c^4$ wall). It modifies the *coupling* between existing matter and the existing gravitational field via a quantum phase transition.

7. The theoretical question of $\lambda_{\rm quintet}$ can be settled experimentally far more quickly than theoretically. **This experiment should be the highest priority in the DFD programme.**

References

[1] J. Tate, B. Cabrera, S.B. Felch, and J.T. Anderson, "Precise determination of the Cooper-pair mass," *Phys. Rev. Lett.* **62**, 845–848 (1989).

[2] J. Tate, B. Cabrera, S.B. Felch, and J.T. Anderson, "Determination of the Cooper-pair mass in niobium," *Phys. Rev. B* **42**, 7885–7893 (1990).

[3] C. Hoang, M. Bajraktari, and M. Tajmar, "High-precision measurement of Cooper-pair mass using rotating spherical-shell superconductor," *Mater. Lett.* **262**, 127178 (2020).

[4] M.P. Ross, S.M. Fleischer, I.A. Paulson, P. Lamb, B.M. Iritani, E.G. Adelberger, C.A. Hagedorn, K. Venkateswara, C. Gettings, E.A. Shaw, S.K. Apple, and J.H. Gundlach, "Test of the equivalence principle for superconductors," arXiv:2407.21232 (2024).

[5] M. Tajmar and C.J. de Matos, "Gravitomagnetic fields in rotating superconductors to solve Tate's Cooper pair mass anomaly," *AIP Conf. Proc.* **813**, 1415 (2006); arXiv:gr-qc/0607086.

[6] M. Tajmar, O. Neunzig, and M. Kößling, "Measurement of anomalous forces from a Cooper-pair current in high-T_c superconductors with nano-Newton precision," *Front. Phys.* **10**, 892215 (2022).

[7] B.S. DeWitt, "Superconductors and gravitational drag," *Phys. Rev. Lett.* **16**, 1092 (1966).

[8] P. Touboul *et al.* (MICROSCOPE Collaboration), "MICROSCOPE mission: Final results of the test of the equivalence principle," *Phys. Rev. Lett.* **129**, 121102 (2022).

[9] G.T. Alcock, *Density Field Dynamics: A Complete Unified Theory*, v3.2 (2026).

A Detailed Numerical Calculations

A.1 Physical Constants Used

$$\alpha = 1/137.035\,999\,084 = 7.2974 \times 10^{-3} \tag{32}$$
$$m_e = 9.1094 \times 10^{-31}\ \text{kg} \tag{33}$$
$$g = 9.8067\ \text{m/s}^2 \tag{34}$$
$$\rho_{\text{Nb}} = 8{,}570\ \text{kg/m}^3 \tag{35}$$
$$T_{c,\text{Nb}} = 9.25\ \text{K} \tag{36}$$

A.2 DFD Anomaly Derivation Summary

$$\alpha^2 = 5.3251 \times 10^{-5} \tag{37}$$
$$\sqrt{3} = 1.7321 \tag{38}$$
$$\delta = \sqrt{3}\,\alpha^2 = 9.2234 \times 10^{-5} = 92.23\ \text{ppm} \tag{39}$$

A.3 Ross Bound Analysis

$$\eta_{\text{CP}} \leq 9.2 \times 10^{-4} \tag{40}$$
$$\delta_{\text{DFD}} = 9.22 \times 10^{-5} \tag{41}$$
$$\frac{\eta_{\text{CP}}}{\delta_{\text{DFD}}} = \frac{9.2 \times 10^{-4}}{9.22 \times 10^{-5}} = 9.98 \approx 10.0 \tag{42}$$
$$|1 - \lambda_{\text{quintet}}| \leq 10.0 \tag{43}$$

A.4 Force Calculation Verification

For $M = 10$ kg, $\lambda_{\text{quintet}} = 0$:

$$\begin{aligned}
\Delta W &= M \cdot g \cdot f_{\text{CP}} \cdot \delta \cdot |1 - \lambda| && (44)\\
&= 10 \times 9.807 \times 0.38 \times 9.223 \times 10^{-5} \times 1.0 && (45)\\
&= 10 \times 9.807 \times 3.505 \times 10^{-5} && (46)\\
&= 10 \times 3.437 \times 10^{-4} && (47)\\
&= 3.437 \times 10^{-3}\ \text{N} = 3.437\ \text{mN} && (48)
\end{aligned}$$

Weight equivalent:

$$\Delta m_f = \frac{\Delta W}{g} = \frac{3.437 \times 10^{-3}}{9.807} = 3.505 \times 10^{-4} \text{ kg} = 350.5 \text{ mg} \tag{49}$$