

A Family of ODE-solver based Fast ADMM with Applications to Sparse Regression

Hao Luo (罗浩)

National Center for Applied Mathematics in Chongqing
Chongqing Normal University
Email: luohao@cqnu.edu.cn



Workshop on Scientific Computing and Data Science
Department of Mathematics Chinese University of Hong Kong
July 11–12, 2024

A joint work with Dr. Zi-Hang Zhang from PKU

Outline

Introduction

PDHG and CP

Primal-Dual Flow Dynamics

Accelerated Primal-Dual Flow

Time Discretization

Numerical Results

Introduction

PDHG and CP

Primal-Dual Flow Dynamics

Accelerated Primal-Dual Flow

Time Discretization

Numerical Results

Problem setting

► Equality constrained separable minimization

$$\min_{u \in \mathcal{U}, v \in \mathcal{V}} F(u, v) := f(u) + g(v) \quad \text{s.t. } Au + Bv = b \quad (1)$$

Assumptions:

- $\mathcal{U}, \mathcal{V}, \Lambda$: Hilbert spaces with inner product $\langle \cdot, \cdot \rangle$ ¹
- $A(B) : \mathcal{U}(\mathcal{V}) \rightarrow \Lambda$: bounded linear operators, $b \in \Lambda$
- $f(g) : \mathcal{U}(\mathcal{V}) \rightarrow (-\infty, +\infty]$: CCP² with constants $\mu_f(\mu_g) \geq 0$
- Consistent condition: $b \in A \text{ dom } f + B \text{ dom } g$

► Composite convex minimization ($b = 0, B = -I$)

$$\min_{u \in \mathcal{U}} P(u) := f(u) + g(Au) \quad (2)$$

¹When no confusion arises, we use the same bracket $\langle \cdot, \cdot \rangle$ for the inner products on \mathcal{U}, \mathcal{V} and Λ .

²CCP means closed, convex and proper.

Preliminary

- ▶ Introduce $\mathbf{x} = (u, v)$, $\mathbf{A} = (A, B)$ and restate the single block form

$$\min_{\mathbf{x} \in \mathcal{U} \times \mathcal{V}} F(\mathbf{x}) \quad \text{s.t. } \mathbf{A}\mathbf{x} = b \quad (3)$$

- ▶ Define the Lagrangian

$$\mathcal{L}(\mathbf{x}, \lambda) := F(\mathbf{x}) + \langle \lambda, \mathbf{A}\mathbf{x} - b \rangle, \quad (\mathbf{x}, \lambda) \in \mathcal{X} := \text{dom } F \times \Lambda^*.$$

- ▶ Saddle-point $(\hat{\mathbf{x}}, \hat{\lambda}) \in \mathcal{X}$:

$$\mathcal{L}(\hat{\mathbf{x}}, \lambda) \leq \mathcal{L}(\hat{\mathbf{x}}, \hat{\lambda}) \leq \mathcal{L}(\mathbf{x}, \hat{\lambda}) \quad \forall (\mathbf{x}, \lambda) \in \mathcal{X}$$

- ▶ Monotone inclusion

$$0 \in M(\hat{\mathbf{x}}, \hat{\lambda}), \quad M(\mathbf{x}, \lambda) = \begin{pmatrix} \partial F(\mathbf{x}) + \mathbf{A}^\top \lambda \\ b - \mathbf{A}\mathbf{x} \end{pmatrix}$$

Applications

Many variational/optimization problems are related to (1)/(2)/(3):

- ▶ Image processing
 - Image denoising: TV-based model, ROF
 - Image deconvolution
 - ...
- ▶ Dynamical optimal transport/Benamou–Brenier problem
- ▶ Sparse regression: Lasso, least absolute deviation (LAD)
- ▶ ...

Existing (Lagrangian-based) methods

- Augmented Lagrangian method (ALM) for (3):

$$\mathbf{x}_{k+1} = \underset{\mathbf{x}}{\operatorname{argmin}} \mathcal{L}_\sigma(\mathbf{x}, \lambda_k), \quad \lambda_{k+1} = \lambda_k + \sigma(\mathbf{A}\mathbf{x}_{k+1} - b)$$

with $\mathcal{L}_\sigma(\mathbf{x}, \lambda) := \mathcal{L}(\mathbf{x}, \lambda) + \sigma/2 \|\mathbf{A}\mathbf{x} - b\|^2$, $\sigma > 0$.

- Hestenes (1969) and Powell (1969)
- Dual formulation = Proximal point algorithm (Rockafellar)
- Uzawa method
- **Not easy** to update $\mathbf{x}_{k+1} = (u_{k+1}, v_{k+1})$

Existing (Lagrangian-based) methods

- ▶ Augmented Lagrangian method (ALM) for (3):

$$\mathbf{x}_{k+1} = \underset{\mathbf{x}}{\operatorname{argmin}} \mathcal{L}_\sigma(\mathbf{x}, \lambda_k), \quad \lambda_{k+1} = \lambda_k + \sigma(\mathbf{A}\mathbf{x}_{k+1} - b)$$

with $\mathcal{L}_\sigma(\mathbf{x}, \lambda) := \mathcal{L}(\mathbf{x}, \lambda) + \sigma/2 \|\mathbf{A}\mathbf{x} - b\|^2$, $\sigma > 0$.

- Hestenes (1969) and Powell (1969)
- Dual formulation = Proximal point algorithm (Rockafellar)
- Uzawa method
- **Not easy** to update $\mathbf{x}_{k+1} = (u_{k+1}, v_{k+1})$

- ▶ Alternating direction method of multipliers (ADMM):

$$\left. \begin{aligned} u_{k+1} &= \underset{u}{\operatorname{argmin}} \mathcal{L}_\sigma(u, v_k, \lambda_k) \\ v_{k+1} &= \underset{v}{\operatorname{argmin}} \mathcal{L}_\sigma(u_{k+1}, v, \lambda_k) \end{aligned} \right\} \text{Decouple } u \text{ and } v$$
$$\lambda_{k+1} = \lambda_k + \sigma(Au_{k+1} + Bv_{k+1} - b)$$

- Numerical solution of PDEs from mechanics, physics and differential geometry (Glowinski, 1976/2014)

- Popular application to image processing, statistical learning, data mining (Boyd et al., *Found Trends Mach Learn*, 2010)
- **Successive computation**, similar with Alternating Direction Method (ADM) and Gauss-Seidel iteration
- Dual formulation = DRSM
- Convergence rate:

$$O(1/\sqrt{k}), \quad O(1/k)(\text{Ergodic}), \quad O(1/k^2)(\mu_f + \mu_g > 0, \text{Ergodic})$$

- Popular application to image processing, statistical learning, data mining (Boyd et al., *Found Trends Mach Learn*, 2010)
- **Successive computation**, similar with Alternating Direction Method (ADM) and Gauss-Seidel iteration
- Dual formulation = DRSM
- Convergence rate:

$$O(1/\sqrt{k}), \quad O(1/k)(\text{Ergodic}), \quad O(1/k^2)(\mu_f + \mu_g > 0, \text{Ergodic})$$

► “**Ergodic**” means average of historical iterates:

$$\bar{\mathbf{x}}_k = \sum_{i=1}^k \alpha_i \mathbf{x}_i, \quad \sum_{i=1}^k \alpha_i = 1$$

- Popular application to image processing, statistical learning, data mining (Boyd et al., *Found Trends Mach Learn*, 2010)
- **Successive computation**, similar with Alternating Direction Method (ADM) and Gauss-Seidel iteration
- Dual formulation = DRSM
- Convergence rate:

$$O(1/\sqrt{k}), \quad O(1/k)(\text{Ergodic}), \quad O(1/k^2)(\mu_f + \mu_g > 0, \text{Ergodic})$$

- “**Ergodic**” means average of historical iterates:

$$\bar{\mathbf{x}}_k = \sum_{i=1}^k \alpha_i \mathbf{x}_i, \quad \sum_{i=1}^k \alpha_i = 1$$

- Primal-dual solvers for (2):
- Arrow–Huricz, PDHG (Zhu and Chan, 2008)
 - Chambolle–Pock’s method (*JMIV*, 2011)
 - Tight connection with ADMM
 - **Ergodic convergence rate** $O(1/k)$

Acceleration and nonergodic rate

► For unconstrained problem: $\min f(x)$

$$\begin{cases} O(1/k) & \mu_f = 0 \\ O(e^{-k/\text{cond}(f)}) & \mu_f > 0 \end{cases} \implies \begin{cases} O(1/k^2) & \mu_f = 0 \\ O(e^{-k/\sqrt{\text{cond}(f)}}) & \mu_f > 0 \end{cases}$$

Acceleration and nonergodic rate

- For unconstrained problem: $\min f(x)$

$$\begin{cases} O(1/k) & \mu_f = 0 \\ O(e^{-k/\text{cond}(f)}) & \mu_f > 0 \end{cases} \implies \begin{cases} O(1/k^2) & \mu_f = 0 \\ O(e^{-k/\sqrt{\text{cond}(f)}}) & \mu_f > 0 \end{cases}$$

- **Why nonergodic?**

Ensure **sparsity** or **low-rankness**; see Li and Lin (JSC, 2019), Tran-Dinh and Zhu (SIOPT, 2020)

Acceleration and nonergodic rate

- For unconstrained problem: $\min f(x)$

$$\begin{cases} O(1/k) & \mu_f = 0 \\ O(e^{-k/\text{cond}(f)}) & \mu_f > 0 \end{cases} \implies \begin{cases} O(1/k^2) & \mu_f = 0 \\ O(e^{-k/\sqrt{\text{cond}(f)}}) & \mu_f > 0 \end{cases}$$

- **Why nonergodic?**

Ensure **sparsity** or **low-rankness**; see Li and Lin (JSC, 2019), Tran-Dinh and Zhu (SIOPT, 2020)

- For problem (1), accelerated ADMM with nonergodic rate:
 - For $\mu_f = \mu_g = 0$
 - Li and Lin (JSC, 2017): $O(1/k)$
 - Ouyang et al. (SIIS, 2015): $O\left(\frac{L}{k^2} + \frac{\|A\|}{k}\right)$ (**mixed-type**)

Acceleration and nonergodic rate

- For unconstrained problem: $\min f(x)$

$$\begin{cases} O(1/k) & \mu_f = 0 \\ O(e^{-k/\text{cond}(f)}) & \mu_f > 0 \end{cases} \implies \begin{cases} O(1/k^2) & \mu_f = 0 \\ O(e^{-k/\sqrt{\text{cond}(f)}}) & \mu_f > 0 \end{cases}$$

- **Why nonergodic?**

Ensure **sparsity** or **low-rankness**; see Li and Lin (JSC, 2019), Tran-Dinh and Zhu (SIOPT, 2020)

- For problem (1), accelerated ADMM with nonergodic rate:
 - For $\mu_f = \mu_g = 0$
 - Li and Lin (JSC, 2017): $O(1/k)$
 - Ouyang et al. (SIIS, 2015): $O\left(\frac{L}{k^2} + \frac{\|A\|}{k}\right)$ (**mixed-type**)
 - For $\mu_f + \mu_g > 0$: $O(1/k^2)$
 - Tran-Dinh et al. (SIOPT, 2020): semi-ergodic rate
 - **Sabach and Teboulle (SIOPT, 2022)**
 - **Zhang et al. (arXiv:2206.05088, 2022)**
 - **He et al. (arXiv:2310.16404, 2023)**

Main contribution

- ▶ A continuous ODE-based framework:
 - A family of accelerated ADMM from a **systematic** way
 - A unified Lyapunov analysis approach
 - Sharp **mixed-type estimate** and **nonergodic** rates
 - Both convex ($\mu_f = \mu_g = 0$) and (partially) strongly convex ($\mu_f + \mu_g > 0$)

Main contribution

- ▶ A continuous ODE-based framework:
 - A family of accelerated ADMM from a **systematic** way
 - A unified Lyapunov analysis approach
 - Sharp **mixed-type estimate** and **nonergodic** rates
 - Both convex ($\mu_f = \mu_g = 0$) and (partially) strongly convex ($\mu_f + \mu_g > 0$)
- ▶ Applications to sparse regression: fast nonergodic convergence and sparsity maintaining

Introduction

PDHG and CP

Primal-Dual Flow Dynamics

Accelerated Primal-Dual Flow

Time Discretization

Numerical Results

PDHG and CP

► PDHG/Arrow–Hurwicz algorithm ³

$$\mathbf{x}_{k+1} = \operatorname{argmin}_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda_k) + \frac{1}{2\tau} \|\mathbf{x} - \mathbf{x}_k\|^2$$

$$\lambda_{k+1} = \operatorname{argmax}_{\lambda} \mathcal{L}(\mathbf{x}_{k+1}, \lambda) - \frac{1}{2\tau} \|\lambda - \lambda_k\|^2$$

³M. Zhu and T. F. Chan. An efficient primal-dual hybrid gradient algorithm for total variation image restoration. CAM Report 08-34, 2008.

PDHG and CP

► PDHG/Arrow–Hurwicz algorithm ³

$$\mathbf{x}_{k+1} = \operatorname{argmin}_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda_k) + \frac{1}{2\tau} \|\mathbf{x} - \mathbf{x}_k\|^2$$

$$\lambda_{k+1} = \operatorname{argmax}_{\lambda} \mathcal{L}(\mathbf{x}_{k+1}, \lambda) - \frac{1}{2\tau} \|\lambda - \lambda_k\|^2$$

- $0 \in S[\mathbf{z}_{k+1} - \mathbf{z}_k] + \tau M(\mathbf{z}_{k+1})$, $S = \begin{pmatrix} I & -\tau \mathbf{A}^\top \\ O & I \end{pmatrix}$
- Preconditioned PPA without symmetric property ⁴

³M. Zhu and T. F. Chan. An efficient primal-dual hybrid gradient algorithm for total variation image restoration. CAM Report 08-34, 2008.

⁴B. He and X. Yuan. Convergence analysis of primal-dual algorithms for a saddle-point problem: from contraction perspective. SIIS, 5(1):119–149, 2012.

PDHG and CP

► PDHG/Arrow–Hurwicz algorithm ³

$$\mathbf{x}_{k+1} = \operatorname{argmin}_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda_k) + \frac{1}{2\tau} \|\mathbf{x} - \mathbf{x}_k\|^2$$

$$\lambda_{k+1} = \operatorname{argmax}_{\lambda} \mathcal{L}(\mathbf{x}_{k+1}, \lambda) - \frac{1}{2\tau} \|\lambda - \lambda_k\|^2$$

- $0 \in S[\mathbf{z}_{k+1} - \mathbf{z}_k] + \tau M(\mathbf{z}_{k+1})$, $S = \begin{pmatrix} I & -\tau \mathbf{A}^\top \\ O & I \end{pmatrix}$
- Preconditioned PPA without symmetric property ⁴
- Asymmetric \Rightarrow Diverge! Counter examples in He et al. *JIMV* (2022)

³M. Zhu and T. F. Chan. An efficient primal-dual hybrid gradient algorithm for total variation image restoration. CAM Report 08-34, 2008.

⁴B. He and X. Yuan. Convergence analysis of primal-dual algorithms for a saddle-point problem: from contraction perspective. *SIIS*, 5(1):119–149, 2012.

PDHG and CP

► PDHG/Arrow–Hurwicz algorithm ³

$$\mathbf{x}_{k+1} = \operatorname{argmin}_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda_k) + \frac{1}{2\tau} \|\mathbf{x} - \mathbf{x}_k\|^2$$

$$\lambda_{k+1} = \operatorname{argmax}_{\lambda} \mathcal{L}(\mathbf{x}_{k+1}, \lambda) - \frac{1}{2\tau} \|\lambda - \lambda_k\|^2$$

- $0 \in S[\mathbf{z}_{k+1} - \mathbf{z}_k] + \tau M(\mathbf{z}_{k+1})$, $S = \begin{pmatrix} I & -\tau \mathbf{A}^\top \\ O & I \end{pmatrix}$
- Preconditioned PPA without symmetric property ⁴
- **Asymmetric** \Rightarrow **Diverge!** Counter examples in He et al. *JIMV* (2022)

► Chambolle-Pock (CP) method ⁵

$$\mathbf{x}_{k+1} = \operatorname{argmin}_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda_k) + \frac{1}{2\tau} \|\mathbf{x} - \mathbf{x}_k\|^2$$

$$\lambda_{k+1} = \operatorname{argmax}_{\lambda} \mathcal{L}(\mathbf{x}_{k+1} + \mathbf{x}_{k+1} - \mathbf{x}_k, \lambda) - \frac{1}{2\tau} \|\lambda - \lambda_k\|^2$$

³ M. Zhu and T. F. Chan. An efficient primal-dual hybrid gradient algorithm for total variation image restoration. CAM Report 08-34, 2008.

⁴ B. He and X. Yuan. Convergence analysis of primal-dual algorithms for a saddle-point problem: from contraction perspective. *SIIS*, 5(1):119–149, 2012.

PDHG and CP

► PDHG/Arrow–Hurwicz algorithm ³

$$\mathbf{x}_{k+1} = \operatorname{argmin}_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda_k) + \frac{1}{2\tau} \|\mathbf{x} - \mathbf{x}_k\|^2$$

$$\lambda_{k+1} = \operatorname{argmax}_{\lambda} \mathcal{L}(\mathbf{x}_{k+1}, \lambda) - \frac{1}{2\tau} \|\lambda - \lambda_k\|^2$$

- $0 \in S[\mathbf{z}_{k+1} - \mathbf{z}_k] + \tau M(\mathbf{z}_{k+1})$, $S = \begin{pmatrix} I & -\tau \mathbf{A}^\top \\ O & I \end{pmatrix}$
- Preconditioned PPA without symmetric property ⁴
- **Asymmetric** \Rightarrow **Diverge!** Counter examples in He et al. *JIMV* (2022)

► Chambolle-Pock (CP) method ⁵

$$\mathbf{x}_{k+1} = \operatorname{argmin}_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda_k) + \frac{1}{2\tau} \|\mathbf{x} - \mathbf{x}_k\|^2$$

$$\lambda_{k+1} = \operatorname{argmax}_{\lambda} \mathcal{L}(\mathbf{x}_{k+1} + \mathbf{x}_{k+1} - \mathbf{x}_k, \lambda) - \frac{1}{2\tau} \|\lambda - \lambda_k\|^2$$

- $0 \in \mathbf{S}[\mathbf{z}_{k+1} - \mathbf{z}_k] + \tau M(\mathbf{z}_{k+1})$, $\mathbf{S} = \begin{pmatrix} I & -\tau \mathbf{A}^\top \\ -\tau \mathbf{A} & I \end{pmatrix}$

³ M. Zhu and T. F. Chan. An efficient primal-dual hybrid gradient algorithm for total variation image restoration. CAM Report 08-34, 2008.

⁴ B. He and X. Yuan. Convergence analysis of primal-dual algorithms for a saddle-point problem: from contraction perspective. *SIIS*, 5(1):119–149, 2012.

Introduction

PDHG and CP

Primal-Dual Flow Dynamics

Accelerated Primal-Dual Flow

Time Discretization

Numerical Results

Primal-dual flow

- Rewriting CP method as a semi-implicit discretization

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\tau} &\in -\partial_{\mathbf{x}}\mathcal{L}(\mathbf{x}_{k+1}, \lambda_k) \\ \frac{\lambda_{k+1} - \lambda_k}{\tau} &= \nabla_{\lambda}\mathcal{L}(\mathbf{x}_{k+1} + \mathbf{x}_{k+1} - \mathbf{x}_k, \lambda_{k+1})\end{aligned}$$

Primal-dual flow

- ▶ Rewriting CP method as a semi-implicit discretization

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\tau} &\in -\partial_{\mathbf{x}}\mathcal{L}(\mathbf{x}_{k+1}, \lambda_k) \\ \frac{\lambda_{k+1} - \lambda_k}{\tau} &= \nabla_{\lambda}\mathcal{L}(\mathbf{x}_{k+1} + \mathbf{x}_{k+1} - \mathbf{x}_k, \lambda_{k+1})\end{aligned}$$

- ▶ We proposed a novel primal-dual flow ⁶

$$\begin{aligned}\gamma \mathbf{x}' &\in -\partial_{\mathbf{x}}\mathcal{L}(\mathbf{x}, \lambda) \\ \theta \lambda' &= \nabla_{\lambda}\mathcal{L}(\mathbf{x} + \mathbf{x}', \lambda)\end{aligned}$$

(PD Flow)

⁶L. A primal-dual flow for affine constrained convex optimization. *ESAIM:COCV* (2022)

Primal-dual flow

- ▶ Rewriting CP method as a semi-implicit discretization

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\tau} &\in -\partial_{\mathbf{x}}\mathcal{L}(\mathbf{x}_{k+1}, \lambda_k) \\ \frac{\lambda_{k+1} - \lambda_k}{\tau} &= \nabla_{\lambda}\mathcal{L}(\mathbf{x}_{k+1} + \mathbf{x}_{k+1} - \mathbf{x}_k, \lambda_{k+1})\end{aligned}$$

- ▶ We proposed a novel primal-dual flow ⁶

$$\boxed{\begin{aligned}\gamma \mathbf{x}' &\in -\partial_{\mathbf{x}}\mathcal{L}(\mathbf{x}, \lambda) \\ \theta \lambda' &= \nabla_{\lambda}\mathcal{L}(\mathbf{x} + \mathbf{x}', \lambda)\end{aligned}} \quad \text{(PD Flow)}$$

- ▶ Scaling parameters $\theta' = -\theta$ and $\gamma' = \mu - \gamma$ with $\mu = \mu_F$

⁶L. A primal-dual flow for affine constrained convex optimization. *ESAIM:COCV* (2022)

Primal-dual flow

- ▶ Rewriting CP method as a semi-implicit discretization

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\tau} &\in -\partial_{\mathbf{x}}\mathcal{L}(\mathbf{x}_{k+1}, \lambda_k) \\ \frac{\lambda_{k+1} - \lambda_k}{\tau} &= \nabla_{\lambda}\mathcal{L}(\mathbf{x}_{k+1} + \mathbf{x}_{k+1} - \mathbf{x}_k, \lambda_{k+1})\end{aligned}$$

- ▶ We proposed a novel primal-dual flow ⁶

$$\boxed{\begin{aligned}\gamma \mathbf{x}' &\in -\partial_{\mathbf{x}}\mathcal{L}(\mathbf{x}, \lambda) \\ \theta \lambda' &= \nabla_{\lambda}\mathcal{L}(\mathbf{x} + \mathbf{x}', \lambda)\end{aligned}} \quad \text{(PD Flow)}$$

- ▶ Scaling parameters $\theta' = -\theta$ and $\gamma' = \mu - \gamma$ with $\mu = \mu_F$
- ▶ Exponential decay of the Lyapunov function

$$\mathcal{E}(\mathbf{x}, \lambda) := \mathcal{L}(\mathbf{x}, \hat{\lambda}) - \mathcal{L}(\hat{\mathbf{x}}, \lambda) + \frac{\gamma}{2} \|\mathbf{x} - \hat{\mathbf{x}}\|^2 + \frac{\theta}{2} \|\lambda - \hat{\lambda}\|^2$$

⁶L. A primal-dual flow for affine constrained convex optimization. *ESAIM:COCV* (2022)

Introduction

PDHG and CP

Primal-Dual Flow Dynamics

Accelerated Primal-Dual Flow

Time Discretization

Numerical Results

Accelerated primal-dual flow

- ▶ **Main idea:** Combining primal-dual form with **acceleration**

Accelerated primal-dual flow

- ▶ **Main idea:** Combining primal-dual form with **acceleration**
- ▶ Nesterov accelerated gradient flow for $\min F(\mathbf{x})$ ⁷

$$\gamma \mathbf{x}'' + (\mu + \gamma) \mathbf{x}' + \partial F(\mathbf{x}) \ni 0 \quad (\text{NAG flow})$$

⁷L. and L. Chen. From differential equation solvers to accelerated first-order methods for convex optimization. MAPR, 195: 735–781, 2022.

Accelerated primal-dual flow

- ▶ **Main idea:** Combining primal-dual form with **acceleration**
- ▶ Nesterov accelerated gradient flow for $\min F(\mathbf{x})$ ⁷

$$\gamma \mathbf{x}'' + (\mu + \gamma) \mathbf{x}' + \partial F(\mathbf{x}) \ni 0 \quad (\text{NAG flow})$$

- ▶ Combining NAG-flow with **(PD Flow)** gives

$$\gamma \mathbf{x}' \in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda) \quad [\text{Primal descent}]$$

$$\theta \lambda' = \nabla_{\lambda} \mathcal{L}(\mathbf{x} + \mathbf{x}', \lambda) \quad [\text{Dual ascent}]$$

⁷L. and L. Chen. From differential equation solvers to accelerated first-order methods for convex optimization. MAPR, 195: 735–781, 2022.

Accelerated primal-dual flow

- ▶ **Main idea:** Combining primal-dual form with **acceleration**
- ▶ Nesterov accelerated gradient flow for $\min F(\mathbf{x})$ ⁷

$$\gamma \mathbf{x}'' + (\mu + \gamma) \mathbf{x}' + \partial F(\mathbf{x}) \ni 0 \quad (\text{NAG flow})$$

- ▶ Combining NAG-flow with **(PD Flow)** gives

$$\gamma \mathbf{x}'' + (\mu + \gamma) \mathbf{x}' \in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda) \quad [\text{Accelerated primal descent}]$$

$$\theta \lambda' = \nabla_{\lambda} \mathcal{L}(\mathbf{x} + \mathbf{x}', \lambda) \quad [\text{Dual ascent}]$$

⁷L. and L. Chen. From differential equation solvers to accelerated first-order methods for convex optimization. MAPR, 195: 735–781, 2022.

Accelerated primal-dual flow

- ▶ **Main idea:** Combining primal-dual form with **acceleration**

- ▶ Nesterov accelerated gradient flow for $\min F(\mathbf{x})$ ⁷

$$\gamma \mathbf{x}'' + (\mu + \gamma) \mathbf{x}' + \partial F(\mathbf{x}) \ni 0 \quad (\text{NAG flow})$$

- ▶ Combining NAG-flow with **(PD Flow)** gives

$$\gamma \mathbf{x}'' + (\mu + \gamma) \mathbf{x}' \in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda) \quad [\text{Accelerated primal descent}]$$

$$\theta \lambda' = \nabla_{\lambda} \mathcal{L}(\mathbf{x} + \mathbf{x}', \lambda) \quad [\text{Dual ascent}]$$

- ▶ Introduce $\bar{\mathbf{x}} = \mathbf{x} + \mathbf{x}'$ to obtain

$$\begin{aligned} \mathbf{x}' &= \bar{\mathbf{x}} - \mathbf{x} \\ \gamma \bar{\mathbf{x}}' &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda) + \mu(\mathbf{x} - \bar{\mathbf{x}}) \\ \theta \lambda' &= \nabla_{\lambda} \mathcal{L}(\bar{\mathbf{x}}, \lambda) \end{aligned}$$

(**APD Flow**)

- First-order system is convenient for discretization and analysis than second-order ODE

⁷L. and L. Chen. From differential equation solvers to accelerated first-order methods for convex optimization. MAPR, 195: 735–781, 2022.

Two block APD flow

- Recall that $\mu = \mu_F = \min\{\mu_f, \mu_g\}$, and $\mu = 0$ when $\mu_f \mu_g = 0$

Two block APD flow

- ▶ Recall that $\mu = \mu_F = \min\{\mu_f, \mu_g\}$, and $\mu = 0$ when $\mu_f \mu_g = 0$
- ▶ For two block problem (1), we introduce **block diagonal scaling parameters**:

$$\gamma \implies \Gamma = \text{diag}(\gamma I, \beta I), \quad \mu \implies \boldsymbol{\mu} = \text{diag}(\mu_f I, \mu_g I)$$

$$\gamma' = \mu - \gamma \implies \Gamma' = \boldsymbol{\mu} - \Gamma$$

Two block APD flow

- ▶ Recall that $\mu = \mu_F = \min\{\mu_f, \mu_g\}$, and $\mu = 0$ when $\mu_f \mu_g = 0$
- ▶ For two block problem (1), we introduce **block diagonal scaling parameters**:

$$\gamma \implies \Gamma = \text{diag}(\gamma I, \beta I), \quad \mu \implies \boldsymbol{\mu} = \text{diag}(\mu_f I, \mu_g I)$$

$$\gamma' = \mu - \gamma \implies \Gamma' = \boldsymbol{\mu} - \Gamma$$

- ▶ Two block APD flow

$$\begin{aligned} \mathbf{x}' &= \bar{\mathbf{x}} - \mathbf{x} \\ \Gamma \bar{\mathbf{x}}' &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda) + \boldsymbol{\mu}(\mathbf{x} - \bar{\mathbf{x}}) \\ \theta \lambda' &= \nabla_{\lambda} \mathcal{L}(\bar{\mathbf{x}}, \lambda) \end{aligned}$$

(2block-APD Flow)

Two block APD flow

- ▶ Recall that $\mu = \mu_F = \min\{\mu_f, \mu_g\}$, and $\mu = 0$ when $\mu_f \mu_g = 0$
- ▶ For two block problem (1), we introduce **block diagonal scaling parameters**:

$$\gamma \implies \Gamma = \text{diag}(\gamma I, \beta I), \quad \mu \implies \boldsymbol{\mu} = \text{diag}(\mu_f I, \mu_g I)$$

$$\gamma' = \mu - \gamma \implies \Gamma' = \boldsymbol{\mu} - \Gamma$$

- ▶ Two block APD flow

$$\begin{aligned} \mathbf{x}' &= \bar{\mathbf{x}} - \mathbf{x} \\ \Gamma \bar{\mathbf{x}}' &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \lambda) + \boldsymbol{\mu}(\mathbf{x} - \bar{\mathbf{x}}) \\ \theta \lambda' &= \nabla_{\lambda} \mathcal{L}(\bar{\mathbf{x}}, \lambda) \end{aligned}$$

(2block-APD Flow)

- ▶ Exponential decay of

$$\mathcal{E}(\mathbf{x}, \lambda) := \mathcal{L}(\mathbf{x}, \hat{\lambda}) - \mathcal{L}(\hat{\mathbf{x}}, \lambda) + \frac{1}{2} \|\bar{\mathbf{x}} - \hat{\mathbf{x}}\|_{\Gamma}^2 + \frac{\theta}{2} \|\lambda - \hat{\lambda}\|^2$$

Introduction

PDHG and CP

Primal-Dual Flow Dynamics

Accelerated Primal-Dual Flow

Time Discretization

Numerical Results

A discretization template

- Denote the step size $\alpha_k > 0$ and consider

$$\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\alpha_k} = \bar{\mathbf{x}}_{k+1} - \mathbf{x}_{k+1}$$

$$\Gamma_k \frac{\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k}{\alpha_k} \in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}_{k+1}, \bar{\lambda}_{k+1}) + \boldsymbol{\mu}(\mathbf{x}_{k+1} - \bar{\mathbf{x}}_{k+1})$$

$$\theta_k \frac{\lambda_{k+1} - \lambda_k}{\alpha_k} = \mathbf{A} \bar{\mathbf{x}}_{k+1} - b$$

A discretization template

- Denote the step size $\alpha_k > 0$ and consider

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\alpha_k} &= \bar{\mathbf{x}}_{k+1} - \mathbf{x}_{k+1} \\ \Gamma_k \frac{\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k}{\alpha_k} &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}_{k+1}, \bar{\lambda}_{k+1}) + \boldsymbol{\mu}(\mathbf{x}_{k+1} - \bar{\mathbf{x}}_{k+1}) \\ \theta_k \frac{\lambda_{k+1} - \lambda_k}{\alpha_k} &= \mathbf{A} \bar{\mathbf{x}}_{k+1} - b\end{aligned}$$

- Parameter equation

$$\frac{\Gamma_{k+1} - \Gamma_k}{\alpha_k} = \boldsymbol{\mu} - \Gamma_{k+1}, \quad \frac{\theta_{k+1} - \theta_k}{\alpha_k} = -\theta_{k+1}$$

A discretization template

- Denote the step size $\alpha_k > 0$ and consider

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\alpha_k} &= \bar{\mathbf{x}}_{k+1} - \mathbf{x}_{k+1} \\ \Gamma_k \frac{\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k}{\alpha_k} &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}_{k+1}, \bar{\lambda}_{k+1}) + \boldsymbol{\mu}(\mathbf{x}_{k+1} - \bar{\mathbf{x}}_{k+1}) \\ \theta_k \frac{\lambda_{k+1} - \lambda_k}{\alpha_k} &= \mathbf{A} \bar{\mathbf{x}}_{k+1} - b\end{aligned}$$

- Parameter equation

$$\frac{\Gamma_{k+1} - \Gamma_k}{\alpha_k} = \boldsymbol{\mu} - \Gamma_{k+1}, \quad \frac{\theta_{k+1} - \theta_k}{\alpha_k} = -\theta_{k+1}$$

- $\bar{\lambda}_{k+1}$ is crucial to decouple \mathbf{x}_{k+1} and $\bar{\mathbf{x}}_{k+1}$:

A discretization template

- Denote the step size $\alpha_k > 0$ and consider

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\alpha_k} &= \bar{\mathbf{x}}_{k+1} - \mathbf{x}_{k+1} \\ \Gamma_k \frac{\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k}{\alpha_k} &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}_{k+1}, \bar{\lambda}_{k+1}) + \boldsymbol{\mu}(\mathbf{x}_{k+1} - \bar{\mathbf{x}}_{k+1}) \\ \theta_k \frac{\lambda_{k+1} - \lambda_k}{\alpha_k} &= \mathbf{A} \bar{\mathbf{x}}_{k+1} - b\end{aligned}$$

- Parameter equation

$$\frac{\Gamma_{k+1} - \Gamma_k}{\alpha_k} = \boldsymbol{\mu} - \Gamma_{k+1}, \quad \frac{\theta_{k+1} - \theta_k}{\alpha_k} = -\theta_{k+1}$$

- $\bar{\lambda}_{k+1}$ is crucial to decouple \mathbf{x}_{k+1} and $\bar{\mathbf{x}}_{k+1}$:
 - $\bar{\lambda}_{k+1} = \lambda_{k+1}$: fully coupled \implies nonlinear & nonsmooth ✗

A discretization template

- Denote the step size $\alpha_k > 0$ and consider

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\alpha_k} &= \bar{\mathbf{x}}_{k+1} - \mathbf{x}_{k+1} \\ \Gamma_k \frac{\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k}{\alpha_k} &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}_{k+1}, \bar{\lambda}_{k+1}) + \boldsymbol{\mu}(\mathbf{x}_{k+1} - \bar{\mathbf{x}}_{k+1}) \\ \theta_k \frac{\lambda_{k+1} - \lambda_k}{\alpha_k} &= \mathbf{A} \bar{\mathbf{x}}_{k+1} - b\end{aligned}$$

- Parameter equation

$$\frac{\Gamma_{k+1} - \Gamma_k}{\alpha_k} = \boldsymbol{\mu} - \Gamma_{k+1}, \quad \frac{\theta_{k+1} - \theta_k}{\alpha_k} = -\theta_{k+1}$$

- $\bar{\lambda}_{k+1}$ is crucial to decouple \mathbf{x}_{k+1} and $\bar{\mathbf{x}}_{k+1}$:
 - $\bar{\lambda}_{k+1} = \lambda_{k+1}$: fully coupled \implies nonlinear & nonsmooth ✗
 - $\bar{\lambda}_{k+1} \sim (u_{k+1}, \bar{u}_{k+1})$: weakly coupled \implies splitting ✓

A discretization template

- Denote the step size $\alpha_k > 0$ and consider

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\alpha_k} &= \bar{\mathbf{x}}_{k+1} - \mathbf{x}_{k+1} \\ \Gamma_k \frac{\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k}{\alpha_k} &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}_{k+1}, \bar{\lambda}_{k+1}) + \boldsymbol{\mu}(\mathbf{x}_{k+1} - \bar{\mathbf{x}}_{k+1}) \\ \theta_k \frac{\lambda_{k+1} - \lambda_k}{\alpha_k} &= \mathbf{A} \bar{\mathbf{x}}_{k+1} - b\end{aligned}$$

- Parameter equation

$$\frac{\Gamma_{k+1} - \Gamma_k}{\alpha_k} = \boldsymbol{\mu} - \Gamma_{k+1}, \quad \frac{\theta_{k+1} - \theta_k}{\alpha_k} = -\theta_{k+1}$$

- $\bar{\lambda}_{k+1}$ is crucial to decouple \mathbf{x}_{k+1} and $\bar{\mathbf{x}}_{k+1}$:

- $\bar{\lambda}_{k+1} = \lambda_{k+1}$: fully coupled \implies nonlinear & nonsmooth ✗
- $\bar{\lambda}_{k+1} \sim (u_{k+1}, \bar{u}_{k+1})$: weakly coupled \implies splitting ✓
- $\bar{\lambda}_{k+1} \sim (v_{k+1}, \bar{v}_{k+1})$: weakly coupled \implies splitting ✓

A discretization template

- Denote the step size $\alpha_k > 0$ and consider

$$\begin{aligned}\frac{\mathbf{x}_{k+1} - \mathbf{x}_k}{\alpha_k} &= \bar{\mathbf{x}}_{k+1} - \mathbf{x}_{k+1} \\ \Gamma_k \frac{\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k}{\alpha_k} &\in -\partial_{\mathbf{x}} \mathcal{L}(\mathbf{x}_{k+1}, \bar{\lambda}_{k+1}) + \boldsymbol{\mu}(\mathbf{x}_{k+1} - \bar{\mathbf{x}}_{k+1}) \\ \theta_k \frac{\lambda_{k+1} - \lambda_k}{\alpha_k} &= \mathbf{A} \bar{\mathbf{x}}_{k+1} - b\end{aligned}$$

- Parameter equation

$$\frac{\Gamma_{k+1} - \Gamma_k}{\alpha_k} = \boldsymbol{\mu} - \Gamma_{k+1}, \quad \frac{\theta_{k+1} - \theta_k}{\alpha_k} = -\theta_{k+1}$$

- $\bar{\lambda}_{k+1}$ is crucial to decouple \mathbf{x}_{k+1} and $\bar{\mathbf{x}}_{k+1}$:

- $\bar{\lambda}_{k+1} = \lambda_{k+1}$: fully coupled \implies nonlinear & nonsmooth ✗
- $\bar{\lambda}_{k+1} \sim (u_{k+1}, \bar{u}_{k+1})$: weakly coupled \implies splitting ✓
- $\bar{\lambda}_{k+1} \sim (v_{k+1}, \bar{v}_{k+1})$: weakly coupled \implies splitting ✓
- $\bar{\lambda}_{k+1} \sim (u_k, v_k)$: decoupled \implies splitting + parallelization ✓

One iteration analysis

Discrete Lyapunov function (From the continuous one)

$$\mathcal{E}_k := \mathcal{L}(\mathbf{x}_k, \hat{\lambda}) - \mathcal{L}(\hat{\mathbf{x}}, \lambda_k) + \frac{1}{2} \|\bar{\mathbf{x}}_k - \hat{\mathbf{x}}\|_{\Gamma_k}^2 + \frac{\theta_k}{2} \|\lambda_k - \hat{\lambda}\|^2$$

Lemma 1

$$\mathcal{E}_{k+1} - \mathcal{E}_k \leq -\alpha_k \mathcal{E}_{k+1} + \frac{\theta_k}{2} \|\lambda_{k+1} - \bar{\lambda}_{k+1}\|^2 - \frac{1}{2} \|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k\|_{\Gamma_k}^2$$

One iteration analysis

Discrete Lyapunov function (From the continuous one)

$$\mathcal{E}_k := \mathcal{L}(\mathbf{x}_k, \hat{\lambda}) - \mathcal{L}(\hat{\mathbf{x}}, \lambda_k) + \frac{1}{2} \|\bar{\mathbf{x}}_k - \hat{\mathbf{x}}\|_{\Gamma_k}^2 + \frac{\theta_k}{2} \|\lambda_k - \hat{\lambda}\|^2$$

Lemma 1

$$\mathcal{E}_{k+1} - \mathcal{E}_k \leq -\alpha_k \mathcal{E}_{k+1} + \frac{\theta_k}{2} \|\lambda_{k+1} - \bar{\lambda}_{k+1}\|^2 - \frac{1}{2} \|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k\|_{\Gamma_k}^2$$

► $\bar{\lambda}_{k+1}$ is also crucial for convergence:

One iteration analysis

Discrete Lyapunov function (From the continuous one)

$$\mathcal{E}_k := \mathcal{L}(\mathbf{x}_k, \hat{\lambda}) - \mathcal{L}(\hat{\mathbf{x}}, \lambda_k) + \frac{1}{2} \|\bar{\mathbf{x}}_k - \hat{\mathbf{x}}\|_{\Gamma_k}^2 + \frac{\theta_k}{2} \|\lambda_k - \hat{\lambda}\|^2$$

Lemma 1

$$\mathcal{E}_{k+1} - \mathcal{E}_k \leq -\alpha_k \mathcal{E}_{k+1} + \frac{\theta_k}{2} \|\lambda_{k+1} - \bar{\lambda}_{k+1}\|^2 - \frac{1}{2} \|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k\|_{\Gamma_k}^2$$

► $\bar{\lambda}_{k+1}$ is also crucial for convergence:

- $\bar{\lambda}_{k+1} - \lambda_{k+1} = 0$: perfect ✓

One iteration analysis

Discrete Lyapunov function (From the continuous one)

$$\mathcal{E}_k := \mathcal{L}(\mathbf{x}_k, \hat{\lambda}) - \mathcal{L}(\hat{\mathbf{x}}, \lambda_k) + \frac{1}{2} \|\bar{\mathbf{x}}_k - \hat{\mathbf{x}}\|_{\Gamma_k}^2 + \frac{\theta_k}{2} \|\lambda_k - \hat{\lambda}\|^2$$

Lemma 1

$$\mathcal{E}_{k+1} - \mathcal{E}_k \leq -\alpha_k \mathcal{E}_{k+1} + \frac{\theta_k}{2} \|\lambda_{k+1} - \bar{\lambda}_{k+1}\|^2 - \frac{1}{2} \|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k\|_{\Gamma_k}^2$$

► $\bar{\lambda}_{k+1}$ is also crucial for convergence:

- $\bar{\lambda}_{k+1} - \lambda_{k+1} = 0$: perfect ✓
- $\bar{\lambda}_{k+1} - \lambda_{k+1} \sim (\bar{u}_{k+1} - \bar{u}_k)$: controllable ✓

One iteration analysis

Discrete Lyapunov function (From the continuous one)

$$\mathcal{E}_k := \mathcal{L}(\mathbf{x}_k, \hat{\lambda}) - \mathcal{L}(\hat{\mathbf{x}}, \lambda_k) + \frac{1}{2} \|\bar{\mathbf{x}}_k - \hat{\mathbf{x}}\|_{\Gamma_k}^2 + \frac{\theta_k}{2} \|\lambda_k - \hat{\lambda}\|^2$$

Lemma 1

$$\mathcal{E}_{k+1} - \mathcal{E}_k \leq -\alpha_k \mathcal{E}_{k+1} + \frac{\theta_k}{2} \|\lambda_{k+1} - \bar{\lambda}_{k+1}\|^2 - \frac{1}{2} \|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k\|_{\Gamma_k}^2$$

► $\bar{\lambda}_{k+1}$ is also crucial for convergence:

- $\bar{\lambda}_{k+1} - \lambda_{k+1} = 0$: perfect ✓
- $\bar{\lambda}_{k+1} - \lambda_{k+1} \sim (\bar{u}_{k+1} - \bar{u}_k)$: controllable ✓
- $\bar{\lambda}_{k+1} - \lambda_{k+1} \sim (\bar{v}_{k+1} - \bar{v}_k)$: controllable ✓

The first IMEX scheme

- ▶ The implicit-explicit choice $\bar{\lambda}_{k+1} = \lambda_k + \alpha_k/\theta_k(A\bar{u}_{k+1} + B\bar{v}_k - b)$

The first IMEX scheme

- ▶ The implicit-explicit choice $\bar{\lambda}_{k+1} = \lambda_k + \alpha_k/\theta_k(A\bar{u}_{k+1} + B\bar{v}_k - b)$
- ▶ Controllable gap

$$\lambda_{k+1} - \bar{\lambda}_{k+1} = \alpha_k/\theta_k A(\bar{u}_{k+1} - \bar{u}_k)$$

The first IMEX scheme

- ▶ The implicit-explicit choice $\bar{\lambda}_{k+1} = \lambda_k + \alpha_k/\theta_k(A\bar{u}_{k+1} + B\bar{v}_k - b)$
- ▶ Controllable gap

$$\lambda_{k+1} - \bar{\lambda}_{k+1} = \alpha_k/\theta_k A(\bar{u}_{k+1} - \bar{u}_k)$$

- ▶ Optimization algorithm (informal)

$$\left\{ \begin{array}{l} u_{k+1} = \underset{u}{\operatorname{argmin}} \left\{ \mathcal{L}_{\sigma_k}(u, v_k, \hat{\lambda}_k) + \frac{\eta_{f,k}}{2\alpha_k^2} \|u - \tilde{u}_k\|^2 \right\} \\ \bar{u}_{k+1} = u_{k+1} + (u_{k+1} - u_k)/\alpha_k \\ v_{k+1} = \mathbf{prox}_{\tau_k g}(\tilde{v}_k - \tau_k B^* \bar{\lambda}_k) \\ \bar{v}_{k+1} = v_{k+1} + (v_{k+1} - v_k)/\alpha_k \\ \lambda_{k+1} = \lambda_k + \alpha_k/\theta_k(A\bar{u}_{k+1} + B\bar{v}_{k+1} - b) \end{array} \right. \quad (\text{Fast-ADMM-1})$$

with last iterate u_k, v_k, λ_k , intermediate sequence $\hat{\lambda}_k, \bar{\lambda}_k, \tilde{u}_k, \tilde{v}_k$ and parameter sequence $\tau_k, \eta_{f,k}, \eta_{g,k}$.

The first IMEX scheme

- ▶ The implicit-explicit choice $\bar{\lambda}_{k+1} = \lambda_k + \alpha_k / \theta_k (A \bar{u}_{k+1} + B \bar{v}_k - b)$
- ▶ Controllable gap

$$\lambda_{k+1} - \bar{\lambda}_{k+1} = \alpha_k / \theta_k A (\bar{u}_{k+1} - \bar{u}_k)$$

- ▶ Optimization algorithm (informal)

$$\left\{ \begin{array}{l} u_{k+1} = \underset{u}{\operatorname{argmin}} \left\{ \mathcal{L}_{\sigma_k}(u, v_k, \hat{\lambda}_k) + \frac{\eta_{f,k}}{2\alpha_k^2} \|u - \tilde{u}_k\|^2 \right\} \\ \bar{u}_{k+1} = u_{k+1} + (u_{k+1} - u_k) / \alpha_k \\ v_{k+1} = \operatorname{prox}_{\tau_k g}(\tilde{v}_k - \tau_k B^* \bar{\lambda}_k) \\ \bar{v}_{k+1} = v_{k+1} + (v_{k+1} - v_k) / \alpha_k \\ \lambda_{k+1} = \lambda_k + \alpha_k / \theta_k (A \bar{u}_{k+1} + B \bar{v}_{k+1} - b) \end{array} \right. \quad (\text{Fast-ADMM-1})$$

with last iterate u_k, v_k, λ_k , intermediate sequence $\hat{\lambda}_k, \bar{\lambda}_k, \tilde{u}_k, \tilde{v}_k$ and parameter sequence $\tau_k, \eta_{f,k}, \eta_{g,k}$.

- ▶ Sequential inner solvers: $\operatorname{prox}_{f+A^*A}$ and prox_g

- Take the step size $\alpha_k^2 \|B\|^2 = \theta_k \beta_k$

- ▶ Take the step size $\alpha_k^2 \|B\|^2 = \theta_k \beta_k$
- ▶ Convergence rate

$$E_k + R_k \lesssim \min \left\{ \frac{\|B\|}{k}, \frac{\|B\|^2}{\mu_g k^2} \right\}$$

where

$$R_k := \underbrace{\mathcal{L}(u_k, v_k, \hat{\lambda}) - \mathcal{L}(\hat{u}, \hat{v}, \lambda_k)}_{\text{Lagrange gap}}$$

$$E_k := \underbrace{\|Au_k + Bv_k - b\|}_{\text{Feasibility violation}} + \underbrace{\left| F(u_k, v_k) - \hat{F} \right|}_{\text{Objective residual}}$$

- ▶ Take the step size $\alpha_k^2 \|B\|^2 = \theta_k \beta_k$
- ▶ Convergence rate

$$E_k + R_k \lesssim \min \left\{ \frac{\|B\|}{k}, \frac{\|B\|^2}{\mu_g k^2} \right\}$$

where

$$R_k := \underbrace{\mathcal{L}(u_k, v_k, \hat{\lambda}) - \mathcal{L}(\hat{u}, \hat{v}, \lambda_k)}_{\text{Lagrange gap}}$$

$$E_k := \underbrace{\|Au_k + Bv_k - b\|}_{\text{Feasibility violation}} + \underbrace{\left| F(u_k, v_k) - \hat{F} \right|}_{\text{Objective residual}}$$

- ▶ When $B = -I, b = 0$,

$$0 \leq P(u_k) - P(\hat{u}) \lesssim \min \left\{ \frac{\|B\|}{k}, \frac{\|B\|^2}{\mu_g k^2} \right\}$$

- ▶ Take the step size $\alpha_k^2 \|B\|^2 = \theta_k \beta_k$
- ▶ Convergence rate

$$E_k + R_k \lesssim \min \left\{ \frac{\|B\|}{k}, \frac{\|B\|^2}{\mu_g k^2} \right\}$$

where

$$R_k := \underbrace{\mathcal{L}(u_k, v_k, \hat{\lambda}) - \mathcal{L}(\hat{u}, \hat{v}, \lambda_k)}_{\text{Lagrange gap}}$$

$$E_k := \underbrace{\|Au_k + Bv_k - b\|}_{\text{Feasibility violation}} + \underbrace{\left| F(u_k, v_k) - \hat{F} \right|}_{\text{Objective residual}}$$

- ▶ When $B = -I, b = 0$,

$$0 \leq P(u_k) - P(\hat{u}) \lesssim \min \left\{ \frac{\|B\|}{k}, \frac{\|B\|^2}{\mu_g k^2} \right\}$$

- ▶ Linearization of $f = f_1 + f_2$

$$\min \left\{ \frac{\|B\|}{\sqrt{\beta_0} k}, \frac{\|B\|^2}{\mu_g k^2} \right\} + \min \left\{ \frac{L_f}{\gamma_0 k^2}, \exp \left(-\frac{k}{4} \sqrt{\frac{\mu_f}{L_f}} \right) \right\}$$

More schemes

- ▶ The symmetric choice $\bar{\lambda}_{k+1} = \lambda_k + \alpha_k / \theta_k (A\bar{u}_k + B\bar{v}_{k+1} - b)$
 - Sequential inner solvers: prox_{g+B^*B} and prox_f :

$$\min \left\{ \frac{\|A\|}{k}, \frac{\|A\|^2}{\mu_f k^2} \right\}$$

- Linearization of $f = f_1 + f_2$

$$\min \left\{ \frac{\|A\|}{\sqrt{\gamma_0} k} + \frac{L_f}{\gamma_0 k^2}, \frac{\|A\|^2}{\mu_f k^2} + \exp \left(-\frac{k}{4} \sqrt{\frac{\mu_f}{L_f}} \right) \right\}$$

More schemes

- The symmetric choice $\bar{\lambda}_{k+1} = \lambda_k + \alpha_k / \theta_k (A\bar{u}_k + B\bar{v}_{k+1} - b)$
 - Sequential inner solvers: \mathbf{prox}_{g+B^*B} and \mathbf{prox}_f :

$$\min \left\{ \frac{\|A\|}{k}, \frac{\|A\|^2}{\mu_f k^2} \right\}$$

- Linearization of $f = f_1 + f_2$

$$\min \left\{ \frac{\|A\|}{\sqrt{\gamma_0} k} + \frac{L_f}{\gamma_0 k^2}, \frac{\|A\|^2}{\mu_f k^2} + \exp \left(-\frac{k}{4} \sqrt{\frac{\mu_f}{L_f}} \right) \right\}$$

- The second choice $\bar{\lambda}_{k+1} = \lambda_k + \alpha_k / \theta_k (A\bar{u}_k + B\bar{v}_k - b)$
 - Parallel inner solvers: \mathbf{prox}_f and \mathbf{prox}_g

$$\min \left\{ \frac{\|B\|}{k}, \frac{\|B\|^2}{\mu_g k^2} \right\} + \min \left\{ \frac{\|A\|}{k}, \frac{\|A\|^2}{\mu_f k^2} \right\}$$

- Linearization of $f = f_1 + f_2$

$$\min \left\{ \frac{\|B\|}{\sqrt{\beta_0} k}, \frac{\|B\|^2}{\mu_g k^2} \right\} + \min \left\{ \frac{\|A\|}{\sqrt{\gamma_0} k} + \frac{L_f}{\gamma_0 k^2}, \frac{\|A\|^2}{\mu_f k^2} + \exp \left(-\frac{k}{4} \sqrt{\frac{\mu_f}{L_f}} \right) \right\}.$$

- All can be extended to $g = g_1 + g_2$

Introduction

PDHG and CP

Primal-Dual Flow Dynamics

Accelerated Primal-Dual Flow

Time Discretization

Numerical Results

LAD regression

Consider the least absolute deviation (LAD) regression problem

$$\min_{x \in \mathbb{R}^n} P(x) := f(x) + \|Ax - b\|_1, \quad (4)$$

where $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$ are given data with $m \ll n$, and f is a regularization function. Consider two types of regularizer:

- Case 1: $f(x) = \lambda \|x\|_1$,
- Case 2: $f(x) = \lambda \|x\|_1 + \mu_f/2 \|x\|^2$.

We generate the matrix A from the standard normal distribution and set $b = Ax^\# + e$, where $x^\#$ is sparse and e is a Gaussian noise.




图 1: LAD regression, Case 1, with $(m, n) = (400, 4000)$ and sparsity 10%.

No worse (or comparable) convergence but good sparsity







图 2: LAD regression, Case 2, with $(m, n) = (400, 4000)$ and sparsity 10%.

No worse (or comparable) convergence but good sparsity

References

This talk is mainly based on



[Hao Luo, and ZiHang Zhang.](#)

A unified differential equation solver approach for separable convex optimization: splitting, acceleration and nonergodic rate.
arXiv:2109.13467v2, 2023. (Under review)

Other related references



[Hao Luo, and Long Chen.](#)

From differential equation solvers to accelerated first-order methods for convex optimization. *Math. Program.* 195:735–781, 2022.



[Hao Luo.](#)

A universal accelerated primal-dual method for convex optimization problems. *JOTA*, 2024.



[Hao Luo.](#)

A primal-dual flow for affine constrained convex optimization.
ESAIM : COCV, 2022.



[Hao Luo.](#)

Accelerated primal-dual methods for linearly constrained convex optimization problems. *arXiv:2109.12604*, 2021.

Thanks for your listening!

Any questions?