

Statistical Inference Chapter 1

Gallant Tsao

July 2, 2024

1. (a) $\Omega = \{(x_1, x_2, x_3, x_4) : x_i \in \{H, T\}\}$.
(b) If there are N leaves on the plant, $\Omega = [N]$.
(c) $\Omega = \{t : t \in \mathbb{R}, t \geq 0\}$.
(d) $\Omega = \{w : w \in \mathbb{R}_+\}$.
(e) If there are n components, $\Omega = \{i/n : i \in \{0, 1, \dots, n\}\}$.
2. (a)

$$\begin{aligned}x \in A \setminus B &\iff x \in A \text{ and } x \notin B \\&\iff x \in A \text{ and } x \notin A \cap B \\&\iff x \in A \setminus (A \cap B).\end{aligned}$$

Also,

$$\begin{aligned}x \in A \setminus B &\iff x \in A \text{ and } x \notin B \\&\iff x \in A \text{ and } x \in B^c \\&\iff x \in A \cap B^c.\end{aligned}$$

Therefore $A \setminus B = A \setminus (A \cap B) = A \cap B^c$.

- (b) By the distributive law,

$$\begin{aligned}(B \cap A) \cup (B \cap A^c) &= B \cap (A \cup A^c) \\&= B.\end{aligned}$$

- (c)

$$\begin{aligned}x \in B \setminus A &\iff x \in B \text{ and } x \notin A \\&\iff x \in B \text{ and } x \in A^c \\&\iff x \in B \cap A^c.\end{aligned}$$

- (d) From part b), we have

$$\begin{aligned}A \cup B &= A \cup ((B \cap A) \cup (B \cap A^c)) \\&= A \cup (B \cap A) \cup A \cup (B \cap A^c) \\&= A \cup A \cup (B \cap A^c) \\&= A \cup (B \cap A^c).\end{aligned}$$

3. (a)

$$\begin{aligned}
 x \in A \cup B &\iff x \in A \text{ or } x \in B \\
 &\iff x \in B \cup A. \\
 x \in A \cap B &\iff x \in A \text{ and } x \in B \\
 &\iff x \in B \cap A.
 \end{aligned}$$

(b)

$$\begin{aligned}
 x \in A \cup (B \cup C) &= x \in A \text{ or } x \in B \cup C \\
 &= x \in A \cup B \text{ or } x \in C \\
 &= x \in (A \cup B) \cup C.
 \end{aligned}$$

(c)

$$\begin{aligned}
 x \in (A \cup B)^c &\iff x \notin A \cup B \\
 &\iff x \in A^c \text{ and } x \in B^c \\
 &\iff x \in A^c \cap B^c. \\
 x \in (A \cap B)^c &\iff x \notin A \cap B \\
 &\iff x \in A^c \text{ or } x \in B^c \\
 &\iff x \in A^c \cup B^c.
 \end{aligned}$$

4. (a) This is $P(A \cup B)$, so we get $P(A) + P(B) - P(A \cap B)$.

(b) This is $P(A \Delta B)$, so we get $P(A) + P(B) - 2P(A \cap B)$.

(c) This is again $P(A \cup B)$, so we get $P(A) + P(B) - P(A \cap B)$.

(d) This is $P((A \cap B)^c)$, so we get $1 - P(A \cap B)$.

5. (a) $A \cap B \cap C = \{\text{a U.S. birth resulting in identical twin females}\}$.

(b) $P(A \cap B \cap C) = \frac{1}{90} \cdot \frac{1}{3} \cdot \frac{1}{2} = \frac{1}{540}$.

6. $p_0 = (1 - u)(1 - w)$, $p_1 = u(1 - w) + w(1 - u)$, $p_2 = uw$. For them to be equal,

$$\begin{aligned}
 p_0 = p_2 &\implies 1 - u - w + uw = uw \\
 &\implies u + w = 1, \\
 p_1 = p_2 &\implies u + w - 2uw = uw \\
 &\implies uw = \frac{1}{3}.
 \end{aligned}$$

The above two equations imply $u(1 - u) = \frac{1}{3}$, which has no real solutions in \mathbb{R} . Therefore we can't choose such u, w satisfying $p_0 = p_1 = p_2$.

7. (a) This is just having an extra case of hitting outside of the dart board. So

$$P(\text{scoring } i \text{ points}) = \begin{cases} 1 - \frac{\pi r^2}{A} & i = 0 \\ \frac{\pi r^2}{A} \cdot \frac{1}{5^2} ((6 - i)^2 - (5 - i)^2) & i = 1, \dots, 5 \end{cases}$$

(b)

$$\begin{aligned}
P(\text{scoring } i \text{ points} | \text{board is hit}) &= \frac{P(\text{scoring } i \text{ points, board is hit})}{P(\text{board is hit})} \\
&= \frac{\pi r^2}{A} \cdot \frac{1}{5^2} ((6-i)^2 - (5-i)^2) / \frac{\pi r^2}{A} \\
&= \frac{1}{5^2} ((6-i)^2 - (5-i)^2), \quad i = 1, \dots, 5
\end{aligned}$$

For $i = 0$, we will definitely score given that we hit the board so $P(\text{scoring } 0 \text{ points} | \text{board is hit}) = 0$, which is consistent with the probability distribution in Example 1.2.7 as well.

8. (a) From the example given,

$$P(\text{scoring } i \text{ points}) = \frac{(6-i)^2 - (5-i)^2}{5^2}, \quad i = 1, \dots, 5.$$

(b) Expanding the above,

$$\frac{(6-i)^2 - (5-i)^2}{5^2} = \frac{11-2i}{r^2},$$

which is a decreasing function of i .

(c)

$$\frac{11-2i}{5^2} > 0 \text{ for } i = 1, \dots, 5$$

hence the first axiom is satisfied.

$$P(S) = P(\text{hitting the board}) = 1,$$

hence the second axiom is satisfied. For $i \neq j$,

$$P(i \cup j) = \text{Area of ring } i + \text{Area of ring } j = P(i) + P(j),$$

hence the third axiom is satisfied so $P(\text{scoring } i \text{ points})$ is a probability function.

9. (a) Suppose $x \in (\cup_{\alpha} A_{\alpha})^c$. Then $x \notin A_{\alpha}$ for all $\alpha \in \Gamma$ so $x \in A_{\alpha}^c$ for all $\alpha \in \Gamma$. Therefore $x \in \cap_{\alpha} A_{\alpha}^c$.

Now suppose $x \in \cap_{\alpha} A_{\alpha}^c$. Then for all $\alpha \in \Gamma$, $x \in A_{\alpha}^c$ hence $x \notin A_{\alpha}$, then $x \notin \cup_{\alpha} A_{\alpha}$ so $x \in (\cup_{\alpha} A_{\alpha})^c$.

(b) Suppose $x \in (\cap_{\alpha} A_{\alpha})^c$. Then $x \notin \cap_{\alpha} A_{\alpha}$ so $x \notin A_{\alpha}$ for some $\alpha \in \Gamma$. Then $x \in A_{\alpha}^c$ for some $\alpha \in \Gamma$. Therefore $x \in \cup_{\alpha} A_{\alpha}^c$.

Now suppose $x \in \cup_{\alpha} A_{\alpha}^c$. Then $x \in A_{\alpha}^c$ for some $\alpha \in \Gamma$ so $x \notin A_{\alpha}$ for some $\alpha \in \Gamma$. Then $x \notin \cap_{\alpha} A_{\alpha}$ thus $x \in (\cap_{\alpha} A_{\alpha})^c$.

10. We have

$$\left(\bigcup_{i=1}^n A_i \right)^c = \bigcap_{i=1}^n A_i^c, \quad \left(\bigcap_{i=1}^n A_i \right)^c = \bigcup_{i=1}^n A_i^c$$

Proof of first equality:

Suppose $x \in (\cup_{i=1}^n A_i)^c$. Then $x \notin \cup_{i=1}^n A_i$ so $x \notin A_i$ for all i , meaning $x \in A_i^c$ for all i . Therefore $x \in \cap_{i=1}^n A_i^c$. Now suppose $x \in \cap_{i=1}^n A_i^c$. Then $x \notin A_i$ for all i , hence $x \notin \cup_{i=1}^n A_i$, hence $x \in (\cup_{i=1}^n A_i)^c$.

Proof of second equality:

Suppose $x \in (\cap_{i=1}^n A_i)^c$. Then $x \notin \cap_{i=1}^n A_i$ so $x \notin A_i$ and so $x \in A_i^c$ for some i , meaning $x \in \cup_{i=1}^n A_i^c$. Now suppose $x \in \cup_{i=1}^n A_i^c$. Then $x \notin A_i$ for some i hence $x \in (\cap_{i=1}^n A_i)^c$.

11. (a) $\emptyset \in \mathcal{B}$ hence property 1 is satisfied. $\emptyset^c = S \in \mathcal{B}$, $S^c = \emptyset \in \mathcal{B}$ hence property 2 is satisfied. $\emptyset \cup S = S \in \mathcal{B}$ hence property 3 is satisfied so \mathcal{B} is a sigma algebra.
- (b) \emptyset is a subset of S hence $\emptyset \in \mathcal{B}$ hence property 1 is satisfied. For any set $A \in \mathcal{B}$, $A^c = S \setminus A \in \mathcal{B}$ hence property 2 is satisfied. Any finite union of elements in \mathcal{B} will be a subset of S , which will be in \mathcal{B} so \mathcal{B} is a sigma algebra.
- (c) Suppose $\mathcal{F}_1, \mathcal{F}_2$ are sigma algebras on the sample space S . Since $\emptyset \in \mathcal{F}_1$ and $\emptyset \in \mathcal{F}_2$, $\emptyset \in \mathcal{F}_1 \cap \mathcal{F}_2$ so property 1 is satisfied. Suppose $A \subseteq \mathcal{F}_1 \cap \mathcal{F}_2$. Then $A \subseteq \mathcal{F}_1$ and $A \subseteq \mathcal{F}_2$. Since $\mathcal{F}_1, \mathcal{F}_2$ are both sigma algebras, $A^c \in \mathcal{F}_1$ and $A^c \in \mathcal{F}_2$. Therefore $A^c \in \mathcal{F}_1 \cap \mathcal{F}_2$ so property 2 is satisfied. Suppose $A_1, A_2, \dots \in \mathcal{F}_1 \cap \mathcal{F}_2$. Then $A_i \in \mathcal{F}_1$ and $A_i \in \mathcal{F}_2$. Since $\mathcal{F}_1, \mathcal{F}_2$ are both sigma algebras, $\cup_i A_i \in \mathcal{F}_1$ and $\cup_i A_i \in \mathcal{F}_2$ hence $\cup_i A_i \in \mathcal{F}_1 \cap \mathcal{F}_2$ hence property 3 is satisfied so $\mathcal{F}_1 \cap \mathcal{F}_2$ is a sigma algebra.
12. (a) 12.1
13. A, B cannot be disjoint. If they are,

$$P(A \cup B) = P(A) + P(B) = \frac{1}{3} + \frac{1}{4} = \frac{13}{12} > 1,$$

which is not possible.

14. For each element, we can choose to include it or exclude it in the subset. Since there are n elements, the number of subsets that can be formed is 2^n . A more formal proof can be done using bijections.
15. Now that the base case of $k = 2$ has been done, assume that this is true for k separate tasks. Then for each of the $n_1 \times n_2 \times \dots \times n_k$ ways, we have n_{k+1} choices for the $(k+1)$ th task. Therefore the entire job can be done in

$$\underbrace{1 \times n_{k+1} + 1 \times n_{k+1} + \dots + 1 \times n_{k+1}}_{n_1 \times \dots \times n_k \text{ terms}} = n_1 n_2 \dots n_{k+1}.$$

16. (a) 26^3
(b) $26^3 + 26^2$
(c) $26^4 + 26^3 + 26^2$
17. This is just choosing 2 numbers out of n of them, which is $\binom{n}{2} = \frac{n(n-1)}{2}$.
18. There are a total of n^n ways of putting n balls into n cells. For exactly one cell to be empty, there will also be another cell which has exactly 2 balls in it. Therefore there are $\binom{n}{2}$ ways of picking these special buckets. Since the order of putting in the balls matters, the answer is $\binom{n}{2} n! / n^n$.

- (a) By part (b), this is $\binom{6}{4} = 15$.
- (b) We can consider the n variables as bins, and the r partial derivatives as balls. Then we are putting r unlabeled balls into n unlabeled bins. There are a total of $\binom{n+r-1}{n-1} = \binom{n+r-1}{r}$ ways of doing this.
19. First of all, there are many different ways such that there is at least one call per day. Staying consistent with Casella's answers, if there is 6 calls on 1 day and 1 call on the other six days, we will denote this configuration as 6111111. All possible configs and the number of ways to form them are shown in the table below:

Config	Number of Ways	Answer
6111111	$7 \binom{12}{7} \cdot 6!$	4656960
5211111	$7 \binom{12}{5} \cdot 6 \binom{7}{2} \cdot 5!$	82825280
4221111	$7 \binom{12}{4} \cdot \binom{6}{2} \binom{8}{2} \binom{6}{2} \cdot 4!$	523908000
4311111	$7 \binom{12}{4} \cdot 6 \binom{8}{3} \cdot 5!$	139708800
3321111	$\binom{7}{2} \binom{12}{3} \binom{9}{3} \cdot 5 \binom{6}{2} \cdot 4!$	698544000
3222111	$7 \binom{12}{3} \cdot \binom{6}{3} \binom{9}{2} \binom{7}{2} \binom{5}{2} \cdot 3!$	1397088000
2222211	$\binom{7}{5} \binom{12}{2} \binom{10}{2} \binom{8}{2} \binom{6}{2} \binom{4}{2} \cdot 2!$	314344800
Total		3162075840

For example, for the config 6111111, there are $\binom{12}{6}$ ways for picking the calls for the day with 6 calls, 7 ways for the 6-call day to be in, and $6!$ ways for rearranging the rest of the 1-call days. A similar reasoning follows for the rest of the configs as well. All in all, the answer is about

$$\frac{3162075840}{7^{12}} \approx 0.2285.$$