

Table S1. Metabolites and their relevant information

Metabolite	Properties	References
2-Aminopimelic acid	<ul style="list-style-type: none"> <li>- forms lysine and contributes to cell wall structure in bacteria.</li> <li>- involved in plant morphogenesis.</li> <li>- likely associated with the cnidarian microbiome or translocated from the symbionts.</li> </ul>	(Berges et al., 1986; Valmaseda et al., 2005; Tabeta and Hirai, 2024)
2-Hydroxyglutaric acid	<ul style="list-style-type: none"> <li>- high levels in humans are associated with metabolic disorders.</li> <li>- metabolite is differentially expressed in Aiptasia colonised with different symbiont types.</li> <li>- implications for T-cell differentiation.</li> </ul>	(Tyrakis et al., 2016; Yamada et al., 2019; Du and Hu, 2021; Lust, 2022; Tsang Min Ching et al., 2022)
3-Aminoglutaric acid	<ul style="list-style-type: none"> <li>- the precursor of glutamine synthase which is important for nitrogen assimilation, amino acid synthesis and cellular metabolism.</li> <li>- commonly found in bacteria and plants.</li> <li>- likely associated with the cnidarian microbiome or translocated from the symbionts.</li> </ul>	(Patrice et al., 2001; Ito et al., 2022)
Galacturonic acid	<ul style="list-style-type: none"> <li>- found in plants and has anti-inflammatory properties.</li> <li>- produced by Symbiodiniaceae and may be important in symbiosis establishment and recognition.</li> <li>- likely translocated from the symbiont to the host.</li> </ul>	(Gerschenson, 2017; Tortorelli et al., 2022)
Gluconic acid	<ul style="list-style-type: none"> <li>- involved in glucose metabolism, and antioxidant NADPH production through the pentose phosphate pathway.</li> <li>- found in the Aiptasia metabolome.</li> <li>- increased only at T4 and not T16 in the current study.</li> </ul>	(MIETTINEN and LESKINEN, 1970; Matthews et al., 2017)
Glycolic acid	<ul style="list-style-type: none"> <li>- increases in aposymbiotic vs. symbiotic Aiptasia.</li> <li>- has anti-inflammatory properties by modulation of NFkB pathways and pro-inflammatory cytokines, however it can also induce apoptosis.</li> </ul>	(Yang et al., 2004; Tang et al., 2017; Lust, 2022)
Pantothenic acid	<ul style="list-style-type: none"> <li>- increases in aposymbiotic Aiptasia and Aiptasia colonised with heterologous symbionts vs. Aiptasia colonised with homologous symbionts.</li> <li>- precursor of CoA.</li> <li>- CoA can lead to the formation of triglycerides, phospholipids, antioxidants, cysteine and methionine.</li> </ul>	(Wada and Takagi, 2006; Matthews, 2017; Aloum et al., 2019; Chandel, 2021; Mignani et al., 2021; Filonenko and Gout, 2023)
Phosphoric acid	<ul style="list-style-type: none"> <li>- increases in aposymbiotic Aiptasia.</li> <li>- involved in cellular signalling, and the production of ATP, phospholipids and triglycerides.</li> </ul>	(Stillwell, 2016; Matthews, 2017; Kritmetapak and Kumar, 2021; Choi et al., 2023)
Rhamnose	<ul style="list-style-type: none"> <li>- usually found in bacteria, plants and fungi.</li> <li>- likely associated with the cnidarian microbiome or translocated from the symbionts.</li> <li>- found in plant and bacterial cell walls and increases in bacteria and plants under stressful conditions to adapt membrane structure or as an antioxidant defense.</li> </ul>	(Williams et al., 2004; Hillyer, 2016; Dastogeer et al., 2017; Jiang et al., 2021; Nguyen et al., 2021; Song et al., 2021)
Ribonic acid	<ul style="list-style-type: none"> <li>- sugar involved in ribose metabolism and production of nucleotides.</li> <li>- found in coral <i>Montipora capitata</i> metabolome.</li> <li>- increases in humans with diabetes</li> </ul>	(Ding et al., 2017; Matthews, 2017; Tofte et al., 2019; Curovic et al., 2020)
Sedoheptulose	<ul style="list-style-type: none"> <li>- sugar that is important for the production of nucleotides and is an intermediate in the pentose phosphate pathway.</li> <li>- mainly produced in plants.</li> <li>- likely translocated from the symbionts.</li> <li>- increased only at T4 and not T16 in the current study.</li> </ul>	(Benson et al., 1951)
Tryptamine	<ul style="list-style-type: none"> <li>- a group of monoamines that includes serotonin and melatonin that are well-studied in vertebrates but less studied in invertebrates.</li> <li>- increases in marine invertebrates, including cnidarians under stressful conditions</li> </ul>	(da Silveira et al., 2007; Liu et al., 2018)

## References

- Aloum, L., Brimson, C. A., Zhyvoloup, A., Baines, R., Baković, J., Filonenko, V., et al. (2019). Coenzyme A and protein CoAlation levels are regulated in response to oxidative stress and during morphogenesis in *Dictyostelium discoideum*. *Biochem. Biophys. Res. Commun.* 511, 294–299. doi:10.1016/j.bbrc.2019.02.031.
- Benson, A. A., Bassham, J. A., and Calvin, M. (1951). Sedoheptulose in photosynthesis by plants. *J. Am. Chem. Soc.* 73, 2970. doi:10.1021/ja01150a543.
- Berges, D. A., DeWolf, W. E., Dunn, G. L., Taggart, J. J., Grappel, S. F., Newman, D. J., et al. (1986). Peptides of 2-Aminopimelic Acid: Antibacterial Agents That Inhibit Diaminopimelic Acid Biosynthesis. *J. Med. Chem.* 29, 89–95. doi:10.1021/jm00151a015.
- Chandel, N. S. (2021). Lipid Metabolism. *Cold Spring Harb. Perspect. Biol.* 13. doi:10.1101/cshperspect.a040576.
- Choi, Y. M., Ajjaji, D., Fleming, K. D., Borbat, P. P., Jenkins, M. L., Moeller, B. E., et al. (2023). Structural insights into perilipin 3 membrane association in response to diacylglycerol accumulation. *Nat. Commun.* 14. doi:10.1038/s41467-023-38725-w.
- Curovic, V. R., Suvitaival, T., Mattila, I., Ahonen, L., Trošt, K., Theilade, S., et al. (2020). Circulating Metabolites and Lipids Are Associated to Diabetic Retinopathy in Individuals With Type 1 Diabetes. *Diabetes* 69, 2217–2226. doi:10.2337/db20-0104.
- da Silveira, R. B., Wille, A. C. M., Chaim, O. M., Appel, M. H., Silva, D. T., Franco, C. R. C., et al. (2007). Identification, cloning, expression and functional characterization of an astacin-like metalloprotease toxin from *Loxosceles intermedia* (brown spider) venom. *Biochem. J.* 406, 355–363. doi:10.1042/BJ20070363.
- Dastogeer, K. M. G., Li, H., Sivasithamparam, K., Jones, M. G. K., Du, X., Ren, Y., et al. (2017). Metabolic responses of endophytic *Nicotiana benthamiana* plants experiencing water stress. *Environ. Exp. Bot.* 143, 59–71. doi:https://doi.org/10.1016/j.envexpbot.2017.08.008.
- Ding, Z., Jia, S., Wang, Y., Xiao, J., and Zhang, Y. (2017). Phosphate stresses affect ionome and metabolome in tea plants. *Plant Physiol. Biochem.* 120, 30–39. doi:https://doi.org/10.1016/j.plaphy.2017.09.007.
- Du, X., and Hu, H. (2021). The Roles of 2-Hydroxyglutarate. *Front. Cell Dev. Biol.* 9, 1–13. doi:10.3389/fcell.2021.651317.
- Filonenko, V., and Gout, I. (2023). Discovery and functional characterisation of protein CoAlation and the antioxidant function of coenzyme A. *BBA Adv.* 3, 100075. doi:10.1016/j.bbadv.2023.100075.
- Gerschenson, L. N. (2017). The production of galacturonic acid enriched fractions and their functionality. *Food Hydrocoll.* 68, 23–30. doi:https://doi.org/10.1016/j.foodhyd.2016.11.030.
- Hillyer, K. E. (2016). Thermal Stress and Bleaching in the Cnidarian-Dinoflagellate Symbiosis: The Application of Metabolomics. 53.
- Ito, K., Miyamoto, H., Matsuura, M., Ishii, C., Tsuboi, A., Tsuji, N., et al. (2022). Noninvasive fecal metabolic profiling for the evaluation of characteristics of

- thermostable lactic acid bacteria, *Weizmannia coagulans* SANK70258, for broiler chickens. *J. Biosci. Bioeng.* 134, 105–115. doi:10.1016/j.jbiosc.2022.05.006.
- Jiang, N., Dillon, F. M., Silva, A., Gomez-Cano, L., and Grotewold, E. (2021). Rhamnose in plants - from biosynthesis to diverse functions. *Plant Sci.* 302, 110687. doi:10.1016/j.plantsci.2020.110687.
- Kritmetapak, K., and Kumar, R. (2021). Phosphate as a Signaling Molecule. *Calcif. Tissue Int.* 108, 16–31. doi:10.1007/s00223-019-00636-8.
- Liu, W., Mo, F., Jiang, G., Liang, H., Ma, C., Li, T., et al. (2018). Stress-induced mucus secretion and its composition by a combination of proteomics and metabolomics of the jellyfish *aurelia coerulea*. *Mar. Drugs* 16. doi:10.3390/md16090341.
- Matthews, J. L. (2017). The Nutritional Implications of Partner Switching in the Cnidarian-Dinoflagellate Symbiosis.
- Matthews, J. L., Crowder, C. M., Oakley, C. A., Lutz, A., Roessner, U., Meyer, E., et al. (2017). Optimal nutrient exchange and immune responses operate in partner specificity in the cnidarian-dinoflagellate symbiosis. *Proc. Natl. Acad. Sci. U. S. A.* 114, 13194–13199. doi:10.1073/pnas.1710733114.
- MIETTINEN, T. A., and LESKINEN, E. (1970). “GLUCURONIC ACID PATHWAY,” in *Metabolic Conjugation and Metabolic Hydrolysis*, ed. W. H. FISHMAN (Academic Press), 157–237. doi:https://doi.org/10.1016/B978-0-12-257601-0.50011-6.
- Mignani, L., Gnutti, B., Zizioli, D., and Finazzi, D. (2021). Coenzyme a Biochemistry: From Neurodevelopment to Neurodegeneration. *Brain Sci.* 11. doi:10.3390/brainsci11081031.
- Nguyen, P. T., Nguyen, T. T., Vo, T. N. T., Nguyen, T. T. X., Hoang, Q. K., and Nguyen, H. T. (2021). Response of *Lactobacillus plantarum* VAL6 to challenges of pH and sodium chloride stresses. *Sci. Rep.* 11, 1–9. doi:10.1038/s41598-020-80634-1.
- Patrice, R., Kelly, N., J., S. H., and F., R. M. (2001).  $\beta$ -Glutamate as a Substrate for Glutamine Synthetase. *Appl. Environ. Microbiol.* 67, 4458–4463. doi:10.1128/AEM.67.10.4458-4463.2001.
- Song, M., Liu, Y., Li, T., Liu, X., Hao, Z., Ding, S., et al. (2021). Plant Natural Flavonoids Against Multidrug Resistant Pathogens. *Adv. Sci.* 8, 1–11. doi:10.1002/advs.202100749.
- Stillwell, W. (2016). “Chapter 5 - Membrane Polar Lipids,” in *An Introduction to Biological Membranes (Second Edition)*, ed. W. Stillwell (Elsevier), 63–87. doi:https://doi.org/10.1016/B978-0-444-63772-7.00005-1.
- Tabeta, H., and Hirai, M. Y. (2024). 1-2-Aminopimelic acid acts as an auxin mimic to induce lateral root formation across diverse plant species. *FEBS Lett.* doi:10.1002/1873-3468.14908.
- Tofte, N., Suvitaival, T., Trost, K., Mattila, I. M., Theilade, S., Winther, S. A., et al. (2019). Metabolomic Assessment Reveals Alteration in Polyols and Branched Chain Amino Acids Associated With Present and Future Renal Impairment in a Discovery Cohort of 637 Persons With Type 1 Diabetes. *Front. Endocrinol. (Lausanne)*. 10, 1–11. doi:10.3389/fendo.2019.00818.
- Tyrakis, P. A., Palazon, A., Macias, D., Lee, K. L., Phan, A. T., Veliça, P., et al. (2016). S-2-hydroxyglutarate regulates CD8<sup>+</sup> T-lymphocyte fate. *Nature* 540, 236–241.

doi:10.1038/nature20165.

- Valmaseda, E. M. M. de, Campoy, S., Naranjo, L., Casqueiro, J., and Martín, J. F. (2005). Lysine is catabolized to 2-aminoadipic acid in *Penicillium chrysogenum* by an omega-aminotransferase and to saccharopine by a lysine 2-ketoglutarate reductase. Characterization of the omega-aminotransferase. *Mol. Genet. Genomics* 274, 272–282. doi:10.1007/s00438-005-0018-3.
- Wada, M., and Takagi, H. (2006). Metabolic pathways and biotechnological production of L-cysteine. *Appl. Microbiol. Biotechnol.* 73, 48–54. doi:10.1007/s00253-006-0587-z.
- Williams, R. J., Spencer, J. P. E., and Rice-Evans, C. (2004). Flavonoids: antioxidants or signalling molecules? *Free Radic. Biol. Med.* 36, 838–849. doi:https://doi.org/10.1016/j.freeradbiomed.2004.01.001.
- Yamada, T., Nabe, S., Toriyama, K., Suzuki, J., Inoue, K., Imai, Y., et al. (2019). Histone H3K27 Demethylase Negatively Controls the Memory Formation of Antigen-Stimulated CD8+ T Cells. *J. Immunol.* 202, 1088–1098. doi:10.4049/jimmunol.1801083.
- Yang, J.-H., Chou, C.-C., Cheng, Y.-W., Sheen, L.-Y., Chou, M.-C., Yu, H.-S., et al. (2004). Effects of glycolic acid on the induction of apoptosis via caspase-3 activation in human leukemia cell line (HL-60). *Food Chem. Toxicol. an Int. J. Publ. Br. Ind. Biol. Res. Assoc.* 42, 1777–1784. doi:10.1016/j.fct.2004.07.004.