

Matrices and Linear Systems Continued

Unit 3

Basic facts

- $Ax = \lambda x$, $(A - \lambda I)x = 0$, for a nonzero x

Basic facts

- $Ax = \lambda x$, $(A - \lambda I)x = 0$, for a nonzero x
- The eigenvalues $\lambda_1, \dots, \lambda_n$ are the roots of the characteristic polynomial $\det(A - \lambda I) = 0$

Basic facts

- $Ax = \lambda x$, $(A - \lambda I)x = 0$, for a nonzero x
- The eigenvalues $\lambda_1, \dots, \lambda_n$ are the roots of the characteristic polynomial $\det(A - \lambda I) = 0$
- The set of eigenvalues $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ is called the spectrum of A .

Basic facts

- $Ax = \lambda x, \quad (A - \lambda I)x = 0, \quad \text{for a nonzero } x$
- The eigenvalues $\lambda_1, \dots, \lambda_n$ are the roots of the characteristic polynomial $\det(A - \lambda I) = 0$
- The set of eigenvalues $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ is called the spectrum of A .
- $\text{trace}(A) = \sum_{i=1}^n a_{ii} = \sum_{i=1}^n \lambda_i, \quad \det(A) = \prod_{i=1}^n \lambda_i$

Basic facts

- $Ax = \lambda x, \quad (A - \lambda I)x = 0, \quad \text{for a nonzero } x$
- The eigenvalues $\lambda_1, \dots, \lambda_n$ are the roots of the characteristic polynomial $\det(A - \lambda I) = 0$
- The set of eigenvalues $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ is called the spectrum of A .
- $\text{trace}(A) = \sum_{i=1}^n a_{ii} = \sum_{i=1}^n \lambda_i, \quad \det(A) = \prod_{i=1}^n \lambda_i$
- $\rho(A) := \max\{|\lambda| : \lambda \in \sigma(A)\}$

Basic facts

- $Ax = \lambda x$, $(A - \lambda I)x = 0$, for a nonzero x
- The eigenvalues $\lambda_1, \dots, \lambda_n$ are the roots of the characteristic polynomial $\det(A - \lambda I) = 0$
- The set of eigenvalues $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ is called the spectrum of A .
- $\text{trace}(A) = \sum_{i=1}^n a_{ii} = \sum_{i=1}^n \lambda_i$, $\det(A) = \prod_{i=1}^n \lambda_i$
- $\rho(A) := \max\{|\lambda| : \lambda \in \sigma(A)\}$
- Work through Ex. 3.1 (1,2,4), Ex. 3.2, and Ex. 3.4. Do not submit.

The QR method

Based on several theoretical results: Eigenvalues of similar matrices, Schur's factorization, Gauss-Schmidt process.

A and B are similar if \exists a nonsingular P such that $B = PAP^{-1}$.

The QR method

Based on several theoretical results: Eigenvalues of similar matrices, Schur's factorization, Gauss-Schmidt process.

A and B are similar if \exists a nonsingular P such that $B = PAP^{-1}$.

Theorem. Similar matrices have the same eigenvalues.

The QR method

Based on several theoretical results: Eigenvalues of similar matrices, Schur's factorization, Gauss-Schmidt process.

A and B are similar if \exists a nonsingular P such that $B = PAP^{-1}$.

Theorem. Similar matrices have the same eigenvalues.

Schur's Theorem: Every square matrix is unitarily similar to a triangular matrix, i.e., for every square A , \exists a unitary U such that $T = UAU^*$, where T is upper triangular.

The QR method

Based on several theoretical results: Eigenvalues of similar matrices, Schur's factorization, Gauss-Schmidt process.

A and B are similar if \exists a nonsingular P such that $B = PAP^{-1}$.

Theorem. Similar matrices have the same eigenvalues.

Schur's Theorem: Every square matrix is unitarily similar to a triangular matrix, i.e., for every square A , \exists a unitary U such that $T = UAU^*$, where T is upper triangular.

QR algorithm produces QR-factorization $A = QR$, where Q is unitary and R is upper triangular.

Gauss-Schmidt process

Column vector form of A $A = [\mathbf{a}_1; \mathbf{a}_2; \dots; \mathbf{a}_n]$

Gauss-Schmidt process

Column vector form of A $A = [\mathbf{a}_1; \mathbf{a}_2; \dots; \mathbf{a}_n]$

Column vector form of Q $\mathbf{q}_i = \frac{\mathbf{a}'_i}{\|\mathbf{a}'_i\|}$

Gauss-Schmidt process

Column vector form of A $A = [\mathbf{a}_1; \mathbf{a}_2; \dots; \mathbf{a}_n]$

Column vector form of Q $\mathbf{q}_i = \frac{\mathbf{a}'_i}{\|\mathbf{a}'_i\|}$

where $\mathbf{a}'_i = \mathbf{a}_i - \sum_{k=1}^{i-1} (\mathbf{q}_k^T \mathbf{a}_i) \mathbf{q}_k, \quad i = 1, 2, \dots, n$

Gauss-Schmidt process

Column vector form of A $A = [\mathbf{a}_1; \mathbf{a}_2; \dots; \mathbf{a}_n]$

Column vector form of Q $\mathbf{q}_i = \frac{\mathbf{a}'_i}{\|\mathbf{a}'_i\|}$

where $\mathbf{a}'_i = \mathbf{a}_i - \sum_{k=1}^{i-1} (\mathbf{q}_k^T \mathbf{a}_i) \mathbf{q}_k, \quad i = 1, 2, \dots, n$

Entries of R $r_{ii} = \|\mathbf{a}'_i\|, \quad r_{ij} = \mathbf{q}_i^T \mathbf{a}_j, \quad i = 1, \dots, n, \quad j = i+1, \dots, n$

hw due next meeting

1. Create a random triadiagonal symmetric matrix of size n
2. Write your own single iteration QR code for triadiagonal symmetric matrices.
3. Test running time of your code vs the Matlab qr for $n = 3, \dots$ as as far as you can go hoping to outrun Matlab.
4. Write you own code to iterate your QR for triadiagonal symmetric matrices. Use several reasonable thresholds for stopping.
5. Test running time of your code vs the Matlab eig for $n = 3, \dots$ as as far as you can go hoping to outrun Matlab.

Operator norm

$$\|A\| = \max_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \max_{\|x\|=1} \|Ax\|$$

$$\|AB\| \leq \|A\| \|B\|$$

Note that it requires a vector norm. We use the following

1. $\|x\|_1 = \sum_{j=1}^n |x_j|$
2. $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$
3. $\|x\|_\infty = \max_{1 \leq j \leq n} |x_j|$

Operator norm

$$\|A\| = \max_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \max_{\|x\|=1} \|Ax\|$$

$$\|AB\| \leq \|A\| \|B\|$$

Note that it requires a vector norm. We use the following

1. $\|x\|_1 = \sum_{j=1}^n |x_j|$
2. $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$
3. $\|x\|_\infty = \max_{1 \leq j \leq n} |x_j|$

What are the subordinate matrix norms?

Subordinate norms

$$\|A\| = \max_{\|x\|=1} \|Ax\|$$

The easiest one to see is $\|x\|_\infty = \max_{1 \leq j \leq n} |x_j|$

Subordinate norms

$$\|A\| = \max_{\|x\|=1} \|Ax\|$$

The easiest one to see is $\|x\|_\infty = \max_{1 \leq j \leq n} |x_j|$

$$\|A\|_\infty = \max_{1 \leq k \leq m} \sum_{j=1}^n |a_{kj}|, \quad \text{the maximum row sum}$$

Try to think about $\|x\|_1 = \sum_{j=1}^n |x_j|$

Subordinate norms

$$\|A\| = \max_{\|x\|=1} \|Ax\|$$

The easiest one to see is $\|x\|_\infty = \max_{1 \leq j \leq n} |x_j|$

$$\|A\|_\infty = \max_{1 \leq k \leq m} \sum_{j=1}^n |a_{kj}|, \quad \text{the maximum row sum}$$

Try to think about $\|x\|_1 = \sum_{j=1}^n |x_j|$

$$\|A\|_1 = \max_{1 \leq j \leq m} \sum_{k=1}^n |a_{kj}|, \quad \text{the maximum column sum}$$

Subordinate and non-subordinate norms

The hardest one is $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$

Subordinate and non-subordinate norms

The hardest one is $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$

$$\|A\|_2 = \sqrt{\rho(A^*A)}, \quad \text{spectral norm}$$

Subordinate and non-subordinate norms

The hardest one is $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$

$$\|A\|_2 = \sqrt{\rho(A^*A)}, \quad \text{spectral norm}$$

Non-subordinate norm

$$\|A\| = \left(\sum_{k,j} a_{kj}^2 \right)^{1/2}, \quad \text{Frobenius Norm}$$

Subordinate and non-subordinate norms

The hardest one is $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$

$$\|A\|_2 = \sqrt{\rho(A^*A)}, \quad \text{spectral norm}$$

Non-subordinate norm

$$\|A\| = \left(\sum_{k,j} a_{kj}^2 \right)^{1/2}, \quad \text{Frobenius Norm}$$

$$\text{cond}(A) = \|A\| \|A^{-1}\|$$

Subordinate and non-subordinate norms

The hardest one is $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$

$$\|A\|_2 = \sqrt{\rho(A^*A)}, \quad \text{spectral norm}$$

Non-subordinate norm

$$\|A\| = \left(\sum_{k,j} a_{kj}^2 \right)^{1/2}, \quad \text{Frobenius Norm}$$

$$\text{cond}(A) = \|A\| \|A^{-1}\|$$

Ex. 4.1 Please write up and turn in