

Homework Assignment 10

Matthew Tiger

November 24, 2016

Problem 10.5.1. Show that the Cantor set C is a fixed point of the map $F : \mathcal{C}(\mathbb{R}) \rightarrow \mathcal{C}(\mathbb{R})$ defined by

$$F(A) = f_1(A) \cup f_2(A)$$

where $f_1(x) = x/3$ and $f_2(x) = x/3 + 2/3$ are contractions on \mathbb{R} .

Solution. Recall that the Cantor set C is defined as

$$C = \left\{ x \in [0, 1] \mid x = \sum_{n=1}^{\infty} \frac{a_n}{3^n}, \quad a_n \in \{0, 2\} \right\}. \quad (1)$$

We wish to show that $F(C) = C$. From the definitions of f_1 and f_2 , we have that

$$\begin{aligned} f_1(C) &= \left\{ x \in [0, 1] \mid x = \sum_{n=1}^{\infty} \frac{a_n}{3^{n+1}}, \quad a_n \in \{0, 2\} \right\} \\ &= \left\{ x \in [0, 1] \mid x = \sum_{n=1}^{\infty} \frac{b_n}{3^n}, \quad b_1 = 0, b_n = a_{n-1} \text{ for } n > 1 \right\} \\ f_2(C) &= \left\{ x \in [0, 1] \mid x = \frac{2}{3} + \sum_{n=1}^{\infty} \frac{a_n}{3^{n+1}}, \quad a_n \in \{0, 2\} \right\} \\ &= \left\{ x \in [0, 1] \mid x = \sum_{n=1}^{\infty} \frac{b_n}{3^n}, \quad b_1 = 2, b_n = a_{n-1} \text{ for } n > 1 \right\} \end{aligned} \quad (2)$$

From (2) and the definition of F , we see that

$$\begin{aligned} F(C) &= f_1(C) \cup f_2(C) \\ &= \left\{ x \in [0, 1] \mid x = \sum_{n=1}^{\infty} \frac{b_n}{3^n}, \quad b_n \in \{0, 2\} \right\}. \end{aligned}$$

But this is precisely the definition of the Cantor set C given in (1). Therefore, $F(C) = C$ and the Cantor set C is a fixed point of F . \square

Problem 10.5.2.*Solution.*

Problem 10.5.4.*Solution.*

Problem 10.5.7.*Solution.*