

Exam 2

Matthew Tiger

April 22, 2017

Problem 1. Find the inverse Laplace transform of the function

$$\bar{f}(s) = \frac{s}{(s-a)(s^2+b^2)}$$

for $a, b > 0$, by using the following three different approaches:

- i. Using partial fraction decomposition,
- ii. Applying the Convolution Theorem,
- iii. Applying Heaviside's Expansion Theorem.

Solution. We will now find the inverse Laplace transform of $\bar{f}(s)$ using the respective approaches listed above:

- i. From the partial fractions method, we see that

$$\bar{f}(s) = \frac{s}{(s-a)(s^2+b^2)} = \frac{c_0}{s-a} + \frac{d_1s+d_0}{s^2+b^2}.$$

Combining the rational fractions on the right side under a common denominator and equating the coefficients in the numerator we arrive at the following system of equations

$$\begin{aligned}c_0 + d_1 &= 0 \\d_0 - ad_1 &= 0 \\c_0b^2 - ad_0 &= 0.\end{aligned}$$

Solving this system, we see that $c_0 = \frac{a}{a^2+b^2}$, $d_1 = -\frac{a}{a^2+b^2}$, and $d_0 = \frac{b^2}{a^2+b^2}$. Thus, we have that

$$\bar{f}(s) = \frac{1}{a^2+b^2} \left[\frac{a}{s-a} - \frac{as}{s^2+b^2} + \frac{b^2}{s^2+b^2} \right].$$

From our table of Laplace transforms, we know that

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{1}{s-a}\right\} &= e^{at} \\ \mathcal{L}^{-1}\left\{\frac{s}{s^2+b^2}\right\} &= \cos bt \\ \mathcal{L}^{-1}\left\{\frac{b}{s^2+b^2}\right\} &= \sin bt.\end{aligned}$$

Therefore, the inverse Laplace transform of $\bar{f}(s)$ is

$$\begin{aligned}f(t) = \mathcal{L}^{-1}\{\bar{f}(s)\} &= \frac{1}{a^2+b^2} \left[a\mathcal{L}^{-1}\left\{\frac{1}{s-a}\right\} - a\mathcal{L}^{-1}\left\{\frac{s}{s^2+b^2}\right\} + b\mathcal{L}^{-1}\left\{\frac{b}{s^2+b^2}\right\} \right] \\ &= \frac{1}{a^2+b^2} [ae^{at} - a\cos bt + b\sin bt].\end{aligned}$$

ii. The Convolution Theorem states that if $\bar{f}(s) = \bar{g}(s)\bar{h}(s)$, then

$$f(t) = \mathcal{L}^{-1}\{\bar{f}(s)\} = \mathcal{L}^{-1}\{\bar{g}(s)\bar{h}(s)\} = (g * h)(t)$$

where

$$(g * h)(t) = \int_0^t g(t-\tau)h(\tau)d\tau.$$

Now, suppose that $\bar{f}(s) = \bar{g}(s)\bar{h}(s)$, where $\bar{g}(s) = \frac{1}{s-a}$ and $\bar{h}(s) = \frac{s}{s^2+b^2}$.

From our table of Laplace transforms we know that $g(t) = \mathcal{L}^{-1}\left\{\frac{1}{s-a}\right\} = e^{at}$ and $h(t) = \mathcal{L}^{-1}\left\{\frac{s}{s^2+b^2}\right\} = \cos bt$.

Thus, by the Convolution Theorem, we have that

$$f(t) = \mathcal{L}^{-1}\{\bar{f}(s)\} = \mathcal{L}^{-1}\{\bar{g}(s)\bar{h}(s)\} = \int_0^t g(t-\tau)h(\tau)d\tau.$$

Therefore, using a computer algebra system, we see that

$$\begin{aligned}f(t) &= \int_0^t g(t-\tau)h(\tau)d\tau \\ &= \int_0^t e^{a(t-\tau)} \cos b\tau d\tau \\ &= e^{at} \int_0^t e^{-a\tau} \cos b\tau d\tau \\ &= \frac{1}{a^2+b^2} [ae^{at} - a\cos bt + b\sin bt].\end{aligned}$$

- iii. Heaviside's Expansion Theorem states that if $\bar{f}(s) = \frac{\bar{p}(s)}{\bar{q}(s)}$, where $\bar{p}(s)$ and $\bar{q}(s)$ are polynomials in s and the degree of \bar{q} is higher than that of \bar{p} , then

$$f(t) = \mathcal{L}^{-1} \{ \bar{f}(s) \} = \sum_{k=1}^n \frac{\bar{p}(\alpha_k)}{\bar{q}'(\alpha_k)} e^{t\alpha_k}$$

where α_k are the distinct root of $\bar{q}(s) = 0$.

For $\bar{f}(s) = \frac{s}{(s-a)(s^2+b^2)}$, we identify $\bar{p}(s) = s$ and $\bar{q}(s) = (s-a)(s^2+b^2)$. Since \bar{p} and \bar{q} are polynomials in s with the degree of \bar{q} greater than that of the degree of \bar{p} , the assumptions of Heaviside's Expansion Theorem are satisfied.

Note that $\bar{q}'(s) = s(3s-2a) + b^2$ and $\alpha_1 = a$, $\alpha_2 = bi$, and $\alpha_3 = -bi$ are the roots of $\bar{q}(s)$.

Therefore, by the Heaviside's Expansion Theorem, we have that

$$\begin{aligned} f(t) &= \mathcal{L}^{-1} \{ \bar{f}(s) \} = \sum_{k=1}^n \frac{\bar{p}(\alpha_k)}{\bar{q}'(\alpha_k)} e^{t\alpha_k} \\ &= \frac{a}{a^2+b^2} e^{at} - \frac{bi}{2bi(a-ib)} e^{bit} - \frac{bi}{2bi(a+ib)} e^{-bit} \\ &= \frac{1}{a^2+b^2} \left[ae^{at} - \frac{a+ib}{2} e^{bit} - \frac{a-ib}{2} e^{-bit} \right] \\ &= \frac{1}{a^2+b^2} [ae^{at} - a \cos bt + b \sin bt] . \end{aligned}$$

□

Problem 2. a. Evaluate the improper definite integral

$$\int_{-\infty}^{\infty} \frac{\cos tx}{x^2 + a^2} dx$$

where $a, t > 0$.

b. Show that

$$\int_0^{\infty} \frac{\sin \pi tx}{x(1+x^2)} dx = \frac{\pi}{2}(1 - e^{-\pi t})$$

where $t > 0$.

Solution. a. Suppose that

$$f(t) = \int_{-\infty}^{\infty} \frac{\cos tx}{x^2 + a^2} dx.$$

In order to evaluate this integral, we take the Laplace transform of $f(t)$ with respect to t . Now, due to uniform convergence, we have that

$$\begin{aligned} \bar{f}(s) = \mathcal{L}\{f(t)\} &= \mathcal{L}\left\{\int_{-\infty}^{\infty} \frac{\cos tx}{x^2 + a^2} dx\right\} = \int_{-\infty}^{\infty} \mathcal{L}\left\{\frac{\cos tx}{x^2 + a^2}\right\} dx \\ &= \int_{-\infty}^{\infty} \frac{1}{x^2 + a^2} \mathcal{L}\{\cos tx\} dx \\ &= \int_{-\infty}^{\infty} \frac{s}{(x^2 + a^2)(x^2 + s^2)} dx. \end{aligned}$$

Using the method of partial fraction decomposition, we see that this last integral becomes

$$\begin{aligned} \bar{f}(s) &= \int_{-\infty}^{\infty} \frac{s dx}{(x^2 + a^2)(x^2 + s^2)} \\ &= \frac{s}{s^2 - a^2} \int_{-\infty}^{\infty} \frac{1}{x^2 + a^2} - \frac{1}{x^2 + s^2} dx. \end{aligned}$$

Thus, we see that

$$\begin{aligned} \bar{f}(s) &= \frac{s}{s^2 - a^2} \int_{-\infty}^{\infty} \frac{1}{x^2 + a^2} - \frac{1}{x^2 + s^2} dx \\ &= \frac{s}{s^2 - a^2} \left[\tan^{-1} \frac{x}{a} \Big|_{-\infty}^{\infty} - \tan^{-1} \frac{x}{s} \Big|_{-\infty}^{\infty} \right] \\ &= \frac{s}{s^2 - a^2} \left[\frac{\pi}{a} - \frac{\pi}{s} \right] \\ &= \frac{\pi}{a} \left[\frac{s}{s^2 - a^2} - \frac{a}{s^2 - a^2} \right]. \end{aligned}$$

Using the table of Laplace transforms, we know that $\mathcal{L}^{-1} \left\{ \frac{s}{s^2 - a^2} \right\} = \cosh at$ and $\mathcal{L}^{-1} \left\{ \frac{a}{s^2 - a^2} \right\} = \sinh at$. Therefore, we have that

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{\cos tx}{x^2 + a^2} dx = f(t) &= \mathcal{L}^{-1} \{ \bar{f}(s) \} = \mathcal{L}^{-1} \left\{ \frac{\pi}{a} \left[\frac{s}{s^2 - a^2} - \frac{a}{s^2 - a^2} \right] \right\} \\ &= \frac{\pi}{a} \left[\mathcal{L}^{-1} \left\{ \frac{s}{s^2 - a^2} \right\} - \mathcal{L}^{-1} \left\{ \frac{a}{s^2 - a^2} \right\} \right] \\ &= \frac{\pi}{a} [\cosh at - \sinh at] \\ &= \frac{\pi}{a} e^{-at}. \end{aligned}$$

b. Suppose that

$$f(t) = \int_0^{\infty} \frac{\sin \pi tx}{x(1+x^2)} dx.$$

In order to evaluate this integral, we take the Laplace transform of $f(t)$ with respect to t . Now, due to uniform convergence, we have that

$$\begin{aligned} \bar{f}(s) = \mathcal{L} \{ f(t) \} &= \mathcal{L} \left\{ \int_0^{\infty} \frac{\sin \pi tx}{x(1+x^2)} dx \right\} = \int_0^{\infty} \mathcal{L} \left\{ \frac{\sin \pi tx}{x(1+x^2)} \right\} dx \\ &= \int_0^{\infty} \frac{1}{x(1+x^2)} \mathcal{L} \{ \sin \pi tx \} dx \\ &= \int_0^{\infty} \frac{\pi}{(x^2 + 1)(\pi^2 x^2 + s^2)} dx. \end{aligned}$$

Using a computer algebra system, we see that this last integral reduces to

$$\begin{aligned} \bar{f}(s) &= \int_0^{\infty} \frac{\pi}{(x^2 + 1)(\pi^2 x^2 + s^2)} dx \\ &= \frac{\pi^2}{2s(\pi + s)} \\ &= \frac{\pi}{2} \left[\frac{1}{s} - \frac{1}{s + \pi} \right]. \end{aligned}$$

Therefore, from our table of Laplace transforms, we have that

$$\begin{aligned} \int_0^{\infty} \frac{\sin \pi tx}{x(1+x^2)} dx = f(t) &= \mathcal{L}^{-1} \{ \bar{f}(s) \} = \frac{\pi}{2} \left[\mathcal{L}^{-1} \left\{ \frac{1}{s} \right\} - \mathcal{L}^{-1} \left\{ \frac{1}{s + \pi} \right\} \right] \\ &= \frac{\pi}{2} (1 - e^{-\pi t}). \end{aligned}$$

□

Problem 3. Apply the Laplace transform to solve the following Initial Value Problems:

- a. $y'' + 2ay' + (a^2 + 4)y = f(t)$
 $y(0) = 1, \quad y'(0) = -a.$
- b. $u_{tt} = c^2 u_{xx} + \sin x, \quad 0 < x < \pi, \quad t > 0$
 $u(0, t) = u(\pi, t) = 1, \quad u(x, 0) = u_t(x, 0) = 0.$

Solution. Recall that if $\bar{y}(s) = \mathcal{L}\{y(t)\}$, then the Laplace transform of the n -th derivative of $y(t)$ is given by

$$\mathcal{L}\{y^{(n)}(t)\} = s^n \bar{y}(s) - \sum_{k=0}^{n-1} s^{n-1-k} y^{(k)}(0). \quad (1)$$

- a. Suppose that $Ly \equiv y''(t) + 2ay'(t) + (a^2 + 4)y(t)$. Using (1), application of the Laplace transform to $Ly = f(t)$ yields that

$$\mathcal{L}\{Ly\} = (s^2 + 2as + a^2 + 4)\bar{y}(s) - 2ay(0) - sy(0) - y'(0) = \bar{f}(s) = \mathcal{L}\{f(t)\}.$$

From the initial data, we see that this reduces to

$$(s^2 + 2as + a^2 + 4)\bar{y}(s) - (s + a) = \bar{f}(s).$$

Solving for $\bar{y}(s)$ yields

$$\bar{y}(s) = \frac{\bar{f}(s) + s + a}{s^2 + 2as + a^2 + 4} = \frac{\bar{f}(s) + s + a}{(s + a + 2i)(s + a - 2i)}.$$

Note that from our table of Laplace transforms that

$$\mathcal{L}^{-1}\left\{\frac{a - b}{(s - a)(s - b)}\right\} = e^{at} - e^{bt}$$

and

$$\mathcal{L}^{-1}\left\{\frac{s}{(s - a)(s - b)}\right\} = \frac{ae^{at} - be^{bt}}{a - b}.$$

Therefore, the solution to the original differential equation is given by

$$\begin{aligned} y(t) &= \mathcal{L}^{-1}\{\bar{f}(s)\} \\ &= \mathcal{L}^{-1}\left\{\frac{\bar{f}(s) + s + a}{(s + a + 2i)(s + a - 2i)}\right\} \\ &= \mathcal{L}^{-1}\left\{\frac{\bar{f}(s)}{(s + a + 2i)(s + a - 2i)}\right\} + \frac{e^{-(2i+a)t}}{4} [(2 - i + ia)e^{4it} + 2 + i - ia]. \end{aligned}$$

- b. Let $u(x, t)$ be a function in x and t . The Laplace transform of $u(x, t)$ with respect to t is given by

$$\mathcal{L}\{u(x, t)\} = \bar{u}(x, s) = \int_0^\infty u(x, t)e^{-st}dt.$$

From this definition, we see from previous theorems that

$$\mathcal{L}\left\{\frac{\partial^n}{\partial t^n}[u(x, t)]\right\} = s^n \bar{u}(x, s) - \sum_{k=0}^{n-1} s^{n-1-k} \frac{\partial^k}{\partial t^k}[u(x, 0)]$$

Similarly, we see from the Leibniz integral rule that

$$\mathcal{L}\left\{\frac{\partial^n}{\partial x^n}[u(x, t)]\right\} = \frac{d^n}{dx^n}[\bar{u}(x, s)].$$

Applying the Laplace transform with respect to t to the differential equation yields that

$$\mathcal{L}\{u_{tt} - c^2 u_{xx}\} = s^2 \bar{u}(x, s) - su(x, 0) - u_t(x, 0) - c^2 \frac{d^2 \bar{u}(x, s)}{dx^2} = \frac{\sin x}{s} = \mathcal{L}\{\sin x\}.$$

In light of the initial data, this equation reduces to

$$s^2 \bar{u}(x, s) - c^2 \frac{d^2 \bar{u}(x, s)}{dx^2} = \frac{\sin x}{s},$$

or, equivalently,

$$\frac{d^2 \bar{u}(x, s)}{dx^2} - \left(\frac{s}{c}\right)^2 \bar{u}(x, s) = -\frac{\sin x}{sc^2}.$$

The homogeneous solution to the above differential equation is easily seen to be

$$\bar{u}_h(x, s) = c_1 \exp\left(-\frac{xs}{c}\right) + c_2 \exp\left(\frac{xs}{c}\right)$$

From the method of undetermined coefficients, assuming the particular solution of the equation is of the form $\bar{u}_p(x, s) = A \sin x$ for some unknown A , the particular solution of the transformed equation is given by

$$\bar{u}_p(x, s) = \frac{\sin x}{s(s^2 + c^2)}.$$

Therefore, the general solution to the transformed equation is given by

$$\bar{u}(x, s) = \bar{u}_h(x, s) + \bar{u}_p(x, s) = c_1 \exp\left(-\frac{xs}{c}\right) + c_2 \exp\left(\frac{xs}{c}\right) + \frac{\sin x}{s(s^2 + c^2)}.$$

Note that the transformed boundary data is given by $\bar{u}(0, s) = \bar{u}(\pi, s) = \frac{1}{s}$. Using the form of the solution to the transformed equation listed above, we see that in light of the transformed boundary data that

$$\begin{aligned} c_1 + c_2 &= \frac{1}{s} \\ c_1 \exp\left(-\frac{\pi s}{c}\right) + c_2 \exp\left(\frac{\pi s}{c}\right) &= \frac{1}{s} \end{aligned}$$

After solving the above system, we therefore see that the solution to the transformed equation is given by

$$\begin{aligned} \bar{u}(x, s) &= c_1 \exp\left(-\frac{xs}{c}\right) + c_2 \exp\left(\frac{xs}{c}\right) + \frac{\sin x}{s(s^2 + c^2)} \\ &= \frac{\exp\left(\frac{\pi s}{c}\right) \exp\left(-\frac{xs}{c}\right)}{s(1 + \exp\left(\frac{\pi s}{c}\right))} + \frac{\exp\left(\frac{xs}{c}\right)}{s(1 + \exp\left(\frac{\pi s}{c}\right))} + \frac{\sin x}{s(s^2 + c^2)} \end{aligned}$$

Therefore, the solution to the original differential equation is given by

$$u(x, t) = \mathcal{L}^{-1} \{ \bar{u}(x, s) \} = \mathcal{L}^{-1} \left\{ \frac{\exp\left(\frac{\pi s}{c}\right) \exp\left(-\frac{xs}{c}\right)}{s(1 + \exp\left(\frac{\pi s}{c}\right))} + \frac{\exp\left(\frac{xs}{c}\right)}{s(1 + \exp\left(\frac{\pi s}{c}\right))} + \frac{\sin x}{s(s^2 + c^2)} \right\}.$$

□

Problem 4.*Solution.*

□

Problem 5.*Solution.*

□

Problem 6.*Solution.*

□

Problem 7.*Solution.*

□

Problem 8.*Solution.*

□