

Exam 2

Matthew Tiger

December 11, 2016

Problem 1. A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is defined by $f(z) = z^8$. Find the fixed points of f . Use your calculations to find the real linear and quadratic factors of the polynomial $p(z) = z^7 - 1$.

Solution.

□

Problem 2. Let K_c be the filled-in Julia set of $f_c(z) = z^2 + c$.

- a. Find the fixed points and the period 2 points of f_{-6} .
- b. Show that $2\sqrt{2} \in K_{-6}$ and find another point in K_{-6} , distinct from those found so far.
- c. Do any of the points you have found lie in the Julia set of f_{-6} ?
- d. Is $-6 \in \mathcal{M}$ where \mathcal{M} is the Mandelbrot set?

Solution.

□

Problem 3. Let $f(z) = z^2 + c$. Find the values of c so that $z = i$ is a period 2 point. Find the fixed points in each case and determine their stability. Is $c \in \mathcal{M}$?

Solution.

□

Problem 4. Show that the function $H(z) = \frac{z-i}{z+i}$ gives a conjugacy between the Newton map N_{f_1} of $f_1(z) = z^2 + 1$ and the function $f_0(z) = z^2$. Deduce the Julia set of N_{f_1} and show that it is chaotic on its Julia set.

Solution.

□

Problem 5. Let $p(z)$ be a polynomial of degree $d > 1$ with Newton function

$$N_p(z) = z - \frac{p(z)}{p'(z)}.$$

- a. If $p(\alpha) = 0$ and $p'(\alpha) \neq 0$, show that α is a fixed point of multiplicity two for N_p , i.e. there is a rational function $k(z) = m(z)/n(z)$ with $n(\alpha) \neq 0$ and $N_p(z) - \alpha = (z - \alpha)^2 k(z)$.
- b. If $p(\alpha) = 0$, $p'(\alpha) \neq 0$, and $p''(\alpha) = 0$, show that α is a fixed point of multiplicity three for N_p .

Solution.

□

Problem 6. a. Show that for $p_\alpha(z) = z(z-1)(z-\alpha)$, the Newton function N_{p_α} has a critical point where $z = (\alpha+1)/3$.

b. For what values of α does p_α satisfy $p(\alpha) = 0$, $p'(\alpha) \neq 0$, and $p''(\alpha) = 0$?

Solution.

□

Problem 7. Let $0 < \mu < \lambda < 1$ and let $h : [0, 1] \rightarrow [0, 1]$ be a homeomorphism with $h \circ L_\mu(x) = L_\lambda \circ h(x)$ for all $x \in [0, 1]$.

- a. Show that h is orientation-preserving.
- b. Show that $h(x) + h(1 - x) = 1$ for all $x \in [0, 1]$. Deduce that $h(1/2) = 1/2$.
- c. Show that $h(\mu/4) = \lambda/4$ and $h(x) > x$ for $0 < x < 1/2$ and $h(x) < x$ for $1/2 < x < 1$.

Solution.

□

Problem 8. Prove that if $f_c(z) = z^2 + c$ has an attracting periodic point, then $c \in \mathcal{M}$, the Mandelbrot set.

Solution.

□