

# Conditioning and Error

## Unit 4

## Spectral radius and norms

For any subordinate matrix norm  $\rho(A) \leq \|A\|$

## Spectral radius and norms

For any subordinate matrix norm  $\rho(A) \leq \|A\|$

*Proof:*

$$Av = \lambda v \rightarrow$$

## Spectral radius and norms

For any subordinate matrix norm  $\rho(A) \leq \|A\|$

*Proof:*

$$Av = \lambda v \rightarrow \|\lambda v\| = \|Av\| \rightarrow$$

## Spectral radius and norms

For any subordinate matrix norm  $\rho(A) \leq \|A\|$

*Proof:*

$$Av = \lambda v \rightarrow \|\lambda v\| = \|Av\| \rightarrow |\lambda| \|v\| = \|\lambda v\| = \|Av\| \leq \|A\| \|v\|$$

Therefore,  $|\lambda| \leq \|A\|$ , and  $\rho(A) \leq \|A\|$ .

Question: why does the norm need to be subordinate?

## Spectral radius of symmetric matrix

For any Hermitian  $A$

$$\|A\|_2 = \rho(A), \quad \|A^{-1}\|_2 = 1/\mu, \quad \text{where } \mu := \min\{|\lambda|, \lambda \in \sigma(A)\}$$

# Spectral radius of symmetric matrix

For any Hermitian  $A$

$$\|A\|_2 = \rho(A), \quad \|A^{-1}\|_2 = 1/\mu, \quad \text{where } \mu := \min\{|\lambda|, \lambda \in \sigma(A)\}$$

*Proof:*

$$A = A^* \rightarrow A^2 = A^*A \rightarrow$$

# Spectral radius of symmetric matrix

For any Hermitian  $A$

$$\|A\|_2 = \rho(A), \quad \|A^{-1}\|_2 = 1/\mu, \quad \text{where } \mu := \min\{|\lambda|, \lambda \in \sigma(A)\}$$

*Proof:*

$$A = A^* \rightarrow A^2 = A^*A \rightarrow$$

$$\|A\|_2 = \sqrt{\rho(A^*A)} = \sqrt{\rho(A^2)} \stackrel{\text{why?}}{=} \sqrt{\rho(A)^2} = \rho(A)$$



## Spectral radius of symmetric matrix

For any Hermitian  $A$

$$\|A\|_2 = \rho(A), \quad \|A^{-1}\|_2 = 1/\mu, \quad \text{where } \mu := \min\{|\lambda|, \lambda \in \sigma(A)\}$$

*Proof:*

$$A = A^* \rightarrow A^2 = A^*A \rightarrow$$

$$\|A\|_2 = \sqrt{\rho(A^*A)} = \sqrt{\rho(A^2)} \stackrel{\text{why?}}{=} \sqrt{\rho(A)^2} = \rho(A)$$

$A^{-1}$  is also Hermitian with  $\rho(A^{-1}) = 1/\mu$ . Thus  $\|A^{-1}\|_2 = 1/\mu$ .

## Comparing norms

- $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

## Comparing norms

- $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

*Proof:*

$$\lambda := \|A\|_2^2 \quad \rightarrow \quad A^* A v = \lambda v \quad \rightarrow$$

## Comparing norms

- $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

*Proof:*

$$\lambda := \|A\|_2^2 \quad \rightarrow \quad A^* A v = \lambda v \quad \rightarrow \quad \|\lambda v\|_1 = \|A^* A v\|_1 \quad \rightarrow$$

## Comparing norms

- $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

*Proof:*

$$\lambda := \|A\|_2^2 \quad \rightarrow \quad A^* A v = \lambda v \quad \rightarrow \quad \|\lambda v\|_1 = \|A^* A v\|_1 \quad \rightarrow$$

$$\lambda \|v\|_1 \leq \|A^*\|_1 \|A\|_1 \|v\|_1 \quad \rightarrow$$

## Comparing norms

- $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

*Proof:*

$$\lambda := \|A\|_2^2 \rightarrow A^* A v = \lambda v \rightarrow \|\lambda v\|_1 = \|A^* A v\|_1 \rightarrow$$

$$\lambda \|v\|_1 \leq \|A^*\|_1 \|A\|_1 \|v\|_1 \rightarrow \|A\|_2^2 \|v\|_1 \leq \|A^*\|_1 \|A\|_1 \|v\|_1$$

Since  $\|A^*\|_1 = \|A\|_\infty$ ,

## Comparing norms

- $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

*Proof:*

$$\lambda := \|A\|_2^2 \rightarrow A^* A v = \lambda v \rightarrow \|\lambda v\|_1 = \|A^* A v\|_1 \rightarrow$$

$$\lambda \|v\|_1 \leq \|A^*\|_1 \|A\|_1 \|v\|_1 \rightarrow \|A\|_2^2 \|v\|_1 \leq \|A^*\|_1 \|A\|_1 \|v\|_1$$

Since  $\|A^*\|_1 = \|A\|_\infty$ , we have  $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

## Comparing norms

- $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

*Proof:*

$$\lambda := \|A\|_2^2 \rightarrow A^* A v = \lambda v \rightarrow \|\lambda v\|_1 = \|A^* A v\|_1 \rightarrow$$

$$\lambda \|v\|_1 \leq \|A^*\|_1 \|A\|_1 \|v\|_1 \rightarrow \|A\|_2^2 \|v\|_1 \leq \|A^*\|_1 \|A\|_1 \|v\|_1$$

Since  $\|A^*\|_1 = \|A\|_\infty$ , we have  $\|A\|_2^2 \leq \|A\|_1 \|A\|_\infty$

- $\|A\|_2 \leq \|A\|_F \leq \sqrt{n} \|A\|_2$  *without proof*



## Basic facts

- The condition number  $\text{cond}(A) = \|A\| \|A^{-1}\|$  depends on the matrix  $A$  and on the norm used. If  $\text{cond}(A)$  is large,  $A$  is called ill-conditioned.

## Basic facts

- The condition number  $\text{cond}(A) = \|A\| \|A^{-1}\|$  depends on the matrix  $A$  and on the norm used. If  $\text{cond}(A)$  is large,  $A$  is called ill-conditioned.
- Since  $\|A\| \|A^{-1}\| \geq \|AA^{-1}\| = \|I\| = 1$ , we always have

$$\text{cond}(A) \geq 1$$

## Basic facts

- The condition number  $\text{cond}(A) = \|A\| \|A^{-1}\|$  depends on the matrix  $A$  and on the norm used. If  $\text{cond}(A)$  is large,  $A$  is called ill-conditioned.
- Since  $\|A\| \|A^{-1}\| \geq \|AA^{-1}\| = \|I\| = 1$ , we always have

$$\text{cond}(A) \geq 1$$

- Since all matrix norms are equivalent, the dependence of  $\text{cond}(A)$  on the norm chosen is less important than the dependence on  $A$ .

## Basic facts

- The condition number  $\text{cond}(A) = \|A\| \|A^{-1}\|$  depends on the matrix  $A$  and on the norm used. If  $\text{cond}(A)$  is large,  $A$  is called ill-conditioned.
- Since  $\|A\| \|A^{-1}\| \geq \|AA^{-1}\| = \|I\| = 1$ , we always have

$$\text{cond}(A) \geq 1$$

- Since all matrix norms are equivalent, the dependence of  $\text{cond}(A)$  on the norm chosen is less important than the dependence on  $A$ .
- The spectral norm is usually the choice for analyzing properties of the condition number. The 1- and  $\infty$ - norms are used in computations.

## Examples

- Almost singular

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 - 10^{-6} \end{pmatrix}$$

$$\text{cond}_2(A) \approx \text{cond}_1(A) \approx \text{cond}_\infty(A) \approx 4 \times 10^6$$

## Examples

- Almost singular

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 - 10^{-6} \end{pmatrix}$$

$$\text{cond}_2(A) \approx \text{cond}_1(A) \approx \text{cond}_\infty(A) \approx 4 \times 10^6$$

- Hilbert matrix  $H(n)$ ,  $h_{ij} := \frac{1}{i+j+1}$ .

$$H(3) = \begin{pmatrix} 1 & 1/2 & 1/3 \\ 1/2 & 1/3 & 1/4 \\ 1/3 & 1/4 & 1/5 \end{pmatrix}$$

$$\text{cond}_2(H(10)) \approx 10^{13}, \quad \text{cond}_1(A) \approx \text{cond}_\infty(A) \approx 3 \times 10^{13}$$

**HW: produce a picture similar to Fig. 4.1 for Hilbert matrix**

## Linear systems, example

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 - 10^{-5}$$

We can find the exact analytic solution easily  $x_1 = x_2 = 10$

## Linear systems, example

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 - 10^{-5}$$

We can find the exact analytic solution easily  $x_1 = x_2 = 10$

A small change now

$$x_1 + x_2 = 20$$

$$x_1 + (1 + 10^{-6})x_2 = 20 - 10^{-5}$$



## Linear systems, example

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 - 10^{-5}$$

We can find the exact analytic solution easily  $x_1 = x_2 = 10$

A small change now

$$x_1 + x_2 = 20$$

$$x_1 + (1 + 10^{-6})x_2 = 20 - 10^{-5}$$

We can still find the exact analytic solution easily

$$x_1 = 30, \quad x_2 = -10.$$

But ... a very small change in one coefficient seems to yield a significant change in the solution. Can we quantify this?

## Linear systems, example, continued

But ... a very small change in one coefficient seems to yield a significant change in the solution. Can we quantify this?

- a small change in  $A$  would be ....

## Linear systems, example, continued

But ... a very small change in one coefficient seems to yield a significant change in the solution. Can we quantify this?

- a small change in  $A$  would be ....  $\frac{\|\delta A\|}{\|A\|} = \frac{2 \times 10^{-6}}{2 \times 10^{-6}} \approx 10^{-6}$

## Linear systems, example, continued

But ... a very small change in one coefficient seems to yield a significant change in the solution. Can we quantify this?

- a small change in  $A$  would be ....  $\frac{\|\delta A\|}{\|A\|} = \frac{2 \times 10^{-6}}{2 \times 10^{-6}} \approx 10^{-6}$
- a significant change in the solution would be  $\frac{\|\delta x\|}{\|x\|} = \frac{20}{10} = 2$
- We have  $\frac{\|\delta x\|}{\|x\|} \approx 2 \times 10^6 \frac{\|\delta A\|}{\|A\|} \approx$

## Linear systems, example, continued

But ... a very small change in one coefficient seems to yield a significant change in the solution. Can we quantify this?

- a small change in  $A$  would be ....  $\frac{\|\delta A\|}{\|A\|} = \frac{2 \times 10^{-6}}{2 - 10^{-6}} \approx 10^{-6}$
- a significant change in the solution would be  $\frac{\|\delta x\|}{\|x\|} = \frac{20}{10} = 2$
- We have  $\frac{\|\delta x\|}{\|x\|} \approx 2 \times 10^6 \frac{\|\delta A\|}{\|A\|} \approx 0.5 \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$

In general, we have  $\frac{\|\delta x\|}{\|x\|} \leq 2 \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$  or...

## Linear systems, example, continued

But ... a very small change in one coefficient seems to yield a significant change in the solution. Can we quantify this?

- a small change in  $A$  would be ....  $\frac{\|\delta A\|}{\|A\|} = \frac{2 \times 10^{-6}}{2 - 10^{-6}} \approx 10^{-6}$
- a significant change in the solution would be  $\frac{\|\delta x\|}{\|x\|} = \frac{20}{10} = 2$
- We have  $\frac{\|\delta x\|}{\|x\|} \approx 2 \times 10^6 \frac{\|\delta A\|}{\|A\|} \approx 0.5 \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$

In general, we have  $\frac{\|\delta x\|}{\|x\|} \leq 2 \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$  or...

$$\frac{\|\delta x\|}{\|x + \delta x\|} \leq \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

## Linear systems, example, continued

But ... a very small change in one coefficient seems to yield a significant change in the solution. Can we quantify this?

- a small change in  $A$  would be ....  $\frac{\|\delta A\|}{\|A\|} = \frac{2 \times 10^{-6}}{2 - 10^{-6}} \approx 10^{-6}$
- a significant change in the solution would be  $\frac{\|\delta x\|}{\|x\|} = \frac{20}{10} = 2$
- We have  $\frac{\|\delta x\|}{\|x\|} \approx 2 \times 10^6 \frac{\|\delta A\|}{\|A\|} \approx 0.5 \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$

In general, we have  $\frac{\|\delta x\|}{\|x\|} \leq 2 \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$  or...

$$\frac{\|\delta x\|}{\|x + \delta x\|} \leq \text{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

## Main theorem I

If  $A$  and  $A + \delta A$  are both invertible, then

$$\|\delta x\| < \|x\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x\|} \leq 2 \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|},$$

$$\text{and} \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

*Proof:* Subtract  $b = Ax$  from  $(A + \delta A)(x + \delta x) = b$  to obtain



## Main theorem I

If  $A$  and  $A + \delta A$  are both invertible, then

$$\|\delta x\| < \|x\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x\|} \leq 2 \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|},$$

$$\text{and} \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

*Proof:* Subtract  $b = Ax$  from  $(A + \delta A)(x + \delta x) = b$  to obtain

$$\delta A(x + \delta x) + A\delta x = 0 \quad \rightarrow$$

## Main theorem I

If  $A$  and  $A + \delta A$  are both invertible, then

$$\|\delta x\| < \|x\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x\|} \leq 2 \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|},$$

$$\text{and} \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

*Proof:* Subtract  $b = Ax$  from  $(A + \delta A)(x + \delta x) = b$  to obtain

$$\delta A(x + \delta x) + A\delta x = 0 \quad \rightarrow \quad -\delta x = A^{-1}(\delta A)(x + \delta x).$$

## Main theorem I

If  $A$  and  $A + \delta A$  are both invertible, then

$$\|\delta x\| < \|x\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x\|} \leq 2 \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|},$$

$$\text{and} \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

*Proof:* Subtract  $b = Ax$  from  $(A + \delta A)(x + \delta x) = b$  to obtain

$$\delta A(x + \delta x) + A\delta x = 0 \quad \rightarrow \quad -\delta x = A^{-1}(\delta A)(x + \delta x).$$

$$\|\delta x\| = \|A^{-1}(\delta A)(x + \delta x)\| \quad \rightarrow$$

## Main theorem I

If  $A$  and  $A + \delta A$  are both invertible, then

$$\|\delta x\| < \|x\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x\|} \leq 2 \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|},$$

$$\text{and} \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

*Proof:* Subtract  $b = Ax$  from  $(A + \delta A)(x + \delta x) = b$  to obtain

$$\delta A(x + \delta x) + A\delta x = 0 \quad \rightarrow \quad -\delta x = A^{-1}(\delta A)(x + \delta x).$$

$$\|\delta x\| = \|A^{-1}(\delta A)(x + \delta x)\| \quad \rightarrow \quad \|\delta x\| \leq \|A^{-1}\| \|\delta A\| \|x + \delta x\|.$$

## Main theorem I

If  $A$  and  $A + \delta A$  are both invertible, then

$$\|\delta x\| < \|x\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x\|} \leq 2 \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|},$$

$$\text{and} \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

*Proof:* Subtract  $b = Ax$  from  $(A + \delta A)(x + \delta x) = b$  to obtain

$$\delta A(x + \delta x) + A\delta x = 0 \quad \rightarrow \quad -\delta x = A^{-1}(\delta A)(x + \delta x).$$

$$\|\delta x\| = \|A^{-1}(\delta A)(x + \delta x)\| \quad \rightarrow \quad \|\delta x\| \leq \|A^{-1}\| \|\delta A\| \|x + \delta x\|.$$

$$\frac{\|\delta x\|}{\|x + \delta x\|} \leq \|A^{-1}\| \|\delta A\| \quad \rightarrow$$

## Main theorem I

If  $A$  and  $A + \delta A$  are both invertible, then

$$\|\delta x\| < \|x\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x\|} \leq 2 \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|},$$

$$\text{and} \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \operatorname{cond}(A) \frac{\|\delta A\|}{\|A\|}$$

*Proof:* Subtract  $b = Ax$  from  $(A + \delta A)(x + \delta x) = b$  to obtain

$$\delta A(x + \delta x) + A\delta x = 0 \quad \rightarrow \quad -\delta x = A^{-1}(\delta A)(x + \delta x).$$

$$\|\delta x\| = \|A^{-1}(\delta A)(x + \delta x)\| \quad \rightarrow \quad \|\delta x\| \leq \|A^{-1}\| \|\delta A\| \|x + \delta x\|.$$

$$\frac{\|\delta x\|}{\|x + \delta x\|} \leq \|A^{-1}\| \|\delta A\| \quad \rightarrow \quad \frac{\|\delta x\|}{\|x + \delta x\|} \leq \|A^{-1}\| \|A\| \frac{\|\delta A\|}{\|A\|}$$

and the second inequality follows.

**HW: prove the first inequality**

## Linear systems, example

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 - 10^{-5}$$

We can find the exact analytic solution easily  $x_1 = x_2 = 10$

## Linear systems, example

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 - 10^{-5}$$

We can find the exact analytic solution easily  $x_1 = x_2 = 10$

A small change now

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 + 10^{-5}$$



## Linear systems, example

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 - 10^{-5}$$

We can find the exact analytic solution easily  $x_1 = x_2 = 10$

A small change now

$$x_1 + x_2 = 20$$

$$x_1 + (1 - 10^{-6})x_2 = 20 + 10^{-5}$$

We can still find the exact analytic solution easily

$$x_1 = 30, \quad x_2 = -10.$$

Once again, a very small change in  $b$  seems to yield a significant change in the solution. Can we quantify this?

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow$

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow \|b\| \leq \|A\| \|x\|.$

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow \|b\| \leq \|A\| \|x\|.$

On the other hand,

$$A(x + \delta x) = b + \delta b \rightarrow A\delta x = \delta b \rightarrow$$

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow \|b\| \leq \|A\| \|x\|.$

On the other hand,

$$A(x + \delta x) = b + \delta b \rightarrow A\delta x = \delta b \rightarrow \delta x = A^{-1}\delta b.$$

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow \|b\| \leq \|A\| \|x\|.$

On the other hand,

$$A(x + \delta x) = b + \delta b \rightarrow A\delta x = \delta b \rightarrow \delta x = A^{-1}\delta b.$$

Then,  $\|\delta x\| = \|A^{-1}\delta b\| \rightarrow$

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow \|b\| \leq \|A\| \|x\|.$

On the other hand,

$$A(x + \delta x) = b + \delta b \rightarrow A\delta x = \delta b \rightarrow \delta x = A^{-1}\delta b.$$

$$\text{Then, } \|\delta x\| = \|A^{-1}\delta b\| \rightarrow \|\delta x\| \leq \|A^{-1}\| \|\delta b\|.$$



## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow \|b\| \leq \|A\|\|x\|.$

On the other hand,

$$A(x + \delta x) = b + \delta b \rightarrow A\delta x = \delta b \rightarrow \delta x = A^{-1}\delta b.$$

$$\text{Then, } \|\delta x\| = \|A^{-1}\delta b\| \rightarrow \|\delta x\| \leq \|A^{-1}\|\|\delta b\|.$$

Multiplying the two inequalities, we have

$$\|b\|\|\delta x\| \leq \|A\|\|A^{-1}\|\|x\|\|\delta b\| \rightarrow$$

## Main theorem II

If  $A$  is invertible, then

$$\frac{\|\delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\delta b\|}{\|b\|}$$

*Proof:*  $b = Ax \rightarrow \|b\| = \|Ax\| \rightarrow \|b\| \leq \|A\|\|x\|.$

On the other hand,

$$A(x + \delta x) = b + \delta b \rightarrow A\delta x = \delta b \rightarrow \delta x = A^{-1}\delta b.$$

Then,  $\|\delta x\| = \|A^{-1}\delta b\| \rightarrow \|\delta x\| \leq \|A^{-1}\|\|\delta b\|.$

Multiplying the two inequalities, we have

$$\|b\|\|\delta x\| \leq \|A\|\|A^{-1}\|\|x\|\|\delta b\| \rightarrow \|b\|\|\delta x\| \leq \text{cond}(A)\|x\|\|\delta b\|$$