

Homework Assignment 2

Matthew Tiger

February 22, 2016

Problem 1. Use the method of variation of parameters to find the general solution of

$$y'' + 2y' + 2y = \sin x.$$

Solution. Suppose that $Ly = y'' + 2y' + 2y$. The general solution to $Ly = \sin x$ is given by $y = y_0 + y_h$ where y_0 is a particular solution of $Ly = \sin x$ and y_h is the solution to the homogeneous equation $Ly = 0$.

The characteristic equation of the equation $Ly = 0$ is $m(x) = x^2 + 2x + 2$, the roots of which are $m_1 = -1 - i$ and $m_2 = -1 + i$. As the roots of the characteristic equation are complex, the solution to $Ly = 0$ is given by

$$y_h = c_1 e^{-x} \sin x + c_2 e^{-x} \cos x. \quad (1)$$

The method of variation of parameters can be used to find a particular solution y_0 . We wish to find functions $u_1(x), u_2(x)$ such that

$$y_0 = u_1(x)y_1(x) + u_2(x)y_2(x) \quad (2)$$

satisfies $Ly_0 = \sin x$ where $y_1(x)$ and $y_2(x)$ are solutions to the homogeneous equation $Ly = 0$. If the functions $u_1(x)$ and $u_2(x)$ are solutions to the system

$$\begin{cases} u_1' y_1 + u_2' y_2 = 0 \\ u_1' y_1' + u_2' y_2' = \sin x \end{cases} \quad (3)$$

then (2) will satisfy the original differential equation $Ly = \sin x$ equation. The solution to the system (3) is

$$u_1(x) = - \int \frac{y_2(x) \sin x}{W[\{y_1, y_2\}]} dx \quad u_2(x) = \int \frac{y_1(x) \sin x}{W[\{y_1, y_2\}]} dx \quad (4)$$

where $W[\{y_1, y_2\}]$ is the Wronskian of the functions y_1 and y_2 .

Using (1), we know that $y_1(x) = e^{-x} \sin x$ and $y_2(x) = e^{-x} \cos x$ so the particular solution has the form $y_0 = u_1(x)e^{-x} \sin x + u_2(x)e^{-x} \cos x$. Further, the Wronskian of y_1 and y_2 is

$$W[\{y_1, y_2\}] = \begin{vmatrix} e^{-x} \sin x & e^{-x} \cos x \\ e^{-x} \cos x - e^{-x} \sin x & -e^{-x} \cos x - e^{-x} \sin x \end{vmatrix} = -e^{-2x}.$$

Thus, using (4), we know that

$$\begin{aligned} u_1(x) &= - \int \frac{y_2(x) \sin x}{W[\{y_1, y_2\}]} dx \\ &= \int \frac{e^{-x} \cos x \sin x}{e^{-2x}} dx \\ &= \frac{e^x}{10} (-2 \cos 2x + \sin 2x) + C \end{aligned}$$

and

$$\begin{aligned} u_2(x) &= \int \frac{y_1(x) \sin x}{W[\{y_1, y_2\}]} dx \\ &= - \int \frac{e^{-x} \sin^2 x}{e^{-2x}} dx \\ &= \frac{e^x}{10} (-5 + \cos 2x + 2 \sin 2x) + C. \end{aligned}$$

Therefore, a particular solution to $Ly = \sin x$ is

$$y_0(x) = \frac{1}{10} (-2 \cos 2x + \sin 2x) \sin x + \frac{1}{10} (-5 + \cos 2x + 2 \sin 2x) \cos x$$

and the general solution to $Ly = \sin x$ is

$$\begin{aligned} y(x) &= y_0(x) + y_h(x) \\ &= \frac{1}{10} (-2 \cos 2x + \sin 2x) \sin x + \frac{1}{10} (-5 + \cos 2x + 2 \sin 2x) \cos x \\ &\quad + c_1 e^{-x} \sin x + c_2 e^{-x} \cos x \end{aligned} \tag{5}$$

□

Problem 2. Find the Green function of the IVP

$$y'' + 2y' + 2y = f(x), \quad y(0) = y'(0) = 0.$$

Solution. Let $Ly = f(x)$ denote the differential equation $y'' + 2y' + 2y = f(x)$ together with the initial conditions $y(0) = y'(0) = 0$. The Green function $G(x, a)$ of the IVP $Ly = f(x)$ is defined by the equations

$$\frac{\partial^2 G(x, a)}{\partial x^2} + \frac{2\partial G(x, a)}{\partial x} + 2G(x, a) = \delta(x - a), \quad G(0, a) = 0, \quad \frac{\partial G}{\partial x}(0, a) = 0$$

where $\delta(x - a)$ is the Dirac Delta function such that $\int_{-\infty}^{\infty} \delta(x - a)f(x)dx = f(a)$. Note that $G(x, a)$ is continuous at $x = a$ and $\partial G/\partial x$ has a jump discontinuity of magnitude 1 at $x = a$.

If y_1 and y_2 are linearly independent solutions of the homogeneous equation $Ly = 0$, then

$$G(x, a) = \begin{cases} A_1 y_1 + A_2 y_2 & \text{if } x < a \\ B_1 y_1 + B_2 y_2 & \text{if } x > a \end{cases}$$

where A_1, A_2, B_1 , and B_2 are undetermined functions. The continuity of $G(x, a)$ at $x = a$ gives the equation

$$A_1 y_1(a) + A_2 y_2(a) = B_1 y_1(a) + B_2 y_2(a).$$

Further, the fact that $\partial G/\partial x$ has a jump discontinuity of magnitude 1 at $x = a$ yields the second equation

$$(B_1 y_1'(a) + B_2 y_2'(a)) - (A_1 y_1'(a) + A_2 y_2'(a)) = 1.$$

Combining these equations, we see that A_1, A_2, B_1 , and B_2 are given by

$$B_1 = A_1 - \frac{y_2(a)}{W[y_1(a), y_2(a)]}$$

$$B_2 = A_2 + \frac{y_1(a)}{W[y_1(a), y_2(a)]}$$

From (1), we know that the linearly independent solutions to the homogeneous equation $Ly = 0$ are $y_1(x) = e^{-x} \sin x$ and $y_2(x) = e^{-x} \cos x$. Also, the Wronskian of these solutions is $W[y_1(a), y_2(a)] = -e^{-2a}$. Thus,

$$B_1 = A_1 - \frac{y_2(a)}{W[y_1(a), y_2(a)]} = A_1 + e^a \cos a$$

$$B_2 = A_2 + \frac{y_1(a)}{W[y_1(a), y_2(a)]} = A_2 - e^a \sin a$$

Using the two initial conditions, we can uniquely determine A_1 and A_2 since $G(x, a) = A_1 y_1(a) + A_2 y_2(a)$ satisfies $LG = f(x)$. Since $y(0) = 0$ we see that $A_2 = 0$ and since $y'(0) = 0$ we see that $A_1 - A_2 = 0$ implying that $A_1 = A_2 = 0$. Therefore, the Green function for the IVP $Ly = f(x)$ is

$$G(x, a) = \begin{cases} 0 & \text{if } x < a \\ e^{a-x} (\sin x \cos a - \sin a \cos x) & \text{if } x > a \end{cases} \quad (6)$$

□

Problem 3. Use your answer to Problem 2 to solve the IVP

$$y'' + 2y' + 2y = \sin x, \quad y(0) = y'(0) = 0.$$

Solution.

□

Problem 4. Show that if y_1 , y_2 , and y_3 are three linearly independent solutions of the linear ODE

$$y''' + p_2(x)y'' + p_1(x)y' + p_0(x)y = 0$$

and u_1 , u_2 , u_3 are solutions of the system

$$\begin{cases} u_1'y_1 + u_2'y_2 + u_3'y_3 = 0, \\ u_1'y_1' + u_2'y_2' + u_3'y_3' = 0, \\ u_1'y_1'' + u_2'y_2'' + u_3'y_3'' = f(x), \end{cases}$$

then the function $u = u_1y_1 + u_2y_2 + u_3y_3$ is a solution of

$$y''' + p_2(x)y'' + p_1(x)y' + p_0(x)y = f(x)$$

Solution.

□

Problem 5. Find the eigenvalues and the respective eigenfunctions for the BVP

$$x^2 y'' + xy' + \lambda y = 0, \quad y'(1) = 0, \quad y'(b) = 0$$

where $b > 1$.

Solution.

□