

Homework Assignment 1

Matthew Tiger

February 8, 2016

Problem 1. Solve the IVP:

$$y' = y^2 \cos(x), \quad y(0) = 2.$$

Solution. Note that this is a separable differential equation and after separating we see that

$$\begin{aligned} \frac{dy}{y^2} &= \cos(x) dx \\ \int \frac{dy}{y^2} &= \int \cos(x) dx \\ -\frac{1}{y} &= \sin(x) + c_1 \end{aligned}$$

so that

$$y = -\frac{1}{\sin(x) + c_1}$$

is the general solution to the differential equation. Using the initial value $y(0) = 2$ and solving for c_1 we see that $c_1 = -1/2$ and the solution to the IVP is given by

$$y = -\frac{1}{\sin(x) - 1/2}.$$

□

Problem 2. Review solutions of first-order linear ODEs (p. 14) and solve the IVP:

$$y' - xy = x^3, \quad y(1) = \frac{1}{2}.$$

Solution. The solution to the first-order linear ODE

$$y'(x) + p_0(x)y(x) = f(x)$$

is given by

$$y(x) = \frac{c_1}{I(x)} + \frac{1}{I(x)} \int_0^x f(t)I(t)dt, \quad I(x) = \exp \left(\int_0^x p_0(t)dt \right).$$

For this problem, we set $p_0(x) = -x$ and $f(x) = x^3$ and see that

$$I(x) = \exp \left(\int_0^x p_0(t)dt \right) = \exp \left(\int_0^x -tdt \right) = \exp \left(-\frac{x^2}{2} \right).$$

Thus the general solution to the ODE $y' - xy = x^3$ is given by

$$\begin{aligned} y &= \frac{c_1}{\exp \left(-\frac{x^2}{2} \right)} + \frac{1}{\exp \left(-\frac{x^2}{2} \right)} \int_0^x t^3 \exp \left(-\frac{t^2}{2} \right) dt \\ &= \frac{c_1}{\exp \left(-\frac{x^2}{2} \right)} - \frac{\exp \left(-\frac{x^2}{2} \right)}{\exp \left(-\frac{x^2}{2} \right)} (2 + x^2) \\ &= \frac{c_1}{\exp \left(-\frac{x^2}{2} \right)} - (2 + x^2) \end{aligned}$$

Using the initial value $y(1) = \frac{1}{2}$, we see that $c_1 = \frac{7}{2} \exp \left(-\frac{1}{2} \right)$ and the solution to the IVP is

$$y = \frac{7 \exp \left(-\frac{1}{2} \right)}{2 \exp \left(-\frac{x^2}{2} \right)} - (2 + x^2).$$

□

Problem 3. Let $Ly = y^{(4)} - 4y''' + 3y'' + 4y' - 4y$.

a. Find the general solutions of the homogeneous ODE $Ly = 0$.

b. Solve the IVP:

$$Ly = 0, \quad y(0) = 0, \quad y'(0) = -7, \quad y''(0) = 5, \quad y'''(0) = 9.$$

c. Solve the BVP:

$$Ly = 0, \quad y(0) = 1, \quad \lim_{x \rightarrow \infty} y(x) = 0.$$

Is this BVP well-posed?

d. Solve the BVP:

$$Ly = 0, \quad y(0) = 1, \quad \lim_{x \rightarrow -\infty} y(x) = 0.$$

Is this BVP well-posed?

Solution. a. The characteristic equation associated to the homogeneous ODE $Ly = 0$ is $m(x) = x^4 - 4x^3 + 3x^2 + 4x - 4$. The roots of the characteristic polynomial are $r_1 = -1$, $r_2 = 1$, $r_3 = 2$, and $r_4 = 2$.

Therefore, the general solution of the homogeneous ODE is

$$y(x) = c_1 e^{-x} + c_2 e^x + c_3 e^{2x} + c_4 x e^{2x}. \quad (1)$$

b. Through an abuse of notation, we note that the matrix associated to the Wronskian of this equation as function of x is given by

$$W(x) = \begin{bmatrix} e^{-x} & e^x & e^{2x} & x e^{2x} \\ -e^{-x} & e^x & 2e^{2x} & e^{2x} + 2x e^{2x} \\ e^{-x} & e^x & 4e^{2x} & 4e^{2x} + 4x e^{2x} \\ -e^{-x} & e^x & 8e^{2x} & 12e^{2x} + 8x e^{2x} \end{bmatrix}.$$

The solution to the IVP is determined by particular values of the coefficients in the general solution (1). These coefficients are found as the solution to the system of equations $W(0)\mathbf{c} = \mathbf{b}$ where

$$\mathbf{c} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ c_4 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} 0 \\ -7 \\ 5 \\ 9 \end{bmatrix}.$$

The solution to this system is given by $\mathbf{c} = \langle 4, -3, -1, 2 \rangle$. Therefore, the solution to the IVP is

$$y(x) = 4e^{-x} - 3e^x - e^{2x} + 2x e^{2x}.$$

c.

d.

□

Problem 4. Read §1.6 and then solve the ODEs:

$$xy' + 2y = x^2\sqrt{y}, \quad y' = \frac{4x^3 - 6xy^2 - 2xy}{x^2 + 6x^2y - 3y^2}, \quad y' + y^2 + (2x + 1)y + 1 + x + x^2 = 0.$$

Solution.

□

Problem 5. a. Use mathematical induction to prove Leibnitz's differentiation rule:

$$D^k(fg) = \sum_{j=0}^k \binom{k}{j} (D^j f)(D^{k-j} g).$$

Here $f = f(x)$ and $g = g(x)$ are k -times differentiable functions and $D^k = \frac{d^k}{dx^k}$.

b. Consider the constant-coefficient ODE

$$D^n y + p_{n-1} D^{n-1} y + \cdots + p_1 D y + p_0 y = 0, \quad (2)$$

where p_0, p_1, \dots, p_{n-1} are real numbers. Let r be a double root of the characteristic polynomial $P(z) = z^n + p_{n-1} z^{n-1} + \cdots + p_1 z + p_0$. Use Leibnitz's rule to show that the function $x e^{rx}$ is a solution of (2).

c. Let r be a triple root of the characteristic polynomial $P(z)$ from part (b). Use Leibnitz's rule to show that the function $x^2 e^{rx}$ is then also a solution of (2).

d. Let r be a real number. Show that the functions e^{rx} , $x e^{rx}$, and $x^2 e^{rx}$ are linearly independent on \mathbb{R} .

Solution.

□

Problem 6. Use the formula for the derivative of a determinant from the lectures, other properties of determinants, and the linear ODE (1.3.1) to verify identity (1.3.4) in the text-book.

Solution.

□