

# Exam 1

Matthew Tiger

March 12, 2017

**Problem 1.** Find the Fourier Transforms of the following functions:

- a.  $f(x) = x^2 e^{-a|x|}$ ,  $a > 0$ ,
- b.  $f(x) = \left(1 - \frac{|x|}{2}\right) H\left(1 - \frac{|x|}{2}\right)$ .

*Solution.* Recall that if  $f(x) \in L^1(\mathbb{R})$ , then the Fourier Transform of  $f$  is defined to be

$$\mathcal{F}\{f(x)\} = F(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ikx} dx. \quad (1)$$

- a. If  $f(x) = x^2 e^{-a|x|}$ ,  $a > 0$ , we see from (1) that, by definition, the Fourier Transform of  $f$  is given by

$$\begin{aligned} \mathcal{F}\{f(x)\} = F(k) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x^2 e^{-a|x|} e^{-ikx} dx \\ &= \frac{1}{\sqrt{2\pi}} \left[ \int_{-\infty}^0 x^2 e^{-(-a+ik)x} dx + \int_0^{\infty} x^2 e^{-(a+ik)x} dx \right]. \end{aligned}$$

Now, we see by integration by parts that

$$\begin{aligned} \int_c^d x^2 e^{-(b+ik)x} dx &= -\frac{x^2}{b+ik} e^{-(b+ik)x} \Big|_c^d + \frac{2}{b+ik} \int_c^d x e^{-(b+ik)x} dx \\ &= -\frac{x^2}{b+ik} e^{-(b+ik)x} \Big|_c^d + \frac{2}{b+ik} \left[ -\frac{x}{b+ik} e^{-(b+ik)x} \Big|_c^d + \frac{1}{b+ik} \int_c^d e^{-(b+ik)x} dx \right] \\ &= -\frac{x^2}{b+ik} e^{-(b+ik)x} \Big|_c^d + \frac{2}{b+ik} \left[ -\frac{x}{b+ik} e^{-(b+ik)x} \Big|_c^d - \frac{1}{(b+ik)^2} e^{-(b+ik)x} \Big|_c^d \right]. \end{aligned}$$

Thus,

$$\int_{-\infty}^0 x^2 e^{-(-a+ik)x} dx = -\frac{2}{(-a+ik)^3}$$

and

$$\int_0^\infty x^2 e^{-(a+ik)x} dx = \frac{2}{(a+ik)^3}.$$

Therefore,

$$\begin{aligned} \mathcal{F}\{f(x)\} = F(k) &= \frac{1}{\sqrt{2\pi}} \left[ \int_{-\infty}^0 x^2 e^{-(-a+ik)x} dx + \int_0^\infty x^2 e^{-(a+ik)x} dx \right] \\ &= \frac{1}{\sqrt{2\pi}} \left[ -\frac{2}{(-a+ik)^3} + \frac{2}{(a+ik)^3} \right] \\ &= \frac{1}{\sqrt{2\pi}} \left[ \frac{4a(a^2-3k)}{(a^2+k^2)^3} \right] \\ &= \sqrt{\frac{2}{\pi}} \left[ \frac{2a(a^2-3k)}{(a^2+k^2)^3} \right]. \end{aligned}$$

b. Recall that the Heaviside function  $H$  is defined as

$$H(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x < 0. \end{cases}$$

Thus,

$$H\left(1 - \frac{|x|}{2}\right) = \begin{cases} 1 & \text{if } |x| < 2 \\ 0 & \text{if } |x| > 2. \end{cases}$$

If  $f(x) = \left(1 - \frac{|x|}{2}\right) H\left(1 - \frac{|x|}{2}\right)$ , we see from (1) that, by definition, the Fourier Transform of  $f$  is given by

$$\begin{aligned} \mathcal{F}\{f(x)\} = F(k) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \left(1 - \frac{|x|}{2}\right) H\left(1 - \frac{|x|}{2}\right) e^{-ikx} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-2}^2 \left(1 - \frac{|x|}{2}\right) e^{-ikx} dx \\ &= \frac{1}{\sqrt{2\pi}} \left[ \int_{-2}^0 \left(1 + \frac{x}{2}\right) e^{-ikx} dx + \int_0^2 \left(1 - \frac{x}{2}\right) e^{-ikx} dx \right]. \end{aligned}$$

Now, we see by integration by parts that

$$\begin{aligned} \int_c^d \left(1 \pm \frac{x}{2}\right) e^{-ikx} dx &= \frac{i}{k} e^{-ikx} \Big|_c^d \pm \frac{1}{2} \int_c^d x e^{-ikx} dx \\ &= \frac{i}{k} e^{-ikx} \Big|_c^d \pm \frac{1}{2} \left[ \frac{ix}{k} e^{-ikx} \Big|_c^d - \frac{i}{k} \int_c^d e^{-ikx} dx \right] \\ &= \frac{i}{k} e^{-ikx} \Big|_c^d \pm \frac{1}{2} \left[ \frac{ix}{k} e^{-ikx} \Big|_c^d + \frac{e^{-ikx}}{k^2} \Big|_c^d \right]. \end{aligned}$$

Thus,

$$\int_{-2}^0 \left(1 + \frac{x}{2}\right) e^{-ikx} dx = \frac{1 - e^{2ik} + 2ik}{2k^2}$$

and

$$\int_0^2 \left(1 - \frac{x}{2}\right) e^{-ikx} dx = \frac{1 - e^{-2ik} - 2ik}{2k^2}$$

Therefore, using the definition of the complex exponential and various trigonometric identities, we have that

$$\begin{aligned} \mathcal{F}\{f(x)\} = F(k) &= \frac{1}{\sqrt{2\pi}} \left[ \int_{-2}^0 \left(1 + \frac{x}{2}\right) e^{-ikx} dx + \int_0^2 \left(1 - \frac{x}{2}\right) e^{-ikx} dx \right] \\ &= \frac{1}{\sqrt{2\pi}} \left[ \frac{1 - e^{2ik} + 2ik}{2k^2} + \frac{1 - e^{-2ik} - 2ik}{2k^2} \right] \\ &= \frac{1}{\sqrt{2\pi}} \left[ \frac{1}{k^2} - \frac{e^{-2ik} + e^{2ik}}{2k^2} \right] \\ &= \frac{1}{\sqrt{2\pi}} \left[ \frac{1 - \cos 2k}{k^2} \right] \\ &= \frac{1}{\sqrt{2\pi}} \left[ \frac{1 - (\cos^2 k - \sin^2 k)}{k^2} \right] \\ &= \frac{2 \sin^2 k}{\sqrt{2\pi} k^2}. \end{aligned}$$

□

**Problem 2.** Find the Laplace Transforms of the following functions:

a.  $f(t) = \int_0^t \frac{\sin ax}{x} dx,$

b.  $f(t) = tH(t - a).$

*Solution.*

□

**Problem 3.** Solve the following integral equations:

a. 
$$\int_0^\infty f(x) \sin kx dx = \begin{cases} 1 - k & k < 1 \\ 0 & k > 1 \end{cases},$$

b. 
$$\int_{-\infty}^\infty \frac{f(t)}{(x-t)^2 + 4} dt = \frac{1}{x^2 + 9}.$$

*Solution.*

□

**Problem 4.** Show that

$$\int_0^\infty F_s(k)G_c(k) \sin kx dx = \frac{1}{2} \int_0^\infty g(\xi) [f(\xi + x) - f(\xi - x)] d\xi$$

where

$$F_s(k) = \sqrt{\frac{2}{\pi}} \int_0^\infty f(x) \sin kx dx$$

and

$$G_c(k) = \sqrt{\frac{2}{\pi}} \int_0^\infty g(x) \cos kx dx.$$

*Solution.*

□

**Problem 5.** Apply the Fourier Transform to solve the following initial value problem for the heat equation:

$$\begin{aligned}\frac{\partial u}{\partial t} &= a^2 \frac{\partial^2 u}{\partial x^2} + f(x, t), \quad -\infty < x < \infty, \\ u(x, 0) &= \phi(x), \quad t > 0.\end{aligned}$$

*Solution.*

□

**Problem 6.** Evaluate the following definite integrals:

a.  $\int_0^\infty \frac{\sin ax \sin bx}{x^2} dx,$

b.  $\int_0^\infty \frac{(a^2 - x^2)^2}{(x^2 + a^2)^4} dx, \quad a > 0.$

*Solution.*

□



**Problem 7.** Use the Fourier Sine Transform to solve the Laplace equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \quad 0 < x < \infty$$

with the boundary data  $0 < y < L$

$$\begin{aligned} u(x, L) &= 0, & u(x, 0) &= f(x), \\ u(0, y) &= 0, & u(x, y) &\rightarrow 0 \text{ as } x \rightarrow \infty \text{ uniformly in } y. \end{aligned}$$

*Solution.*

□

**Problem 8.** Apply the Fourier Transform to solve the 3-dimensional wave problem

$$\frac{\partial^2 u}{\partial t^2} = a^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right), \quad -\infty < x, y, z < \infty,$$

subject to the initial conditions

$$\begin{aligned} u(x, y, z, t)|_{t=0} &= 0 \\ \frac{\partial u(x, y, z, t)}{\partial t} \Big|_{t=0} &= \delta(x, y, z). \end{aligned}$$

*Solution.*

□

**Problem 9.** Show that if  $E$  is a solution of an  $m$ -th order partial differential equation

$$P(\partial)u = \sum_{k=0}^m a_k \partial^k u = \delta,$$

where  $\delta$  is the Dirac delta function, then  $E * f$  is the solution of the partial differential equation  $P(\partial)u = f$ , where  $*$  is the convolution.

*Solution.*

□