

Homework Assignment 11

Matthew Tiger

December 4, 2016

Problem 14.2.2. Let $f_c : \mathbb{C} \rightarrow \mathbb{C}$, $f_c(z) = z^2 + c$ for $c \in \mathbb{C}$. Recall that a period n point z_0 is super attracting if $(f^n)'(z_0) = 0$.

- i. If z_0 and z_1 are the fixed points of f_c , show that $f'(z_0) + f'(z_1) = 2$. Deduce that there can be at most one attracting fixed point. Give an example to show that f_c may not have any attracting fixed points.
- ii. Show that if f_c has a super-attracting fixed point z_0 , then $z_0 = 0$ and $c = 0$.
- iii. Find the value of c such that f_c has a super-attracting 2-cycle and give the associated 2-cycle.
- iv. Why is it that $z = 0$ is a point in the orbit of a cycle if and only if the cycle is super-attracting?
- v. If f_c has a super-attracting 3-cycle, show that c satisfies the equation

$$c^3 + 2c + c + 1 = 0$$

Solution. i. If z_0 and z_1 are fixed points of f_c , then they are the roots of the equation $z^2 - z + c = 0$. Note that the solutions to this equation are of the form

$$z_0 = \frac{1 + \sqrt{1 - 4c}}{2}, \quad z_1 = \frac{1 - \sqrt{1 - 4c}}{2}. \quad (1)$$

Since $f'_c(z) = 2z$, we see from (1) that

$$f'_c(z_0) + f'_c(z_1) = 2(z_0 + z_1) = 2 \left(\frac{1 + \sqrt{1 - 4c}}{2} + \frac{1 - \sqrt{1 - 4c}}{2} \right) = 2.$$

Suppose that $z_0 = r_0 e^{i\theta_0}$ is an attracting fixed point and let $z_1 = r_1 e^{i\theta_1}$. Since z_0 is attracting, we have that $|f'_c(z_0)| = 2|z_0| \leq 1$ which implies that $r_0 \leq 1/2$. Note that by the relation $f'_c(z_0) + f'_c(z_1) = 2$ we have that $z_1 = 1 - z_0 = 1 - r_0 e^{i\theta_0}$. Thus,

$$|z_1| = |1 - r_0 e^{i\theta_0}| \geq ||1| - |r_0 e^{i\theta_0}|| = |1 - r_0|.$$

If $r_0 = 1/2$, then $z_1 = z_0$ and there is at most one fixed point, otherwise if $r_0 < 1/2$, then $|z_1| \geq |1 - r_0| > 1/2$ which implies that $|f'_c(z_1)| > 1$ or that z_1 is repelling.

If $c = 5/4$, then we see that $|f'_c(z_0)| = |f'_c(z_1)| = \sqrt{5}/2$ where $\sqrt{5}/2 > 1$. Thus, f_c may not have any attracting fixed points.

- ii. Suppose that z_0 is a super attracting fixed point of f_c . Then $|f'_c(z_0)| = 2|z_0| = 0$. Since $|z_0| = 0$ if and only $z_0 = 0$, we readily see that $z_0 = 0$. Note that z_0 is of the form presented in (1). Thus, $(1 \pm \sqrt{1-4c})/2 = 0$ which implies that $1 - 4c = 1$ or that $c = 0$.
- iii. The 2-cycles of f_c are solutions of the equation $f_c^2(z) - z = 0$ that are also not solutions of $f_c(z) - z = z^2 - z + c = 0$. Factoring $f_c^2(z) - z$ we see that

$$f_c^2(z) - z = (z^2 - z + c)(z^2 + z + c + 1) = 0$$

if and only if z is a fixed point or if

$$z_2 = \frac{-1 - \sqrt{-3-4c}}{2}, \quad z_3 = \frac{-1 + \sqrt{-3-4c}}{2}.$$

Thus, $\{z_2, z_3\}$ forms a 2-cycle of f_c . This 2-cycle will be super-attracting if and only if

$$|(f_c^2)'(z_2)| = |f'_c(z_2)f'_c(z_3)| = |(-1 - \sqrt{-3-4c})(-1 + \sqrt{-3-4c})| = 4|1+c| = 0$$

Thus, the 2-cycle is super-attracting if and only if $c = -1$. Therefore, the super attracting 2-cycle is $\{0, -1\}$.

- iv. For an n -cycle $\{z_0, \dots, z_{n-1}\}$ of f_c we see that

$$|(f_c^n)'(z_0)| = |f'_c(z_0) \cdots f'_c(z_{n-1})| = 2^n |z_0 \cdots z_{n-1}|. \quad (2)$$

Thus, from (2), we have that $\{z_0, \dots, z_{n-1}\}$ is a super-attracting n -cycle of f_c if and only if $|(f_c^n)'(z_0)| = 0$ if and only if $z_i = 0$ for some $i = 0, \dots, n-1$.

- v. Suppose that $\{z_0, z_1, z_2\}$ is a super-attracting 3-cycle of f_c . Thus, we must have that

$$|f'_c(z_0)f'_c(z_1)f'_c(z_2)| = 2^3 |z_0 z_1 z_2| = 0$$

Without loss of generality, we may assume that $z_0 = 0$. Using the fact that $f_c(z_0) = z_1 \neq z_0$ and $f_c^2(z_0) = z_2 \neq z_0$, we see that

$$\begin{aligned} z_1 &= f_c(z_0) = z_0^2 + c = c \\ z_2 &= f_c^2(z_0) = (z_0^2 + c)^2 + c = c^2 + c \end{aligned}$$

In order for this to be a 3-cycle, we require that $f_c^3(z_0) = z_0 = 0$, i.e. we require that

$$\begin{aligned} f_c^3(z_0) &= f_c(f_c^2(z_0)) = (c^2 + c)^2 + c \\ &= c^4 + 2c^3 + c^2 + c \\ &= c(c^3 + 2c^2 + c + 1) = 0. \end{aligned}$$

However, we must have that $c \neq 0$ or $z_0 = 0$ would not generate a 3-cycle. Therefore, $\{z_0, z_1, z_2\}$ is a super-attracting 3-cycle if and only if $c^3 + 2c^2 + c + 1 = 0$. □

Problem 14.3.1.*Solution.*

Problem 14.3.3.*Solution.*