



GANARND

G GANA COSMETIC

dp Dr.PARK

GANARND MEDICAL DEVICES

We also supply the research and development service for customers. We develop and manufacture any products in dermal fillers. Mesotherapy and medical cosmetics field according to customer's need Our MOQ of R&D is reasonable with competitive price

www.ganarnd.co.kr



Our business division is divided into three fields

- Medical devices
- Cosmeceuticals and gene therapy products
- R&D and manufacturing

We have three companies

GANA R&D Co.,Ltd

- Producing dermal fillers, mesotherapy products, and clinic care products

GANA COSMETIC Co.,Ltd

- Producing cosmeceutical and conventional cosmetics

Dr PARK Co.,Ltd

- R&D of gene therapy products
 - DNA sequencing (NGS, Sanger sequencing)
 - DNA, RNA synthesis
 - Bioinformatics
 - RNA engineering (si RNA)
 - Gene engineering (CRISPR Cas 9, TALENS)
 - Peptide sequencing
 - Peptide synthesis
 - Protein synthesis
 - Exosome drug delivery system
 - Induced pluripotent stem cell
 - Human tissue culture using bio compatible polymer 3D printing technique and IPSC
-

GANA R&D

GANA COSMETIC

Dr.PARK

Address

555, sunil technopia, Dunchon-daero, Jungwon-gu,
Seongnam-si, Gyeonggi-do, South Korea
Office #1108, Factory 1 #103, Factory 2 #905, Factory 3 #902,
Factory 4 # 812, Factory 5 # 909, Factory 6 #904,
Factory 7 #1206, Factory 8 #1208, Factory 9 #913
Zip Code : 13215

TEL : 82-31-732-0242

FAX : 82-2-6455-7512

E-mail : dydgh416@naver.com

Homepage : www.ganarnd.co.kr

Dermal fillers

04	GANA MHS (Semi permanent)
05	Filler X (PCL + HA filler)
08	GANA HA (HA filler)
10	GANA PH (PLLA + HA filler)
13	GANA PN (PDRN syringe type)
15	GANA FILL (CE approved PLLA filler)
18	GANA CH (Calcium + HA filler)
19	GANA PNS (PDRN filler)
20	GANA PFS (Peptide filler)
21	GANA HA body (Hyaluronic acid dermal fillers)
26	GANA booster (Biorevitalization)



We also supply the research and development service for customers. We develop and manufacture any products in dermal fillers. Mesotherapy and medical cosmetics field according to customer's need Our MOQ of R&D is reasonable with competitive price

www.ganarnd.co.kr

GANA MHS (Semi permanent)

Dermal filler composed of

- Poly methyl methacrylate (PMMA) 100mg/cc
- HA (Cross linked hyaluronic acid) 20mg/cc

Characterized by semi permanent duration

How to use

1.2ml prefilled syringe

1. Attach enclosed needle
2. Injection layer-subcutaneous layer
3. Can be used for Nose, forehead, cheek, chin, Nasolabial fold
4. Contraindication Lip eye

PMMA dermal filler-GANA MHS

- PMMA (Polymethyl-methacrylate) fillers contain about 10 percent of 40um sized PMMA microspheres that are suspended in 90 percent of cross linked hyaluronic acid gel.
- One year after HA gel breaks down and your body produces its own natural collagen to fill out the space under the skin.

What is PMMA

Poly(methyl methacrylate)(PMMA), also known as acrylic or acrylic glass as well as by the trade names Plexiglas, Acrylite, Lucite, and Perspex among several others (see below), is a transparent thermoplastic often used in sheet form as a lightweight or shatter-resistant alternative to glass. The same material can be utilized as a Medical devices, casting resin, in inks and coatings, and has many other uses.



GANA MHS is 1.2ml prefilled syringe type

GANA MHS vs Existing PMMA fillers

	Existing PMMA	GANA MHS	
PMMA	20%	10%	
Bovine collagen	3.5%	None	Bovine collagen makes initial volume
HA	none	2%	HA makes initial volume for one year
Correction by hyaluronidase	Impossible	Possible	Initial removal of injected filler is possible

Filler X

(PCL + HA filler)

FILLER X(Patented Dermal filler)

New generation of dermal fillers




1.2ml syringe × 1EA/box

Major component

- Made of polycaprolactone 50mg/cc
- Cross linked hyaluronic acid gel 20mg/cc (HA dermal fillers)
- PCL 5%+ HA 95%
- Average 50um sized polycaprolactone

1.2ml prefilled syringe


What is PCL

Polycaprolactone	
	
Names	
IUPAC name (1,7)-polyoxepan-2-one	
Other names 2-Oxepanone homopolymer 6-Caprolactone polymer	
Identifiers	
CAS Number	24980-41-4
Abbreviation	PCL
Properties	
Chemical formula	(C ₆ H ₁₀ O ₂) _n
Density	1.145 g/cm ³
Melting point	60°C (140°F)
Except where otherwise noted, data are given for materials in their standard state (at 25°C[77°F], 100 kPa). Inpobox references	

Polycaprolactone (PCL) is a biodegradable polyester with a low melting point of around 60°C and a glass transition temperature of about -60°C. The most common use of polycaprolactone is in the manufacture of speciality polyurethane. Polycaprolactones impart good water, oil, solvent and chlorine resistance to the polyurethane produced.

This polymer is often used as an additive for resins to improve their processing characteristics and their end use properties (e.g., impact resistance). Being compatible with a range of other materials, PCL can be mixed with starch to lower its cost and increase biodegradability or it can be added as a polymeric plasticizer to PVC.

Polycaprolactone is also used for splinting, modeling, and as a feedstock for prototyping systems such as Fused Filament Fabrication 3D Printers.



PCL collagen stimulator

For those clients looking for a longer lasting dermal filler, PCL is an excellent option. PCL fillers work by not only correcting the wrinkles and folds in your skin, but also by stimulating your body's own collagen for a long-lasting, natural result. It is also fully resorbed by your body.

How it works

Collagen is one of the important support structures found naturally in our skin, helping to maintain the appropriate tightness, elasticity and suppleness.


Over the course of time, our skin gradually loses its natural collagen, leading to facial laxity and the formation of wrinkles and folds.

Filler X works differently to traditional dermal fillers, which only address lines, wrinkles and folds. Thanks to its unique properties, it not only corrects wrinkles and folds but by stimulating your own collagen to really treat the underlying causes of facial ageing, the effects are gradual and longer-lasting giving you a natural youthful appearance.

FILLER X vs HA and Ellanse

	HA fillers	Pcl filler	FILLER X
Duration	1yr	2yr and 4yr	2yr and 4yr
Migration	Yes	No	No
Collagen stimulation	No	Yes	Yes
Elimination by hyaluronidase	Yes	No	Yes
Swelling	No	Yes	No
Correction	Yes	No	Yes

FILLER X



Disadvantages of pre existing PCL fillers

1. Initial swelling
2. Initial volume loss
3. Impossibility of correction

The main complaints of HA dermal fillers

Short duration of HA fillers

The merits of FILLER X

1. The Even and longer volume augmentation over than 4years by cross linked hyaluronic acid gel and Later volume augmentation by the collagen formation by PCL
2. The lower initial swelling by the absence of carboxy methyl cellulose
3. The immediate correction is possible by hyaluronidase
4. Injected calcium PCL induce collagen synthesis So that the duration of FILLER X is much longer than HA fillers
5. Newly formed collagen prohibit the migration of FILLER X
6. Patented products-exclusiveness

Precaution

- All kinds of collagen stimulating dermal filler start to make collagen in 4-6 weeks.
- So be careful before use.
- Once you inject collagen stimulating dermal filler on forehead, lip There will be lump growing

Among our products

- GANA V, X
- GANA PH
- FILLER X
- Are collagen stimulating filler

**Do not use these products on forehead,
lip And periorcular area**

Recommended dose


- Both Nasolabial fold : 1-2ml
- Both Cheeck : 2-6ml
- Chin : 1-3ml
- Nose : 1-2ml

FILLER X - Patented



######

Depth



Recommended application of GANA HA


A type : cheek, chin, forehead, nose

B type : nasolabial fold, lips

C type : fine wrinkle, around eyes

Recommended dose

1. Nose 1-2ml
2. Nasolabial fold 2-3ml
3. Forehead 3-5ml
4. Lip each side over 1ml
5. Cheek 4-10ml
6. Short chin 2-4ml
7. Sunken eye 1ml
8. Temporal area depression each over 2ml



Specification

Test Items	Standard Provision	Result
Appearance	Colorless, transparent No visible foreign matter, no air bubbles.	Qualified
Hyaluronate Content	24.22.20mg/ml Booster : 10mg/ml	Qualified
PH	6.5~7.6	Qualified
Osmotic Pressure(mOsmol/L)	0.9-1.1	Qualified
Particle size distribution(μm)	Type A : 1mm : 23Guage Type B : 200-500um : 27Guage Type C : 50-100um : 30Guage Booster : under 30um	Qualified
Crosslinking agent BDDE residue(μg/g)	≤2.0	Qualified
Protein	≤1.0%	Qualified
Heavy metal	≤20ppm	Qualified
Bacteria Test	Should have no bacterial	Qualified
Bacteria Endotoxin Test	≤0.5EU/mL	Qualified
Viscosity	4-6X10 ⁶ CPs	Qualified
Hemolytic	Should be no hemolysis ring	Qualified
Conclusion	Conclusion : The produces were tested conform to the standard of GANARND.CO.LTD	

Recommended dose

GANA HA	Volume	Injection layer	Recommended guage	HA concentration (mg/cc)	Cross linking ratio	Recommended use
A type	1.2ml	Sub Q	25	24	10%	Large volume augmentation
B type	1.2ml	Deep dermis	27	22	10%	Moderate wrinkle
C type	1.2ml	Dermis	30	20	10%	Fine wrinkle
booster	2.5ml	Eipdermis	30	10	5%	Mestoherapy

GANA PH

(PLLA + HA filler)

Major component

GANA PH is dermal fillers

- Made of Poly-L-lactic acid 20mg/cc
- Cross linked hyaluronic acid gel 20mg/cc (HA dermal fillers)




PLLA 2% + HA 98%
Average 50um sized PLLA

1.2ml syringe × 1EA/box


What is PLLA

Poly(L-Lactide) is a biodegradable material with good bio-absorbability and bio-compatibility. After degradation in the body, PLLA products are excreted by metabolism and thus has no harmful side effects to the human body. It is for these reasons that PLLA products are currently widely used in the field of medicine, such as for one-time infusion equipment, medical surgical suture, drug delivery carriers, tissue engineering scaffolds, bone anchors, injectable microcapsules, microspheres, implants, elastomer materials for organ support, etc.

Mechanism



PLLA degradation




PLLA mechanism



PLLA replaces lost collagen, corrects your imperfection.
It lasts more than 2 years.


Freeze dried PLLA has micropore
Micropore absorb water to break down into small pieces
The broken PLLA particles induce controlled inflammation
So that there will be collagen synthesis
This SEM picture is of our PLLA



GANA PH

	HA fillers	PLLA filler	GANA PH
Duration	1yr	3yr	3yr
Migration	Yes	No	No
Collagen stimulation	No	Yes	Yes
Elimination by hyaluronidase	Yes	No	Yes
Swelling	No	Yes	No
Correction	Yes	No	Yes
Dissolution with water	Not needed	Many hours waiting	Not needed
Volume augmentation	Immediate effect	2 months	Immediate

GANA PH



Dismertis of pre exsiting PLLA fillers

1. Initial swelling
2. Initial volume loss
3. Impossibility of correction
4. Long dissolution time
5. 2 months of Time lag before volume augmentation

The main complaints of HA dermal fillers

Short duration of HA fillers

The merits of GANA PH

1. Early and longer volume augmentation over than 2years by cross linked hyaluronic acid gel and Later volume augmentation by the collagen formation by PLLA
2. The lower initial swelling by the absence of carboxymethylcellulose
3. The immediate correction is possible by hyaluronidase
4. Injected PLLA induce collagen synthesis duration of GANA PH is much longer than HA fillers
5. Newly formed collagen prohibit the migration of GANA PH

Recommended dose

- Both Nasolabial fold : 1-2ml
- Both Cheeck : 2-6ml
- Chin : 1-3ml
- Nose : 1-2ml

Precaution

- All kinds of collagen stimulating dermal filler start to make collagen in 4-6 weeks
- So be careful before use
- Once you inject collagen stimulating dermal filler on forehead, lip There will be lump growing

Among our products

- GANA V, X
- GANA PH
- FILLER X
- Are collagen stimulating filler

Do not use these products on forehead, lip And periocular area

GANA PN (PDRN syringe type)



Major component

- PDRN (poly deoxyribonucleotide from salmon sperm) 2%
- Glutathione 2mg/cc
- Cross linked Hyaluronic acid 10mg/cc

PDRN -polydeoxyribonucleotide

Definition

PDRN is the low molecular weight DNA complex fragments from salmon or trout sperm. It play a role in anti inflammatory, cell proliferation and tissue remodeling.



4 kinds of PDRN products

	Volume	Package	PDRN %	HA concentration	Cross linking HA	Glutathione
GANA PN	2.5ml × 2EA	Syringes	2%	10mg/cc	yes	2mg/cc
GANA PNS	1.2ml × 1EA	Syringe	2%	20mg/cc	yes	None
GANA PNV	3ml × 5EA	Vials	0.5%	5mg/cc	yes	2mg/cc
GANA PNV plus	3ml × 5EA	Vials	2%	10mg/cc	yes	2mg/cc

GANA PN, Prefilled syringe		GANA PNV, Prefilled vial	
Volume	2.5cc filled in 3cc syringe	2.5cc filled in 5cc vial	
Packaging	2 syringes in a box	5 vials in a box	
Concentration of PDRN	2%	0.5%	
Other component	Cross linked HA glutathione Lidocaine Q.S-PBS		Same as above

GANA PN vs Pre existing PDRN

	Effect	Existing products	GANA PN
PDRN content	Wound healing Cell regeneration	2%	2%
Glutathione	Whitening	None	2mg/cc
Cross linked hyaluronic acid	Duration 6month	None	10mg/cc

PDRN effect

- Safe and simple injection
- Protecting wrinkles
- Skin lifting effect
- Skin brightening
- Improving elasticity, turgidity and tonicity of skin
- Repairing damaged dermal cells
- High physical compatibility of Polynucleotide extracted from salmon

How to use

- Unpack box
- Clean the application area
- Attach enclosed needle
- Injection level will be dermis layer
- If there is bleeding, compress area for one minute
- Second session will be needed in a month

GANA PN contain cross linked hyaluronic acid so GANA PN can be used like dermal fillers for face augmentation. This is the merit of GANA PN compared to other PDRN products

Protocol

1. 2-3 session is needed
2. Interval one month
3. After second session
The effect can be seen

Preparation

Apply ice bad and make anesthesia with lidocaine cream both to reduce pain

Precaution

This product contain
Cross linked hyaluronic acid(HA)
Not non cross linked hyaluronic acid

Once injected to epidermis
This can cause nodule and papule
So injection layer will be dermis layer
And use mesotherapy gun needle depth more than 1.5mm

Do not inject at one point too much
Inject widely deeply and evenly

After injection

Compress treated area with dry gauze

Contraindication

1. Any kinds of active skin disease
2. Any kinds of general illness
3. Hypersensitive to this products
4. Bleeding disorder
5. Pregnancy, breast feeding

Side effect

- Bruising
- Hematoma
- Atrophy & Skin hypertrophy
- Erythema
- Urticaria

Use Mesogun. Derma roller or enclosed 30G needles



What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process. The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation.

Most importantly, PDRN provides the nucleotides necessary for cell regeneration. In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries

GANAFILL

(CE approved PLLA filler)



CE and ANVISA approved

What is biodegradable material

The term biodegradable is used to describe materials that decompose through the actions of bacteria, fungi, and other living organisms—simply put it has the ability to break down or decompose back into the natural environment without causing harm. Examples of well-known biodegradable materials used in medical industries are PLLA, PLGA, PGA, PDO and PCL.

What is PLLA

Poly(L-lactide) is a biodegradable material with good bio-absorbability and bio-compatibility. After degradation in the body, PLLA products are excreted by metabolism and thus has no harmful side effects to the human body. It is for these reasons that PLLA products are currently widely used in the field of medicine, such as for one-time infusion equipment, medical surgical suture, drug delivery carriers, tissue engineering scaffolds, bone anchors, injectable microcapsules, microspheres, implants, elastomeric materials for organ support, etc.

Mechanism

- PLLA (Poly-L-lactic acid)
- Polymer of Lactic acid
- Lactic acid $\rightarrow$ CO₂ + H₂O


PLLA immune reaction

Some PLLA Eaten by macrophage

- By immune reaction activate fibroblast to produce collagen. The other PLLA remain in human body—half life 3 years.

Mechanism and technique - II

Difference with existing PLLA infection



PLLA replaces lost collagen, corrects your imperfection. It lasts more than 2 years.

Injectable PLLA, the main component of GANAFill works in the deep dermis to create collagen.

**GANA FILL****INJECTABLE POLY-L-LACTIC ACID POWDER**

Reconstitute with 18ml of sterile water for injections
One vial contains Injectable PLLA 520mg
Made in Korea

**GANA FILL-Face**

This product is made from a mixture of PLLA powder FDA approved materials. GANA FILL is more innovative and creative than all existing products. It induces the formation of collagen which in turn has the effect of increasing volume of the part where the surgical procedure was done. The augmentation will continue until the 8th week and will last for 2 years after the procedure. Main areas of use are temple, cheek, This product uses the same main ingredients as Sculptra.

Reconstitute with distilled water

- Min 10cc
- Max 18cc

GANA FILL-X Body

This GANAFILL X contains large amounts of PLLA 1560mg PLLA, GANAFILL X is made for body augmentation. Breast, hip, male and female genital organ.

**Merits of GANAFILL vs Fat implantation**

1. Quicker dissolution time-under one hour
 - If you have no time, You can reconstitute water and shake well for 10minutes.
 - if you use vortex mixer, dissolution time will be under 60 seconds, shake 60seconds then just use it
2. Cheaper price
3. Larger quantity. GANAFILL 520mg VS other PLLA fillers 150mg (GANAFILL × 1560mg)
4. Better contouring effect
 - because GANAFILL has more PLLA
5. Less volume loss at first injection
 - In a few days of sculptra injection. There will be no augmentation, You will see Initial volume loss, You have to wait 1-2months until you see augmentation but GANAFILL has lower Volume loss.
6. Low nodule formation-more water more dilution lower grnuloma

Merits of GANAFILL vs Other PLLA fillers

1. Quicker procedure time under 1 hour
2. Cheaper price
 - Two or three GANAFILL are enough for full face augmentation
3. Faster recovery time under 2days
 - Recovery time of fat implantation is over 1 month



Effect and various application

- Face augmentation
- Face lifting
- Breast and hip augmentation
- Hand and neck wrinkle improvement
- Scar improvement
- Female and male genital organ
- Vagina tightening and penis shaft thickening

Vortex mixer



Face



First session

- for full face
- one vial is max

Second session

- One month later
- Half dose of first session
- Depth of injection
- Subcutaneous layer or periosteum
- Injection technique
- Fan technique using cannula

Precaution

1. Do not treatment prohibited area. (lip, nose, forehead)
2. Use distilled water only. (Do not use a saline solution or water.)
3. Don't inject at one spot too much, inject widely and deeply to avoid granuloma formation
4. Patient have to get second treatment. (Check progress)

	First session	Second session	Depth of injection	Injection technique
Temporal area	0.5 vial max	One month later half dose of first session	Subcutaneous layer or Periosteum	Fan shape injection using cannula
Nasolabial fold	0.5 vial max			
Cheek	1 vial max			
Ant zygomatic area	0.5 vial max			

GANA FILL procedure

1. Reconstitute with 18ml of sterile water at least 1hours before injection.
2. If you use vortex mixer, dissolution time will be under 60 seconds.
3. Use blunt needle larger than 21G.
4. For subcutaneous injection. For better performance inject deeper.
5. Use fan technique and avoid injecting too much in one area.
6. Do not use for forehead, lips, nose or areas near the eyes.
7. Use for face, breast and hip volume augmentation and acne scar recovery.
8. Effects can be seen 4 weeks after treatment up until 8 weeks after treatment. Volume augmentation lasts for 2 years.
9. Vigorous massaging is necessary around the area of treatment for 30 minutes every day for the first week
10. Two or three session is needed

GANA CH (Calcium + HA filler)



Main component

- Calcium hydroxyapatite 5%
- Cross linked hyaluronic acid 95%

New generation of dermal fillers

Made of Calcium hydroxyapatite + cross linked hyaluronic acid gel (HA dermal fillers)

- The main complaints of calcium dermal fillers are initial swelling, initial volume loss and impossibility of correction
- The main complaints of HA dermal are the short duration of HA fillers
- This GANA CH has only the merits of calcium and HA fillers but the demerits of calcium and HA dermal fillers

The merits of GANA CH

1. The Even and longer volume augmentation over than 2years by initial volume augmentation by cross linked hyaluronic acid gel and Later volume augmentation by the collagen formation by calcium hydroxyapatite
2. The lower initial swelling by the absence of carboxy methyl cellulose

3. The immediate correction is possible by hyaluronidase
4. This GANA CH has the merits of HA and calcium dermal fillers
5. Injected calcium hydroxyapatite induce collagen synthesis So that the duration of GANA CH is much longer than HA fillers
6. newly formed collagen prohibit the migration of GANA CH

Recommended are of GANA CH

- Nose
- Chin
- Forehead

Do not use in

- Cheek
- Lip

Injection layer

- Deep dermis

	Ha fillers	Calcium fillers	GANA CH
Duration	1yr	3yr	3yr
Migration	Yes	No	No
Collagen stimulation	No	Yes	Yes
Elimination by hyaluronidase	Possible	Impossible	Possible
Swelling	None	Mild swelling	None
Correction	Possible	Impossible	Possible

GANA PNS (PDRN filler)



Major component

- PDRN (poly deoxyribonucleotide from salmon sperm) 2%
- Cross linked Hyaluronic acid 20mg/cc

1.2ml syringe × 1EA/box

GANA PNS is HA dermal fillers containing PDRN

GANA PNS application

- Face rejuvenation
- Wrinkle removal
- Face volume augmentation
- Preventing sagging
- Improving elasticity, turgidity and tonicity of skin
- Repairing damaged dermal cells

How to use

- Unpack box
- Clean the application area
- Attach enclosed needle
- Injection level will be subcutaneous or dermis layer
- If there is bleeding, compress area for one minute

What is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process. The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation.

Most importantly, PDRN provides the nucleotides necessary for cell regeneration. In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries.

4 kinds of PDRN products

	Volume	Package	PDRN %	HA concentration	Cross linking HA	Glutathione
GANA PN	2.5ml × 2EA	Syringes	2%	10mg/cc	Yes	2mg/cc
GANA PNS	1.2ml × 1EA	Syringe	2%	20mg/cc	Yes	None
GANA PNV	3ml × 5EA	Vials	0.5%	5mg/cc	Yes	2mg/cc
GANA PNV plus	3ml × 5EA	Vials	2%	10mg/cc	Yes	2mg/cc

GANA PFS

(Peptide filler)

Peptide filler in prefilled syringe


1.2ml X 1EA

Main components

1. Cross linked HA 20mg/cc
2. Acetyl hexapeptide 8 (AH8)
3. Palmitoyl tetrapeptide 7
4. Palmitoyl pentapeptide 4
5. PBS

Main effects and use

1. Cross HA fill depressed wrinkle
2. AH8 paralyze facial muscle
3. Effective for nasolabial fold
4. Crow feets wrinkle and forehead wrinkle



GANA HA body

(Hyaluronic acid dermal fillers)

HA filler for body augmentation

GANA HA body is

GA dermal filler specially made for body contouring For example Breast augmentation buttock augmentation Genital organ

Main component is

Cross linked hayluronic acid 20mg/cc
Cross linking ratio 10% 50ml prefilled vial

Biphasic HA dermal fillers are characterized by

- Adding of non cross linked HA
 - Non cross linked HA disappear in 1-2wks
- Big particle size
- Relatively short duration
- Low cross linking ratio
- Easy to mold shape

Monophasic HA dermal fillers are characterized by

- Non adding of non cross linked HA
- Longer duration
- Relatively high duration
- Difficult to mold shape
- Relatively high cross linking ratio

GANA HA is characterized by

1. Non adding of non cross linked HA
2. Highest cross linking ratio
3. Highest HA concentration
4. GANA HA has small particle size so that doctors are easy to make shape
5. Highest viscosity than any other dermal fillers
6. Longest duration
7. Low extruding force



GANA HA body

Cross linked HA
50cc
HA concentration
20mg/cc
Cross linking ratio
10%

How to use

1. Remove aluminum cap and rubber stopper
2. Withdraw HA body with 10cc syringe
3. You will have 5 syringes for one vial
4. Make anesthesia at puncture
5. Puncture with 16G or 18G needle
6. Attache 16G or 18G canula to prefilled 10ml syringe
7. Inject GANA HA body to breast or hip
8. Injection are will be subcutaneous fat layer in the hip
9. For breast augmentation. Injection layer will be just above pectoralis major muscle
10. For breast injection. be careful not to damage mammary glands
11. for each breast and hip. minimum dose will be 50ml maximum dose for each breast will be 200ml

Recommended dose

- For vaginal tightening 50ml
- For partial breast augmentation 50-100ml each
- For full breast augmentation 200ml each
- For partial hip augmentation 50-150ml
- For full hip augmentation 200-400ml each

Merits of Hyaluronic acid (HA)dermal filler for body augmentation

	Removal	Price	Safety	Procedure time	Minimal invasive procedure?
HA dermal fillers	O	Expensive	O	Long	O
Conventional fillers (PMMA,PAAG)	X	Expensive	O X	Short	O
Implants	O But need major surgery	Expensive	O X	Long	X
GANA HA body	O	Reasonable price	O	Short	O

GANA HA body vs Existing HA fillers

	Monophasic (ex.Juvederm)	Biphasic (ex.restylane)	GANA HA body
Cross linking ratio %	6%	1-2%	10%
HA concentration (mg/cc)	20-24	20-24	20
Cross linking agent	BDDE	BDDE	BDDE
Duraron month	6-12	6-18	24
Particle size	None	100-2000um	500um
Viscosity. CPS.	2-4 X 106CPS	1-4 X 106CPS	2-4 X 106CPS
Package size	1-3ml	1-3m	50ml

Specification

Test Items	Standard Provision	Result
Appearance	Colorless,transparent No visible foreign matter, no air bubbles.	Qualified
Hyaluronate content	.20mg/ml	Qualified
PH	6.5~7.5	7.3
Osmotic pressure(mOsmol/L)	0.9-1.1	1
Particle size distribution(μ m)	Body : 500um : 23Guage	Qualified
Crosslinking agent BDDE residue(μ g/g)	≤ 2.0	Qualified
Protein	$\leq 0.1\%$	Qualified
Heavy metal	≤ 20 ppm	Qualified
Bacteria test	Shoule have no bacterial	Qualified
Bacteria endotoxin test	≤ 0.5 EU/mL	Qualified
Viscosity	2-4X10 ⁶ CPs	Qualified
Hemolytic	Should be no hemolysis ring	Qualified
Conclusion	Conclusion : The produces were tested conform to the standard of GANARND.CO.LTD	

How to withdraw

- GANA HA body is prefilled vial type
- And has high viscosity
- Need special technique to withdraw

Prepare normal saline,
18g needle and six 10ml syringe



Open caps of NS and body



Withdraw 10ml normal saline



Mix normal saline with body




Then you can withdraw Body easily



How to use for vaginal tightening

1. Pull out solution, 10ml syringe
2. Prepare 10ml X 4 syringes
3. Locally make anesthesia vaginal inlet puncture point
4. Inject 10ml ha body at submucosal layer.
5. Injection location is 5cm inside from vaginal inlet
6. Inject each 10ml ha body to 3,6,9,12 clock location, Totally 40ml
7. In two month second session may be needed half dose of first session
8. Be careful not to damage urethra and rectum


Vagina treatment of procedure




Injection layer

1. Cooper's ligaments
2. Breast lobule
3. Extralobular duct
4. Ductal ampulla (Reservoir)
5. Main duct
6. Nipple
7. Skin
8. Subcutaneous fat
9. Mammary layer fatty tissue
10. Retromammary fat
11. Lymph nodes
12. Pectoralis major muscle
13. Pectoralis minor muscle
14. Rib

Breast




Puncture point



Various depression on buttock

Buttock augmentation




Progress


1. For 3-4 days there might be swelling and pain
2. Prescribe pain killer, antibiotics and anti inflammatory for 5days
3. The swelling and pain is normal cause this products has very high cross linking ration almost 10%, it absorb water and getting bigger for a few days It maybe cause swelling and pain
4. second session in a month is necessary
The injected dermal filler can migrate to adjacent area so in second session practitioner must correct deformed shape

Breast filler


Before and after




Before and after (300ml used)




Before and after (300ml used)



Before and after (300ml used)




Before and after (300ml used)



Before and after(200ml both used)



Left 150ml VS right before



Buttock filler

Before and after (300ml used)



News paper articles in JAPAN



GANA booster (Biorevitalization)



2.5ml × 1EA/box

GANA booster (biorevitalization)

The most of hyaluronic acid gel products for mesotherapy are non cross linked HA. But GANA booster (biorevitalization) is cross linked HA gel same as dermal fillers.

- So has much better duration and safety
- Non cross linked HA gel is can not be autoclave sterilized
- Also GANA booster (biorevitalization) has lidocaine to lower pain

The merits of GANA booster (biorevitalization)

Prefilled syringe

- Cross linked hyaluronic acid 10mg/cc (longer duration and better efficacy)
- Autoclave sterilized (better safety)
- Glutathione 0.1% for whitening

Preparation

Apply ice bad and make anesthesia with lidocaine cream both to reduce pain

Precaution

This GANA booster (biorevitalization) contain

- Cross linked hyaluronic acid (HA)
- Not non cross linked hyaluronic acid

Once injected to epidermis

- This can cause nodule and papule
- So injection layer will be dermis layer
- And use mesotherapy gun. needle depth more than 1.5mm

Do not inject at one point too much

Inject widely deeply and evenly

After injection

Compress treated area with dry gauze

GANA booster (biorevitalization) vs Other HA solution

	GANA HA hydro	Other
Duration	6 month	2 wks
Safety	Autoclave sterilization	No sterilization
Pain	Less painful	Painful
Whitening	Yes	No

PROTOCOL


1. 2-3 session is needed
2. Interval one month
3. After second session the effect can be seen

GANA booster is usually used by meso gun
If you have not meso gun
You can use this meso needle



Certificate of analysis

- Products name : GANA HA Hyrdo
- Production date : 2015-9-15
- Expiratory date : 2016-9-15
- LOT No :
- CAS No. : 9004-61-9
- Molecular formula : $(C_{14}H_{21}NO_{11})_n$
- Content : 99.9 %
- Grade : Medicine grade.
- Molecular Structure :



Specification

Test items	Standard provision	Result
Appearance	Colorless, transparent No visible foreign matter, no air bubbles.	Qualified
Hyaluronate content	Hydro : 10mg/ml	Qualified
PH	6.5 ~7.5	7.3
Osmotic pressure (mOsmol/L)	0.9 ~1.1	1
Particle size distribution (μm)	Hydro : under 30um	Qualified
Crosslinking agent BDDE residue (μg/g)	≤ 2.0	Qualified
Protenin	≤ 0.1 %	Qualified
Heavy metal	≤ 20ppm	Qualified
Bacteria test	Shoule have no bacterial	Qualified
Bacteria endotoxin test	≤ 0.5EU/mL	Qualified
Viscosity	10-50X10 CPs	Qualified
Hemolytic	Should be no hemolysis ring	Qualified

Cosmetics

29	ANA PHA (PHA peeling)
31	GINO FACE/GINO BODY (Acne/Keratois pilaris)
33	ANA TOX (FDA registered spraying peptide)
36	ANA PNV (PDRN vial type)
39	REGEN PEEL (TCA peeling)
41	ANA Cocktail (Whitening)
43	ANA SCALP (Hair care)
45	ANA PNV+ (High PDRN)
49	ANA EYE (Eye bag removal)
50	ANA DM (PLLA meso)
53	ANA PF (Expression wrinkle removal)
54	ANA GYNO (Vaginal whitening with vegetable estrogen)
55	ANA TM (FDA registered mask pack)
59	ANA MASK (Facial burn mask)
62	ANA TC plus (PDRN, PHA.AH8 ointments)
64	ANA TC (PDRN, PHA.AH8 ointments)
66	ANA BB (BB with PDRN ointments)
67	ANA KA (Kojic acid ointment)
68	ANA V line (Face contour improval)
69	Dr park's TELO (Telomere)
70	ANA DA (Lipolysis)
71	ANA booster-V
72	ANA HA-V
73	ANA peptides (APDN)
74	ANA peptides (Carnosine)
75	ANA peptides (AH8)
76	ANA peptides (PP3)
77	ANA peptides (PT1)
78	ANA peptides (MP17)
79	ANA peptides (AGH1)
80	ANA peptides (AT5)
81	ANA peptides (NP1)
82	ANA AM

We also supply the research and development service for customers.
We develop and manufacture any products in dermal fillers.
Mesotherapy and medical cosmetics field according to customer's
need Our MOQ of R&D is reasonable with competitive price

www.ganarnd.co.kr

GANA PHA

(PHA peeling)

PHA peeling

Ingredients

- Polyhydroxy acid 10%
- Alpha arbutin 1% for whitening

PHA is new generation AHA



GANA PHA 40 ml

Safety of PHA

Even very sensitive skin types can tolerate PHAs, including atopic and rosacea skin. It is thought that the gentle, nonirritating action of PHAs may be partially related to a larger molecular composition than that of conventional AHAs.

Larger molecules are believed to penetrate the skin at a slower rate, causing less irritation. This reduced irritation, coupled with natural antioxidant properties, gives PHAs a gentle, competitive edge over other anti-aging treatments.

What is PHA (Exfoliant, Antioxidants)

Polyhydroxy acids are ingredients such as gluconolactone and lactobionic acid. They're supposed to be as effective as AHAs, but less sensitizing.

Gluconolactone and lactobionic acid are chemically and functionally similar to AHAs (such as glycolic acid).

The significant difference between them and AHAs is that gluconolactone and lactobionic acid have larger molecular structures, which limits their ability to penetrate the skin, resulting in less potential for sensitizing side-effects.

Supposedly, this reduced absorption into the skin does not hamper their effectiveness. Does that mean gluconolactone and lactobionic acid are better for your skin than AHAs in the form of glycolic acid or lactic acid?

According to an Internet-published class lecture by Dr. Mark G. Rubin, a board-certified dermatologist and assistant clinical professor of dermatology at the University of California, San Diego, research on gluconolactone demonstrated only a "6% decrease in dermal penetration" in comparison to glycolic acid, which "isn't a dramatic improvement."

Gluconolactone may be slightly less sensitizing for some skin types, but this isn't the magic bullet for exfoliation some cosmetics companies have been extolling. Polyhydroxy acids can also function as antioxidants and may promote some amount of improvement in skin's surface strength.

PHA effect

- PHAs have been found to be compatible with clinically sensitive skin, including rosacea and atopic dermatitis, and can be used after cosmetic procedures.
- PHAs provide additional humectant and moisturization properties compared with AHAs and can enhance stratum corneum barrier function, therefore increasing the skin's resistance to chemical challenge.
- Most PHAs also possess antioxidant properties.
- PHAs such as gluconolactone or lactobionic acid may be used in combination with other products, ingredients, or procedures such as laser and microdermabrasion to provide additional benefits to therapy or to enhance the therapeutic effect.
- Several studies were conducted in support of this, and methods and results are discussed. In summary, PHA-containing products were used in combination with retinoic acid in treating adult facial acne and were found to be well tolerated.
- PHAs plus retinylacetate (pro-vitamin A) in a cream base exhibited significant antiaging skin benefits such as skin smoothing and plumping.
- PHAs plus hydroquinone showed excellent improvement in antiaging and skin lightening parameters.
- Finally, PHA-containing products were shown to be compatible with African American, Caucasian, and Hispanic/Asian skin and provided significant improvements in photoaging in these populations.

PHA face peel is indicated for :

- Acne-prone skin
- Comedones
- Enlarged pores
- Hyperpigmentation and uneven skin tone
- Skin aging prevention
- Early signs of photoaging such as skin dryness,
- Loss of elasticity and fine lines
- Hyperkeratosis (rosacea, seborrheic dermatitis)
- Actinic dermatitis
- Correction and prevention of natural aging signs such as withering and first wrinkles pigmentation

Benefit of PHA

- Promote cell turnover. By sloughing off dead skin cells, revealing smoother, healthy new cells underneath.
- PHAs have humectant effects that allow for increased moisturisation. When skin is sufficiently moisturised, it looks smoother and healthier.
- Enhancing skin smoothness and elasticity.
- Antioxidant effects. Gluconolactone is an antioxidant known for its free radical-fighting ability, protecting skin from harmful UV exposure.
- Increased skin barrier function. Our skin barrier, or stratum corneum, plays a vital role in keeping out irritating chemicals and substances.
- Improving skin clarity and brightness; reducing the appearance of fine lines, wrinkles, and pore size
- Reduce flare-ups in inflammatory conditions. PHA products can lower inflammatory episodes of acne, rosacea, dermatitis, psoriasis and melasma

How to use

1. Wash face before go to bed
2. Apply GANA PHA
3. Interval 1 week
4. 2-3 sessions

Prepare GANA PHA, brush and bowl



Soak brush with GANA PHA



Pump GANA PHA into bowl



Spread GANA PHA



GINO FACE GINO BODY (Acne/Keratois pilaris)

GINO FACE/GINO BODY (Acne/Keratois pilaris)

- GINO face and GINO body contain fresh water sponge spicules
- This spicules induce mild inflammation so that skin cell turn over very fast
- Eventually make healthy new skin, make skin younger
- Also these spicules make holes,
These small holes help acne pustules to come out of pores

GINO FACE is developed for acne treatment

GINO BODY is developed for keratosis pilaris



GINO Face. 45gram



GINO Body. 45gram



Spongilla is a genus of freshwater sponges in the family Spongillidae found in lakes and slow streams. Sponges of the genus Spongilla attach themselves to rocks and logs and filter the water for various small aquatic organisms such as protozoans, bacteria, and other free-floating pond life. Unlike marine sponges, freshwater sponges are exposed to far more adverse and variable environmental conditions, so they have developed gemmules as a means of dormancy. When exposed to excessively cold or otherwise harsh situations, the sponges form these gemmules, which are highly resistant "buds" that can live dormant after the mother sponge has died. When conditions improve, the gemmules "germinate" and a new sponge is born.

GINNO peel Series

It is a highly effective acne treatment that regenerates the skin by penetrating the meticulously refined spicules

Main ingredients

GINO Face. 45gram

- Fresh water Poriferan spicules
- Salicylic Acid
- Alpha Arbutin
- Niacin
- Sodium Hyaluronic Acid

GINO Body. 45gram

- Fresh water Poriferan spicules
- Glycolic Acid
- Urea
- Hyaluronic Acid

GINO Peel Treatment

This treatment procedure turns the skin smooth and healthy by solving intensified troubles in depth and efficiently increasing skin improvement ability, to take care of troubles and form an immune system to prevent recurrence.

By using micro needles called spicules, skin cells are stimulated and regenerated, and naturally eliminate dead skin cells by improving immunity and promoting regeneration.

GINO face is Recommended for

- Papule or inflamed acne
- Excessive excretion of sebum
- Dry skin with many fine wrinkles
- Those with red pimples
- Those who want simple treatment with small troubles

GINO body is Recommended for

- Keratosis pilaris

Advantage of GINO Peel

- It can be performed on all skin types such as sensitive, dry, or oily skin.
- It improves sebum, dead skin cells, acne, and inflammation.
- It is a peeling procedure using natural spicules, so has almost no side effects.
- It is also effective on keratosis pilaris, also called goose flesh.

Process of GINO Peel

- Gino Peel effective on acne is penetrated into the skin
- Penetrated spicule causes cell divisions for 72 hours to induce loss of dead skin cells, suppress sebum, eliminate acne germs, and reduce inflammation
- Dead skin cells start to come off 3 days later
- Skin becomes thicker and maintains the healthy condition

How to apply

- Clean up that area where you would apply.
- Apply proper volume of GINO Peel and massage by finger tip with soft pressure.
- It might feel tingly, it is natural reaction cause of implanting spicule into skin.
- Massage for 5 min and leave it until dry.
- Next morning, Rinse with water gently, no rubbing.
- Repeat same procedure Once or twice a week as tolerated
- Do apply facial pack to apply enough moisturizing.
- Avoid rinsing for 72 hrs, keep apply moisturizing item for helping regeneration.

GANA TOX

(FDA registered spraying peptide)

FDA VCRP Registration number : 1007236

GANA TOX : CPNP number 3028590

GANA TOX (spraying argireline)

- Argireline and PDRN solution
- Using air spray gun (spraying skin argireline)
- Do not use this products as mesotherapy
- 5ml vials × 10EA/box

GANA TOX include

- PDRN (polydeoxyribobucleotide)
- 3 kinds of peptides
 - Argireline-acetyl hexapeptide-8
 - Palmitoyl tetra peptide 7
 - Palmitoyl pentapeptide 4
- Adenosine
- Hyaluronic acid
- Glutathione
- Niacinamide
- Beta glucan

GANA TOX indication

- Effective for
- Cell healing
- Wrinkle removal
- Skin regeneration
- Face rejuvenation
- Scar removal
- Whitening



5ml vials × 10EA/box

GANA TOX vs Existing peptide products

	Argireline	PDRN	Adenosine
GANA TOX	100 ppm	Included	Included
Existing products	50 ppm	None	None
	Key effect for wrinkle removal	Cell healing	

Argireline

Argireline is made up of peptides, which are chains of amino acids that can affect how cells in the body work. When Argireline is applied to the skin, it is absorbed into the body and the cream's chemical composition causes the facial muscles to contract with less force. Argireline prevents the formation of expression wrinkles by somehow inhibiting muscle movement.

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process. The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation. Most importantly, PDRN provides the nucleotides necessary for cell regeneration. In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries.

Adenosine

In Anti-Aging Skin Care. Adenosine Triphosphate, also referred to as ATP, is a type of molecule that is responsible for being the leading source of energy for the majority of our cellular and muscular functions. It is capable of boosting the levels of proteins such as Collagen and Elastin in the body.

Palmitoyl tetrapeptide-7

Palmitoyl tetrapeptide-7 is a type of synthetic peptide compound which combines several chains of amino acids to reduce skin inflammation and the accompanying skin damage. This peptide compound is also believed to stimulate the regeneration of collagen fibers in the dermis by acting as sort of cellular messenger.

Palmitoyl tetrapeptide-7(Pal-GQPR) is. (It was also formerly known and marketed as palmitoyl tetrapeptide-3.)

Palmitoyl tetrapeptide-7 consists of a short chain of four amino acids (a.k.a. GQPR peptide or glycine-glutamine-proline-arginine) connected to palmitic acid. Palmitic acid is a fatty acid added to improve the peptide's oil solubility and thus skin penetration.

Palmitoyl tetrapeptide-7 is believed to work by reducing the production of interleukin-6 (IL-6) by the the key skin cells, keratinocytes and fibroblasts.

IL-6 is a molecule that promotes inflammation, which, in turn, leads to faster degradation of the skin matrix and thus contributes to the development of wrinkles and loss of skin firmness and elasticity.

By reducing the levels of IL-6 and possibly other inflammation mediators, palmitoyl tetrapeptide-7 is thought to slow down the degradation of the skin matrix and may also stimulate its replenishment.

Palmitoyl Pentapeptide-4

Pentapeptide-4 activates certain genes involved in the process of extracellular matrix renewal and cell proliferation.

Its 16-carbon aliphatic chain is for improving the penetration of the molecule through the lipid structures of the skin.

Stimulates collagen I, III and VI fibronectin, elastin, glucosaminoglycans production.

Palmitoyl Pentapeptide is a compound that is derived from a five-part chain of amino acids known as pro-collagen I pentapeptide.

By itself, this amino chain has been shown to increase the production of collagen in laboratory cultures.

Palmitoyl pentapeptide is a synthetic derivative of this five-part amino acid chain, and it is normally marketed under the brand name Matrixyl, which is simply a combination of palmitic acids and amino-acid chains.

Functionally, Palmitoyl Pentapeptide promotes the production of two compounds: collagen and hyaluronic acid. Combined, these compounds help the skin recover and heal from wounds and penetrative damage by generating new specialized skin cells that bind the damaged tissue together.

How to use

Topical apply

1. Half vial is for full face
One vial is for full face and neck
2. Apply GANA TOX before go to bed
3. Interval 2-3 times a week
4. 5-6 session is needed
5. You will see the effect after 5th session

- Use air brush gun
Or various derma roller
Needle length not longer than 0.5mm



Open GANA TOX vial



Pour GANA TOX in the cup of air spray gun



Turn on the air spray gun



Push the button of the air spray gun



GANA PNV (PDRN vial type)



3ml vials × 5EA/box

GANA PNV is low PDRN concentration solution, prefilled in vials

Major component

- PDRN (poly deoxyribonucleotide from salmon sperm) 0.5%
- Glutathione 2mg/cc
- Cross linked Hyaluronic acid 5mg/cc

GANA PNV is mesotherapysolution, prefilled in vials

PDRN

- Help all kinds of wound healing,
- Regeneration of cell and tissue

Glutathione

- Whitening

Cross linked Hyaluronic acid 5mg/cc

- Prevent water loss
- Keep skin moisturized

PDRN (polydeoxyribonucleotide)

Definition

PDRN is the low molecular weight DNA complex fragments from salmon or trout sperm. It play a role in anti inflammatory, cell proliferation and tissue remodeling.



4 kinds of PDRN products

	Volume	Package	PDRN %	HA concentration	Cross linking HA	Glutathione
GANA PN	2.5ml × 2EA	Syringes	2%	10mg/cc	Yes	2mg/cc
GANA PNS	1.2ml × 1EA	Syringe	2%	20mg/cc	Yes	None
GANA PNV	3ml × 5EA	Vials	0.5%	5mg/cc	Yes	2mg/cc
GANA PNV plus	3ml × 5EA	Vials	2%	10mg/cc	Yes	2mg/cc

GANA PNV vs Existing products

	Effect	Existing products	GANA PNV
PDRN content	Wound healing Cell regeneration	0.2%	0.5%
Lidocaine	Pain	None	None
Glutathione	Whitening	None	2mg/cc
Cross linked hyaluronic acid	Duration 6month	None	10mg/cc

Application

- Face rejuvenation
- Acne scar
- Wrinkle removal
- Whitening
- Preventing sagging
- Improving elasticity, turgidity and tonicity of skin
- Repairing damaged dermal cells

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process. The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation. Most importantly, PDRN provides the nucleotides necessary for cell regeneration.

In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries

How to use and various application

1. Topical apply

- It can be used for topical apply for skin rejuvenation
- Wash face, apply one vial on face and go to bed

After 3 sessions, you will see effect

2. Derma roller

- Use 0.2mm to 0.5mm needle length derma roller
- Interval 1wk
- 3 sessions

Preparation

Apply ice bad and make anesthesia with lidocaine cream both to reduce pain

Precaution

This product contain

Cross linked hyaluronic acid (HA)
Not non cross linked hyaluronic acid

Once injected to epidermis

This can cause nodule and papule
So injection layer will be dermis layer
And use mesotherapy gun .needle depth more than 1.5mm

Do not inject at one point too much
Inject widely deeply and evenl

After injection

Compress treated area with dry gauze

Contraindication

1. Any kinds of active skin disease
2. Any kinds of general illness
3. Hypersensitive to this products
4. Bleeding disorder
5. Pregnancy, breast feeding

Precaution

- This products contain cross linked hyaluronic acid
- Be careful not to penetrate artery
- Cross linked HA can cause arterial occlusion

Side effect

- Bruising
- Hematoma
- Atrophy & Skin hypertrophy
- Erythema
- Urticaria

Post procedure care

1. Apply mask pack for skin suppression
2. Avoid direct sun light
3. In case of mild pain, patients can take NSAID




Before and after

One week after first session



Before and after



REGEN PEEL

(TCA peeling)

Ingredient

REGEN PEEL

30% Trichloroacetic acid
purified water

Post peel neutralizer

7% sodium bicarbonate
Purified water

Warning

- This is a strongest TCA peeling
- The beginner never use it
- This is only for dermatologists

Explanation

This is a professional grade TCA peel with an unbuffered pH of 1.3. TCA is the peel that all other peels are compared to! It has been around for decades, is easy to use, and very predictable. Most dermatologists prefer to use TCA over other acids due to its effectiveness and dependability. If you are looking for long-lasting, dramatic changes in your skin, TCA is the correct choice for you.

We do NOT EVER recommend the direct application of this undiluted solution as a facial peel.

TCA is a highly regarded acid and is perfect for treating common issues such as:

- Acne
- Skin Tone Irregularities
- Texture Problems
- Age Spots
- Fine Wrinkles
- Scarring and Acne Scars



REGEN PEEL 30ml
30% Trichloroacetic acid

Post peel neutralizer 50ml
7% NAHCO3

This peel is very easy to use. You simply cleanse, apply, time it, neutralize and rinse. We include very clear instructions on how to use this peel.

After you have applied the peel, your skin may be red, this is normal, and you may wear makeup to tone in down if you would like. About 2 days after the peel your skin will start to dry out and look leathery, brownish and aged, but don't panic, this is what we want! Your skin is preparing to shed off, and that will occur over the next 3-5 days for most people. Depending on the depth of peel, you may want to hide out some during this time as the peeling can be heavy. We call this down time. You can encourage peeling by using warm compresses and taking hot showers, but you cannot pick at the skin! It must come off when it is ready to, not when you are ready for it to!

By peeling off the skin you risk scarring by exposing skin that is not ready for the world yet, so allow the skin to shed on its own. As with all exfoliating products, the use of sunscreen is essential to obtaining the results you are looking for.

If you are not able to commit to using a sunscreen, this is not the treatment you should choose as you will just damage your skin all over again!

Please Read

Chemical peels should not be used on any sensitive areas of the body, including but not limited to the underarms and groin/genitals.

- People who are not candidates for Chemical peels
- Pregnant or Breast Feeding
- Have Herpes
- Using Accutane
- Are immune impaired
- Have had problems with atypical scarring in the past (keloid, etc)

This is a medical quality peel, and by purchasing this item you are accepting all responsibility for its use. You may need to contact a professional for your particular skin concern. Misuse of chemical products can cause injury and burns.

How to use

- Trichloroacetic acid (TCA), is applied to skin for chemical peels.
- During a peel, TCA is applied to your face for one minutes to remove the outer layers of the skin and reveal fresh layers of skin underneath.
- TCA peel treats surface wrinkles, skin pigment problems and blemishes.
- TCA peels with acid concentrations of 30 percent or more are usually applied in a doctor's office.

Step 1

Test the TCA on a small patch of skin before applying it to your face. Wait 24 hours to see if you tolerate the peel.

Step 2

Apply a thin layer of TCA to your skin with a facial brush. Avoid the eye and lip areas.

Step 3

Neutralize regen peel with neutralizer. In a minute Time is important.

Step 4

Apply a generous amount of post-peel neutralizer to your skin. Cover every area on which you applied TCA to ensure that you neutralize the chemical.

Step 5

Wash your face with cleanser and apply antibiotic ointment to your skin for the first 48 hours after your peel.

Step 6

Protect your skin with a sunblock with SPF 30 or higher for a week after your peel.



Things You'll Need

- Skin cleanser
- Latex gloves
- Facial brush
- Kitchen timer
- Post-peel neutralizer
- Antibiotic ointment
- SPF 30 sunblock



GANA Cocktail

(Whitening)



Specially made for only whitening!

This is solution for dermaroller or mesotherapy

Glutathione : 4ml × 5vials
Vitamin C (ascorbic acid) : 4ml × 5vials

Ingredients

- Glutathione : 4ml × 5vials
- Vitamin C (ascorbic acid) : 4ml × 5Vials

Attention

Never mix Glutathione and Vitamin C

Glutathione is a low molecular weight thiol-tripeptide that plays a prominent role in maintaining intracellular redox balance. In addition to its remarkable antioxidant properties, the discovery of its antimelanogenic properties has led to its promotion as a skin-lightening agent. It is widely used for this indication in some ethnic populations.

Glutathione is present intracellularly in its reduced form and plays an important role in various physiological functions. Its skin-lightening effects result from direct as well as indirect inhibition of the tyrosinase enzyme and switching from eumelanin to pheomelanin production. It is available in oral, parenteral and topical forms.

Vitamin C, a revitalizing & antioxidant solution.

A very effective antioxidant that neutralizes free radicals—the unstable molecules that age the skin as a result of pollution, smoking and exposure to the sun. It inhibits melanin, stimulates growth of the hydro-lipid film and enhances collagen synthesis, an important skin protein that strengthens the structural support and elasticity of the skin.

Vitamin C Revitalizing & Antioxidant Solution is used mainly to treat and prevent photoageing, uneven complexions and for helping the skin to fight wrinkles. Ascorbic acid is one of the main non-enzymatic, water-soluble antioxidant. Vitamin C is required for the synthesis of collagen, an important structural component of blood vessels, tendons, ligaments and bones.

In addition, vitamin C is required for the synthesis of carnitine, a small molecule that is essential for the transport of fat to cellular organelles called mitochondria, for conversion to energy.

It plays a critical role in dermal construction, stimulating collagen synthesis of dermal fibroblasts.

This is thought to occur as a result of ascorbate directly stimulating collagen synthesis, directly and specifically activating collagen gene regulation both by increasing the transcription rate and stabilizing procollagen messenger RNA, thus genetically signaling collagen synthesis.

Another mechanism is the initiation of lipid peroxidation, which leads to an increase in a by-product, which stimulates collagen gene expression.

In addition, ascorbic acid is necessary to form lysyl hydroxylase an enzyme necessary for cross-linking one collagen molecule to another providing tissue strength.

The topical use of ascorbic acid prevents and treats photo-ageing.

UV light exposure depletes up to two thirds of cutaneous ascorbic acid stores.

Caution

Glutathione and Vitamin C can not be mixed for use

How to use

- Apply biweekly
- Glutathione - 1st 3rd 5th week
- Ascorbic acid - 2nd 4th 6th week



GANA SCALP

(Hair care)

PDRN × 5 vials

GHK_CU × 5 vials



3ml × 10EA
- use for hair growth

Caution

- Do not use for other use
- Do not use on the face and neck

Ingredients

- PDRN (poly deoxyribonucleotide)
- Copper peptide

Effect of GANA SCALP

1. Scar healing by Copper peptide
2. Hair growth by copper peptide
3. Cell regeneration by PDRN and copper peptide
4. Blood vessel regrowth by PDRN and copper peptide

What is copper peptide

Copper peptide GHK-Cu is a naturally occurring copper complex of a glycyl-L-histidyl-L-lysine peptide. Since it has three amino acids it is called tripeptide. The GHK-Cu tripeptide has strong affinity for copper(II) and was first isolated from human plasma. It can be found also in saliva and urine.

Copper peptides are naturally occurring small protein fragments that have high affinity to copper ions. In human plasma, the level of GHK-Cu is about 200 µg/ml at age 20. By the age of 60, the level drops to 80 µg/ml.

In humans, tripeptide GHK-Cu can promote activation of wound healing, attraction of immune cells, antioxidant and anti-inflammatory effects, stimulation of collagen and glycosaminoglycan synthesis in skin fibroblasts and promotion of blood vessels growth.

Recent studies revealed its ability to modulate expression of a large number of human genes, generally reversing gene expression to a healthier state. Synthetic GHK-Cu is used in cosmetics as a reparative and anti-aging ingredient.

Cosmetic use

Facial studies

Copper peptide GHK-Cu is widely used in anti-aging cosmetics (INCI name : Copper tripeptide-1).

Several controlled facial studies confirmed anti-aging, firming and anti-wrinkle activity of copper peptide GHK-Cu.

Abdulghani et al. established that facial cream containing GHK-Cu increased collagen in photoaged skin of 20 female volunteers, performing better than vitamin C and retinoic acid.

Leyden et al. conducted a 12-week facial study of GHK-Cu containing face and eye cream, reporting significant improvement of skin laxity, clarity and appearance, reduced fine lines and the depths of wrinkles and increased skin density and thickness comparing to placebo.

GHK-Cu eye cream performed better than vitamin K cream.

Finkley et al. conducted 12 week facial study on 67 women and reported that GHK-Cu cream applied twice daily improved aged skin appearance, increased thickness, reduced wrinkles and strongly stimulated dermal keratinocyte proliferation as determined by histological analysis of biopsies. The same study found copper peptide GHK-Cu to be non-toxic and non-irritating.

Hair growth

Copper peptide GHK-Cu and its analogues were found to stimulate hair growth. In some circumstances, the efficiency of synthetic analog of GHK-Cu was similar to that of 5% minoxidil.

A commercial product GraftCyte was clinically proven to improve hair transplantation outcome.

Shown to promote collagen production, using copper peptides topically on the scalp will help strengthen already existing hair, while stimulating growth in areas that are lacking thickness.

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process.

The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation. Most importantly, PDRN provides the nucleotides necessary for cell regeneration.

In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries

How to use

1. Use with derma roller
2. Interval 2wks
3. PDRN × 5 vials
GHK_CU × 5 vials
Apply each vials in turn biweekly
4. Total 10 sessions



GANA PNV + (high PDRN)



3ml vials × 5EA/box

GANA PNV plus : CPNP number 3030186

**GANA PNV plus is high PDRN concentration solution,
prefilled in vials**

Major component

- PDRN (poly deoxyribonucleotide from salmon sperm) 2%
- Glutathione 2mg/cc
- Cross linked Hyaluronic acid 10mg/cc
- GANA PNV plus is mesotherapysolution, prefilled in vials

PDRN

- Help all kinds of wound healing,
- Regeneration of cell and tissue

Glutathione

- Whitening

Cross linked Hyaluronic acid 10mg/cc

- Prevent water loss
- Keep skin moisturized

PDRN(polydeoxyribonucleotide)

Definition

PDRN is the low molecular weight DNA complex fragments from salmon or trout sperm. It play a role in anti inflammatory, cell proliferation and tissue remodeling.



4 kinds of PDRN products

	Volume	Package	PDRN %	HA concentration	Cross linking HA	Glutathione
GANA PN	2.5ml × 2EA	Syringes	2%	10mg/cc	Yes	2mg/cc
GANA PNS	2.5ml × 1EA	Syringe	2%	20mg/cc	Yes	None
GANA PNV	3ml × 5EA	Vials	0.5%	5mg/cc	Yes	2mg/cc
GANA PNV plus	3ml × 5EA	Vials	2%	10mg/cc	Yes	2mg/cc

GANA PNV plus vs Existing products

	Effect	Existing products	GANA PNV+
PDRN content	Wound healing Cell regeneration	0.2%	2.0%
Glutathione	Whitening	None	2mg/cc
Cross linked hyaluronic acid	Duration 6month	None	10mg/cc

Application

- Face rejuvenation
- Acne scar
- Wrinkle removal
- Whitening
- Preventing sagging
- Improving elasticity, turgidity and tonicity of skin
- Repairing damaged dermal cells

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process. The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation. Most importantly, PDRN provides the nucleotides necessary for cell regeneration.

In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries.

How to use and various application

1. Topical apply

- It can be used for topical apply for skin rejuvenation
- Wash face, apply one vial on face and go to bed
- One vial for each session

After 3 sessions, you will see effect

2. Derma roller

- Use 0.2mm to 0.5mm needle length derma roller
- Interval 1wk
- 3 sessions
- One vial for each session
- Dose : cheek 5 to 15ml
forehead 5 to 10ml
second session in a month needed

Preparation

Apply ice bad and make anesthesia with lidocaine cream both to reduce pain

Precaution

This product contain

Cross linked hyaluronic acid (HA)
Not non cross linked hyaluronic acid

Once injected to epidermis

This can cause nodule and papule
So injection layer will be dermis layer
And use mesotherapy gun. needle depth more than 1.5mm


Do not inject at one point too much
Inject widely deeply and evenly

After injection

Compress treated area with dry gauze

Procedure with mesotherapy gun

needle depth 1.5mm above
mesotherapy gun setting is below picture



Contraindication

1. Any kinds of active skin disease
2. Any kinds of general illness
3. Hypersensitive to this products
4. Bleeding disorder
5. Pregnancy, breast feeding

Precaution

- This products contain cross linked hyaluronic acid
- Be careful not to penetrate artery
- Cross linked HA can cause arterial occlusion

Side effect


- Bruising
- Hematoma
- Atrophy & Skin hypertrophy
- Erythema
- Urticaria



Use Mesogun.
MTS or enclosed 30G needles



Injection layer : dermis



CASE 1

Used GANA PNV plus

(=same components as GANA PN)

Skin tone and skin irregularity were improved

Procedured by Dr Vita Mastrolonardo

Aesthetic and Anesthesologist Dr Importaciones CruzAng

Before and after



CASE 2

Used GANA PNV plus

(=same components as GANA PN)

Sagging jaw line

Skin irregularity on cheek

much improved

Procedured by Dr. Oxmerari Chirivella

Plastic Surgeon of the Venezuelan Society of Plastic Surgeon
(SVCPREM)

Importaciones CruzAng Venezuela

Before and after



GANA EYE

(Eye bag removal)



2ml vials × 5EA in a box

**Specially made for removing eye bag,
dark circle, only!**

Main components

- Carnitine : 2%
- Acetyl tetrapeptide-5 : 50ppm
- Phosphate buffer solution-q.s

Packaging

- 2ml/5ml prefilled vials × 5EA in a box
One box GANA EYE can be used for more than 20 sessions

GANA DM **(PLLA meso)**



Freeze dried Poly L lactic acid powder **induce collagen synthesis in human body**

- Cross linked HA make skin moisturized more than 6 months
- Conventional HA meso products are non cross linked HA, non cross linked HA last in human body just 1-2wks,

GANA DM set
 - Composed of freeze dried PLLA powder (50mg/5ml)
 - Cross linked hyaluronic acid (5ml/5ml)

What is PLLA

- PLLA is a powerful collagen stimulator in human body
- PLLA induce collagen synthesis in 6 weeks
- Once collagen made it last more than 5 years
- Newly formed collagen make volume for depressed area
- And induce lifting by tightening of face
- So newly formed PLLA recover depressed area, like acne scar, stretch marks

Poly(L-lactide) is a biodegradable material with good bio-absorbability and biocompatibility. After degradation in the body, PLLA products are excreted by metabolism and thus has no harmful side effects to the human body. It is for these reasons that PLLA products are currently widely used in the field of medicine, such as for one-time infusion equipment, medical surgical suture, drug delivery carriers, tissue engineering scaffolds, bone anchors, microcapsules, microspheres, implants, elastomer materials for organ support, etc.

The characteristics of the PLLA and hyaluronic acid in Gana dualmesosolution

1. The PLLA in Gana DM
 - The size of PLLA particles in the conventional products is average 30 to 50um size.
2. The hyaluronic acid in GANA DM
 - 1) Conventional mesotherapy products only contain non cross linked hyaluronic acid but GANA DM contain cross linked hyaluronic acid last more than 6 months
 - 2) Safety, cross linked hyaluronic acid can be autoclaved—contain no preservative
 - 3) Cross linked HA can be used like dermal filler—give us immediate volume correction

Hyaluronic acid in GANA DM vs Existing mesotherapyproducts

	Conventional	GANA DM
Cross linking ratio %	0 %	5 %
HA concentration (mg/cc)	10	10
Cross linking agent	None	BDDE
Duraron month	1-2 wks	6-12 month
Particle size	None	Under 30um

Application

- Stretch mark (striae)
- Acne scar
- Big pore
- Face rejuvenation
- Face lifting by collagen synthesis
- Neck wrinkle
- Hand wrinkle

After procedure

Skin tone was improved
Fine wrinkle were removed

Before and after**Stretch mark****Before procedure**

Procedure by Russian distributor.

BIOMED co ltd

Stretch mark procedure
After procedure
Only left belly was done
One session
After 6weeks

**Acne scar****Before procedure**

Procedured by Russian distributor.

BIOMED co ltd

After procedure

Acne scar on glabella and cheek was removed a lot


Neck wrinkle**Hand wrinkle**

HAND before

Before and after

How it works

- Freeze dried PLLA has micropore
- Micropore absorb water to break down into small pieces
- The broken PLLA particles induce controlled inflammation
- So that there will be collagen synthesis
- This SEM picture is of our PLLA



Mechanism and technique - II



GANA DM set

PLLA powder for mesotherapy

- Main component is poly L lactic acid
PLLA is collagen stimulator, induce collagenesis by activating fibroblast
- GANA DM set is autoclaved cross linked hyaluronic acid

The merits of GANA DM set. Prefilled vial

Major component

- Cross linked hyaluronic acid gel (longer duration and better efficacy)
- Autoclave sterilized (better safety)

How to use for face rejuvenation

- Mix ganafill hydro with gana ha hydro
- Shake well for 5 minutes
- Use Vortex mixer to mix
- Please see picture below

How to stop Aging process

The older people have lower collagen and lower hyaluronic acid consequently make wrinkle, fold, sagging that make person look old

- GANA FILL hydro make collagen in human body
- GANA HA hydro bind water in skin
- So that two newly formed collagen and hyaluronic acid lower wrinkle fold and sagging
- Consequently make person look younger

Aging

1. Water loss
2. Collagen loss
3. Elasticity loss

- GANA FILL hydro induce collagen
- GANA HA hydro prevent water loss

VORTEX MIXER



How to use for stretch marks, acne scar and neck wrinkle

- Prepare DM set solution
- Interval one month
- Total 2-3 sessions will be needed

GANA PF (Expression wrinkle removal)

GANA PF (peptide filler)

GANA PF is made to remove wrinkle effectively
It contain cross linked HA and peptides good for wrinkle removal



Packing unit : 2ml × 5 vials

Characterized by two main components

- Cross linked HA-main component of dermal filler
- Acetyl hexapeptide-8 : act like botulinum toxin, relaxing facial muscle

Ingredients

- Cross linked hyaluronic acid 15mg/cc
- Acetyl hexapeptide 50ppm
- PBS-qs

Precaution

1. Use only for fine shallow wrinkle
ex) forehead, crow feet, under eye fine wrinkle
2. Do not use this product for deep wrinkle
like nasolabial fold and lips
3. Do not use this products for augmenation
4. Cause of acetyl hexa peptide
The some person might feel paralyzed
facial expression
It is normal expectation
Reassure patients

What is cross linked hyaluronic acid and acetyl hexapeptide

Cross linked hyaluronic acid

Cross linked HA is not normal HA of cosmetics. This is mani components of dermal fillers.

Cross HA is linked by BDDE (cross linker) to stay in human skin more than 6 months up to 18 months.

Restore depressed soft tissue so that to elevate sunken wrinkle and store water in dermis.

Acetyl hexapeptide

Is made up of peptides, which are chains of amino acids that can affect how cells in the body work.

When Acetyl hexapeptide is applied to the skin, it is absorbed into the body and the cream's chemical composition causes the facial muscles to contract with less force.

Acetyl hexapeptide prevents the formation of expression wrinkles by somehow inhibiting muscle movement.

GANA GYNO (Vaginal whitening with vegetable estrogen)



This products is for labia whitening and vaginal dryness also can be used as lubricants

Especially effective for menopausal woman

GANA GYNO 50ml

Ingredients

- Glutathione
- Isoflavone pharmaceutical grade
- Xanthan gum
- Glycerine
- Lactic acid

What is glutathione

Glutathione is the body's most powerful antioxidant, unknown by most. It is popular when used as a way to help whiten skin though. Glutathione skin whitening is growing in popularity, helping people achieve the skin pigmentation they desire with no harmful side effects when taken in proper dosages.

What is Isoflavone

1) Soyisoflavones is a standardized natural extract containing soy isoflavone aglycones, genistein and daidzein. It has anti-aging and antioxidants benefits. Soy isoflavones exhibit estrogen-like properties that may help maintain skin texture and integrity, prevent wrinkles and stimulate hyaluronic acid. The antioxidant effects of soy isoflavones are also useful in protecting the skin against photo damage thereby preventing aging, as well as reducing the appearance of current age spots. The estrogen-like qualities of this product can also be used in hair care products to enhance hair regrowth.

2) Isoflavone vaginal gel is effective for relief of vaginal dryness and dyspareunia symptoms and an increase in the intermediate and superficial cells was noted. These results were similar to the effects with use of estrogens and superior to placebo gel.

GANA TM

(FDA registered mask pack)

(Facial mask) FDA registered

FDA VCRP REGISTRATION NUMBER : 1007236

Effects

Very effective for

1. Skin rejuvenation by PDRN (salmon sperm DNA)
2. Mild and non irritable peeling by PHA
3. Wrinkle removal and face lifting by acetyl hexapeptide-8
4. Immediate hydration and skin defect recovering
by Cross linked hyaluronic acid

Main ingredients

- PDRN (poly deoxy ribo nucleotide)
- PHA (poly hydroxy acid)
- Acetyl hexapeptide-8
(similar function to botulinum toxin)
- Cross linked hyaluronic acid
(main components of dermal fillers)

Detail ingredients

- Lactobionic acid(=PHA)
- Acetyl hexapeptide-8(=arigirelline)
- Alpha arbutin(goof for whitening)
- Adenosin
- Palmitoyl tetrapeptide7
- Palmitoyl pentapeptide 4
- 1.2 hexandiol
- Bio cellulose
- Cross linked hyaluronic acid




3 mask packs/box



Attention and How to use

GANA TM mask packs are three layered bio cellulose remove upper and lower layers and use middle bio cellulose



GANA TM is completely differently from conventional products so when you use GANA TM you must know below

1. You can feel skin tightened just after apply.
It is because GANA TM has acetyl hexapeptide-8 in large amounts hexapeptide-8 directly relax facial muscles
2. After using GANA TM
You might feel skin dried but polyhydroxy acid make moistness for long time.
So after 3 sessions, roughly one week
You can feel moistness
3. After removing GANA TM
Do not apply any moisture cream or lotion.
It will prohibit GANA TM effect
4. Before use, wash face and apply GANA TM for 10 minutes. Then go to bed
5. One GANA TM has 3 packs.
Use three packs in a week
6. The effect will be seen after at least 3 packs
7. Main effect of GANA TM are
 - removing fine wrinkle, face lifting for 4-6 months by acetyl hexapeptide-8
 - hydration and skin tone improvement by polyhydroxy acid
 - damaged skin repair by PDRN
 - long lasting hydration by cross linked hyaluronic acid
8. Cause of peeling effect of polyhydroxy acid (PHA)
You might feel little irritated skin and slight redness for a hour
But it is normal process of soft peeling so you don't have to concern

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process. The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation. Most importantly, PDRN provides the nucleotides necessary for cell regeneration. In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries.

Cross linked hyaluronic acid

Cross linked HA is not normal HA of cosmetics, This is main components of dermal fillers. Cross HA is linked by BDDE (cross linker) to stay in human skin more than 6 months up to 18 months. Restore depressed soft tissue so that to elevate sunken wrinkle and store water in dermis.

Safety of PHA

Even very sensitive skin types can tolerate PHAs, including atopic and rosacea skin. It is thought that the gentle, nonirritating action of PHAs may be partially related to a larger molecular composition than that of conventional AHAs. Larger molecules are believed to penetrate the skin at a slower rate, causing less irritation. This reduced irritation, coupled with natural antioxidant properties, gives PHAs a gentle, competitive edge over other anti-aging treatments.

What is PHA (Exfoliant, Antioxidants)

Polyhydroxy acids are ingredients such as gluconolactone and lactobionic acid. They're supposed to be as effective as AHAs, but less sensitizing. Gluconolactone and lactobionic acid are chemically and functionally similar to AHAs (such as glycolic acid).

The significant difference between them and AHAs is that gluconolactone and lactobionic acid have larger molecular structures, which limits their ability to penetrate the skin, resulting in less potential for sensitizing side-effects. Supposedly, this reduced absorption into the skin does not hamper their effectiveness. Does that mean gluconolactone and lactobionic acid are better for your skin than AHAs in the form of glycolic acid or lactic acid?

According to an Internet-published class lecture by Dr. Mark G. Rubin, a board-certified dermatologist and assistant clinical professor of dermatology at the University of California, San Diego, research on gluconolactone demonstrated only a "6% decrease in dermal penetration" in comparison to glycolic acid, which "isn't a dramatic improvement."

Gluconolactone may be slightly less sensitizing for some skin types, but this isn't the magic bullet for exfoliation some cosmetics companies have been extolling.

Polyhydroxy acids can also function as antioxidants and may promote some amount of improvement in skin's surface strength

- PHAs plus hydroquinone showed excellent improvement in antiaging and skin lightening parameters.
- Finally, PHA-containing products were shown to be compatible with African American, Caucasian, and Hispanic/Asian skin and provided significant improvements in photoaging in these populations.

PHA face peel is indicated for :

- Acne-prone skin
- Comedones
- Enlarged pores
- Hyperpigmentation and uneven skin tone
- Skin aging prevention
- Early signs of photoaging such as skin dryness
- Loss of elasticity and fine lines
- Hyperkeratosis (Rosacea, seborrheic dermatitis)
- Actinic dermatitis
- Correction and prevention of natural aging signs such as withering and first wrinkles pigmentation

PHA effect

- PHAs have been found to be compatible with clinically sensitive skin, including rosacea and atopic dermatitis, and can be used after cosmetic procedures.
- PHAs provide additional humectant and moisturization properties compared with AHAs and can enhance stratum corneum barrier function, therefore increasing the skin's resistance to chemical challenge.
- Most PHAs also possess antioxidant properties.
- PHAs such as gluconolactone or lactobionic acid may be used in combination with other products, ingredients, or procedures such as laser and microdermabrasion to provide additional benefits to therapy or to enhance the therapeutic effect.
- Several studies were conducted in support of this, and methods and results are discussed. In summary, PHA-containing products were used in combination with retinoic acid in treating adult facial acne and were found to be well tolerated.
- PHAs plus retinylacetate (pro-vitamin A) in a cream base exhibited significant antiaging skin benefits such as skin smoothing and plumping.

Benefit of PHA

- Promote cell turnover. By sloughing off dead skin cells, revealing smoother, healthy new cells underneath.
- PHAs have humectant effects that allow for increased moisturisation. When skin is sufficiently moisturised, it looks smoother and healthier.
- Enhancing skin smoothness and elasticity.
- Antioxidant effects. Gluconolactone is an antioxidant known for its free radical-fighting ability, protecting skin from harmful UV exposure.
- Increased skin barrier function. Our skin barrier, or stratum corneum, plays a vital role in keeping out irritating chemicals and substances.
- Improving skin clarity and brightness; reducing the appearance of fine lines, wrinkles, and pore size
- Reduce flare-ups in inflammatory conditions. PHA products can lower inflammatory episodes of acne, rosacea, dermatitis, psoriasis and melasma

Acetyl hexapeptide-8, Argireline

Argireline is made up of peptides, which are chains of amino acids that can affect how cells in the body work.

When Argireline is applied to the skin, it is absorbed into the body and the cream's chemical composition causes the facial muscles to contract with less force.

Argireline prevents the formation of expression wrinkles by somehow inhibiting muscle movement.

Adenosine

In Anti-Aging Skin Care.

Adenosine Triphosphate, also referred to as ATP, is a type of molecule that is responsible for being the leading source of energy for the majority of our cellular and muscular functions.

It is capable of boosting the levels of proteins such as Collagen and Elastin in the body.

Palmitoyl tetrapeptide-7

Is a type of synthetic peptide compound which combines several chains of amino acids to reduce skin inflammation and the accompanying skin damage.

This peptide compound is also believed to stimulate the regeneration of collagen fibers in the dermis by acting as sort of cellular messenger.

Palmitoyl tetrapeptide-7 (Pal-GQPR) is.

(It was also formerly known and marketed as palmitoyl tetrapeptide-3.)

Palmitoyl tetrapeptide-7 consists of a short chain of four amino acids (a.k.a. GQPR peptide or glycine-glutamine-proline-arginine) connected to palmitic acid.

Palmitic acid is a fatty acid added to improve the peptide's oil solubility and thus skin penetration.

Palmitoyl tetrapeptide-7 is believed to work by reducing the production of interleukin-6 (IL-6) by the the key skin cells, keratinocytes and fibroblasts.

IL-6 is a molecule that promotes inflammation, which, in turn, leads to faster degradation of the skin matrix and thus contributes to the development of wrinkles and loss of skin firmness and elasticity.

By reducing the levels of IL-6 and possibly other inflammation mediators, palmitoyl tetrapeptide-7 is thought to slow down the degradation of the skin matrix and may also stimulate its replenishment.

Palmitoyl Pentapeptide-4

Pentapeptide-4 activates certain genes involved in the process of extracellular matrix renewal and cell proliferation.

Its 16-carbon aliphatic chain is for improving the penetration of the molecule through the lipid structures of the skin.

Stimulates collagen I, III and VI fibronectin, elastin, glucosaminoglycans production.

Palmitoyl Pentapeptide is a compound that is derived from a five-part chain of amino acids known as pro-collagen I pentapeptide. By itself, this amino chain has been shown to increase the production of collagen in laboratory cultures.

Palmitoyl pentapeptide is a synthetic derivative of this five-part amino acid chain, and it is normally marketed under the brand name Matrixyl, which is simply a combination of palmitic acids and amino-acid chains.

Functionally, Palmitoyl Pentapeptide promotes the production of two compounds: collagen and hyaluronic acid.

Combined, these compounds help the skin recover and heal from wounds and penetrative damage by generating new specialized skin cells that bind the damaged tissue together.



GANA MASK

(Facial burn mask)



Package : one sheet/pack

Development objective

GANA MASK is developed for curing any kinds of burn damage of face including laser therapy and chemical peeling

What is difference,

- As soon as applied to patient GANA MASK can keep moisturizing from evaporation
- Prevent infection
- Mechanical-chemical protection
- Reduce pain
 - PLLA Powder in GANA MASK stimulate the collagen synthesis of soft tissue defect
- High concentration Cross linked hyaluronic acid gel absorb watery content from wound and fill defected skin area.
- PDRN in GANA MASK help to heal wounded skin

Main component of GANA MASK

1. Bio cellulose
2. Poly-L-lactic acid. PLLA
3. Cross linked hyaluronic acid
4. PDRN (polydeoxyribonucleotide from salmon sperm)



GANA MASK

Introduction of contents

BIO Cellulose Sheet

Application field of sheet

- Applicable plastic surgery, rhino, genital augmentation.
- In case of surgical operation substitute for diaphragm.
- The fascia from every part of the body or peritoneum can be replaced.

*** Bio Implantable Cellulose sheet itself works as existing Alloderm.**

- Microbial cellulose (MC) synthesized in abundance by *Acetobacter xylinum* shows vast potential as a novel wound healing system. The high mechanical strength and remarkable physical properties result from the unique nanostructure of the never-dried membrane. This article attempts to briefly summarize the recent developments and applications of MC in the emerging field of novel wound dressings and skin substitutes. It considers the properties of the synthesized material, its clinical performance, as well as progress in the commercialization of MC for wound care products. Efficient and inexpensive fermentation techniques, not presently available, will be necessary to produce large quantities of the polymer.

Epithelium and Soft tissues regeneration

- The function of GANA MASK powder (PLLA, PDRN)
- Improve soft tissue regeneration
- Improve epidermis regeneration
- Absorb exudate
- In case of major burns is accompanied by soft tissue loss and epithelium damage. The frozen dried powder has functions for epithelium and soft tissue regeneration. It can be applied on burn injury before applying the sheet is soaked gel up.

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process. The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation.

Most importantly, PDRN provides the nucleotides necessary for cell regeneration. In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries.

It's still a very new treatment that is being explored in clinical tests.

Hyaluronic acid (HA;conjugate base hyaluronate),

also called hyaluronan, is an anionic, non sulfated glycosaminoglycan distributed widely throughout connective, epithelial, and neural tissues.

It is unique among glycosaminoglycans in that it is nonsulfated, forms in the plasma membrane instead of the Golgi apparatus, and can be very large, with its molecular weight often reaching the millions.

One of the chief components of the extracellular matrix, hyaluronan contributes significantly to cell proliferation and migration, and may also be involved in the progression of some malignant tumors.

The average 70 kg(154 lb) person has roughly 15 grams of hyaluronan in the body, one-third of which is turned over (degraded and synthesized) every day.


Hyaluronic acid is also a component of the group A streptococcal extracellular capsule, and is believed to play a role in virulence.

Poly(L-lactide) is a biodegradable material with good bio-absorbability and bio-compatibility.


After degradation in the body, PLLA products are excreted by metabolism and thus has no harmful side effects to the human body.

It is for these reasons that PLLA products are currently widely used in the field of medicine, such as for one-time infusion equipment, medical surgical suture, drug delivery carriers, tissue engineering scaffolds, bone anchors, injectable microcapsules, microspheres, implants, elastomer materials for organ support, etc.


Mechanism



PLLA degradation



Mechanism and technique - II



How to use

1. Clean face
2. Apply GANA MASK for 20min
3. Remove pack
4. Do not wash face after using GANA MASK



GANA TC plus

(PDRN, PHA.AH8 ointments)

Effects

Very effective for

1. Skin rejuvenation by PDRN (salmon sperm DNA)
2. Mild and non irritable peeling by PHA
3. Wrinkle removal and face lifting by acetyl hexapeptide-8
4. Immediate hydration and skin defect recovering by Cross linked hyaluronic acid



Tube type - 45gram

Main ingredients

- PDRN (poly deoxy ribo nucleotide) 2%
- Lactobionic acid 2%
- Acetyl hexapeptide-8 (similar function to botulinum toxin) 50ppm
- Cross linked hyaluronic acid (main components of dermal fillers) 1%

	PHA	HA	Acetyl hexapeptide-8	PDRN
GANA TC plus	2 %	1 %	50 ppm	2 %
GANA TC	0.5 %	1 %	25 ppm	0.5 %

Acetyl hexapeptide-8, Argireline

Argireline is made up of peptides, which are chains of amino acids that can affect how cells in the body work. When Argireline is applied to the skin, it is absorbed into the body and the cream's chemical composition causes the facial muscles to contract with less force.

Argireline prevents the formation of expression wrinkles by somehow inhibiting muscle movement.

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process.

The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation.

Most importantly, PDRN provides the nucleotides necessary for cell regeneration. In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries

Cross linked hyaluronic acid

Cross linked HA is not normal HA of cosmetics. This is many components of dermal fillers.

Cross HA is linked by BDDE (cross linker) to stay in human skin more than 6 months up to 18 months.

Restore depressed soft tissue so that to elevate sunken wrinkle and store water in dermis.

What is PHA (Exfoliant, Antioxidants)

Polyhydroxy acids are ingredients such as gluconolactone and lactobionic acid. They're supposed to be as effective as AHAs, but less sensitizing.

Gluconolactone and lactobionic acid are chemically and functionally similar to AHAs (such as glycolic acid).

The significant difference between them and AHAs is that gluconolactone and lactobionic acid have larger molecular structures, which limits their ability to penetrate the skin, resulting in less potential for sensitizing side-effects.

Supposedly, this reduced absorption into the skin does not hamper their effectiveness.

Does that mean gluconolactone and lactobionic acid are better for your skin than AHAs in the form of glycolic acid or lactic acid?

According to an Internet-published class lecture by Dr. Mark G. Rubin, a board-certified dermatologist and assistant clinical professor of dermatology at the University of California, San Diego, research on gluconolactone demonstrated only a "6% decrease in dermal penetration" in comparison to glycolic acid, which "isn't a dramatic improvement."

Gluconolactone may be slightly less sensitizing for some skin types, but this isn't the magic bullet for exfoliation some cosmetics companies have been extolling.

Polyhydroxy acids can also function as antioxidants and may promote some amount of improvement in skin's surface strength

GANA TC

(PDRN, PHA.AH8 ointments)

Effects

Very effective for

1. Skin rejuvenation by PDRN (salmon sperm DNA)
2. Mild and non irritable peeling by PHA
3. Wrinkle removal and face lifting by acetyl hexapeptide-8
4. Immediate hydration and skin defect recovering by Cross linked hyaluronic acid



Tube type - 45gram

Main ingredients

- PDRN (poly deoxy ribo nucleotide) 0.5%
- Lactobionic acid 0.5%
- Acetyl hexapeptide-8 (similar function to botulinum toxin) 25ppm
- Cross linked hyaluronic acid (main components of dermal fillers) 1%

	PHA	HA	Acetyl hexapeptide-8	PDRN
GANA TC plus	2 %	1 %	50 ppm	2 %
GANA TC	0.5 %	1 %	25 ppm	0.5 %

Acetyl hexapeptide-8, Argireline

Argireline is made up of peptides, which are chains of amino acids that can affect how cells in the body work. When Argireline is applied to the skin, it is absorbed into the body and the cream's chemical composition causes the facial muscles to contract with less force.

Argireline prevents the formation of expression wrinkles by somehow inhibiting muscle movement.

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process.

The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation.

Most importantly, PDRN provides the nucleotides necessary for cell regeneration. In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries

Cross linked hyaluronic acid

Cross linked HA is not normal HA of cosmetics. This is many components of dermal fillers.

Cross HA is linked by BDDE (cross linker) to stay in human skin more than 6 months up to 18 months.

Restore depressed soft tissue so that to elevate sunken wrinkle and store water in dermis.

What is PHA (Exfoliant, Antioxidants)

Polyhydroxy acids are ingredients such as gluconolactone and lactobionic acid. They're supposed to be as effective as AHAs, but less sensitizing.

Gluconolactone and lactobionic acid are chemically and functionally similar to AHAs (such as glycolic acid).

The significant difference between them and AHAs is that gluconolactone and lactobionic acid have larger molecular structures, which limits their ability to penetrate the skin, resulting in less potential for sensitizing side-effects.

Supposedly, this reduced absorption into the skin does not hamper their effectiveness.

Does that mean gluconolactone and lactobionic acid are better for your skin than AHAs in the form of glycolic acid or lactic acid?

According to an Internet-published class lecture by Dr. Mark G. Rubin, a board-certified dermatologist and assistant clinical professor of dermatology at the University of California, San Diego, research on gluconolactone demonstrated only a "6% decrease in dermal penetration" in comparison to glycolic acid, which "isn't a dramatic improvement."

Gluconolactone may be slightly less sensitizing for some skin types, but this isn't the magic bullet for exfoliation some cosmetics companies have been extolling.

Polyhydroxy acids can also function as antioxidants and may promote some amount of improvement in skin's surface strength

GANA BB

(BB with PDRN ointments)

GANA BB is

BB cream containing PDRN

And made for Caucasian

Color and flavor are customized to european

PDRN is good for skin regeneration

SPF 35



Tube type - 45gram

What Is PDRN?

Polydeoxyribonucleotide is derived from sperm trout through an extraction process.

The compounds are able to stimulate Vascular Endothelial Growth Factors and certain receptors in the body that trigger the creation of new artery formation, improves blood flow, and reduces inflammation. Most importantly, PDRN provides the nucleotides necessary for cell regeneration.

In addition to its use for diabetic foot ulcers, PDRN has proven effective in treating PAOD, chronic wounds, hind limb ischemia, osteoarthritis disease, coronary artery disease, and burn injuries

BB cream is short for 'blemish balm' cream.

It's a versatile combination of products such as moisturizer, primer, SPF, foundation, skin treatment, and concealer. It was originally developed in Germany as an all-in-1 cream to be used by patient's post-laser treatment for protection, coverage, and hydration.

GANA KA

(Kojic acid ointment)

Kojic acid 4% ointment for whitening

Kojic acid is a chelation agent produced by several species of fungi, especially *Aspergillus oryzae*, which has the Japanese common name koji.

Kojic acid is a by-product in the fermentation process of malting rice, for use in the manufacturing of sake, the Japanese rice wine. It is a mild inhibitor of the formation of pigment in plant and animal tissues, and is used in food and cosmetics to preserve or change colors of substances. It forms a bright red complex with ferric ions.



Tube type - 45gram

Effect of kojic acid

1. Whitening

Kojic acid is made from several different types of fungi. It's also a byproduct when certain foods ferment, including Japanese sake, soy sauce, and rice wine. Kojic acid inhibits and prevents the formation of tyrosine, which is an amino acid that's needed to produce melanin. Melanin is the pigment that affects hair, skin, and eye color. Because it inhibits the production of melanin, kojic acid can have a lightening effect.

2. Protective effects from radiation-induced damage

Kojic acid appears to possess marked protective (radioresistance) effects from radiation-induced damage. It increased the 30-day survival rate of mice after exposed to a lethal dose of gamma radiation and had significant radioprotective effects on the hematopoietic system, the immune system and DNA of mice exposed to a 4 Gy sublethal dose of radiation. Pretreatment with kojic acid has been shown to protect Chinese hamster ovary cells against ionizing radiation-induced damage. Dogs pretreated with kojic acid after whole-body exposure to a lethal dose of 3 Gy gamma radiation had a 51-day survival rate of 66.7% versus the dogs in the 3 Gy irradiation only group, which all died within 16 days of postirradiation.

GANA V line

(Face contour improval)

GANA V line

Face contour improvement solution

Ingredients

- Phosphatidylcholine (PPC)
- Deoxycholic acid (DA)
- Acetylhexapeptide-8
- Glutathione
- Benzyl alcohol

Main effect

- Dissolve facial fat by PPC and DA
- Wrinkle removal and face lifting by Acetyl hexapeptide-8
- Whitening by glutathione



Package unit : 5ml × 10vials/box

Dr park's TELO


(Telomere)



Package unit : 3ml X 5vials/box

Ingredients

1. Synthesized Telomere. 50ng/ml
Repeted TTAGGG sequence. 60 base pair
2. PDRN (polydeoxyribonucleotide) 2000ppm
3. Cross linked hyaluronic acid 10mg/ml
4. Glutathione 1%



Composed of repeated TTAGGG sequence

Telomere are the caps at the end of each strand of DNA that protect our chromosomes, like the plastic tips at the end of shoelaces. Without the coating, shoelaces become frayed until they can no longer do their job, just as without telomeres, DNA strands become damaged and our cells can't do their job.

Each time a cell divides, the telomere get shorter. When they get too short, the cell can no longer divide; it becomes inactive or "senescent" or it dies. This shortening process is associated with aging, cancer, and a higher risk of death. So telomere also have been compared with a bomb fuse.

Oligonucleotide synthesizer

Oligonucleotide synthesis is the chemical synthesis of relatively short fragments of nucleic acids with defined chemical structure (sequence).

The technique is extremely useful in current laboratory practice because it provides a rapid and inexpensive access to custom-made oligonucleotides of the desired sequence. Whereas enzymes synthesize DNA and RNA only in a 5' to 3' direction, chemical oligonucleotide synthesis does not have this limitation, although it is, most often, carried out in the opposite, 3' to 5' direction. Currently, the process is implemented as solid-phase synthesis using phosphoramidite method and phosphoramidite building blocks derived from protected 2'-deoxynucleosides (dA, dC, dG, and T), ribonucleosides (A, C, G, and U), or chemically modified nucleosides, e.g. LNA or BNA.

GANA DA

(Lipolysis)

Lipolysis solution for double chin and body

Do not use it for face

Ingredients

Deoxycholic acid 10mg/cc
Q.S-Phosphate buffer solution

Purpose

Emulsifying subcutaneous fat in double chin



Packaging : 5ml × 10 vials

What is deoxycholic acid



Deoxycholic acid has been used since its discovery in various fields of human medicine. In the human body deoxycholic acid is used in the emulsification of fats for absorption in the intestine. It has, in some countries (including Switzerland) been licensed as an emulsifier in food industry, but it is no longer common. Outside the body it is used in experimental basis of cholagogues and is also in use to prevent and dissolve gallstones.

In research deoxycholic acid is used as a mild detergent for the isolation of membrane associated proteins. The critical micelle concentration for deoxycholic acid is approximately 2.4-4 mM.

Sodium deoxycholate, the sodium salt of deoxycholic acid, is often used as a biological detergent to lyse cells and solubilise cellular and membrane components. Sodium deoxycholate mixed with phosphatidylcholine, is used in mesotherapy injections to produce lipolysis, and has been used as an alternative to surgical excision in the treatment of lipomas.

Deoxycholates and bile acid derivatives in general are actively being studied as structures for incorporation in nanotechnology. They also have found application in microlithography as photoresistant components.

Deoxycholic acid



GANA booster-V

(GANA booster in vial)



Biorevitalization

Ingredients

- Cross linked hyaluronic acid 10mg/CC
- Glutathione 1mg/cc

Packaging unit

- 3ml/5ml vials × 2EA/box

GANA HA-V

(GANA HA-B type in vial)



Ingredients

- Cross linked hyaluronic acid
- Concentration : 22mg/cc

Packaging unit

- 3ml/5ml vial × 2EA/box

GANA peptides (APDN)



3ml × 2vials/box

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collage formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA peptides APDN

- Pure peptides solution
- Component : β -Ala-Pro-Dab-NHBn
- Packing unit : 3ml × 2vials/box
- Concentration : 500ppm

What is SYN-AKE

- SYN-AKE is synthesized viper venom tripeptide
- Effect : Anti-wrinkle effect in a month
- Antagonist of acetylcholine receptor
- Relax muscle

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for whole face with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect

GANA peptides (Carnosine)



3ml × 2vials/box

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collage formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA PEPTIDES (L- CARNOSINE)

- Pure peptides solution
- Packing unit : 3ml × 2vials/box
- Concentration : 50ppm

what is carnoisine?

L-Carnosine CAS : 305-84-0

beta-alanyl-L-histidine

Carnosine can increase the Hayflick limit in human fibroblasts, as well as appearing to reduce the telomere shortening rate.

It is also considered as a geroprotector.

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for whole face with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect

GANA peptides (AH8)



3ml × 2vials/box

What is peptide?

The peptides are chain of amino acids
 The each peptides has different amino acids sequence
 Each peptides has different function

For example

- Whitening
- Collagen formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA PEPTIDES (AH8)

- Acetyl hexapeptides 8 (AH8)
- Pure peptides solution
- Packing unit : 3ml × 2vials/box
- Concentration : 20ppm

What is AH8?

Acetyl hexapeptide is made up of peptides, which are chains of amino acids that can affect how cells in the body work. When Acetyl hexapeptide is applied to the skin, it is absorbed into the body and the cream's chemical composition causes the facial muscles to contract with less force. Acetyl hexapeptide prevents the formation of expression wrinkles by somehow inhibiting muscle movement.

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for whole face with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
 So each peptides can be mixed with other peptides
 To make peptides cocktail or other skin booster
 For better effect

GANA peptides (PP3)

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collagen formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA peptides PP3

- Pure peptides solution
- Palmitoyl pentapeptide-3
- Packing unit : 3ml × 2vials/box
- Concentration : 150ppm

Palmitoyl Pentapeptide-3 is a lipophilic modified peptide originally developed for wound healing research and currently being investigated as an anti-aging ingredient. It helps to prevent fine lines induced by repeated facial movements and increased collagen and elastin matrix.

Sequence

- H-Gly-Pro-Arg-Pro-Ala-OH

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for whole face with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect



3ml × 2vials/box

GANA peptides (PT1)



3ml × 2vials/box

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collage formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA peptides PP3

- Palmitoyl Tripeptide-1
- Pure peptides solution
- Packing unit : 3ml × 2vials/box
- Concentration : 100ppm

Palmitoyl Tripeptide-1

palmitoyl tripeptide-1. Also known as pal-GHK and palmitoyl oligopeptide, palmitoyl tripeptide-1 is a synthetic fatty acid-linked peptide that can help repair visible skin damage and strengthen skin's underlying supportive elements.

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for whole face with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect

GANA peptides (MP17)

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collagen formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA peptides MP17

- Myristoyl-Pentapeptide-17
- Pure peptides solution
- Packing unit : 3ml × 2vials/box
- Concentration : 500ppm

mainly based on Myristoyl Pentapeptide-17. It significantly stimulates keratin genes, guides cell growth and growth factor secretion to thicken and enhance the appearance of eyelashes.

Peptide sequence

N2-(1-Oxotetradecyl)-L-lysyl-L-leucyl-L-alanyl-L-lysyl-L-lysineamide

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - Half vial for one session with eye brush be careful not to peptide get in to eyes
3. Interval 2 weeks
4. Total 3-4 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect



3ml × 2vials/box

GANA peptides (AGH1)

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collage formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA peptides AGH1

- Acetyl Glutamyl Heptapeptide-1 (snap-8)
- Pure peptides solution
- Packing unit : 3ml × 2vials/box
- Concentration : 500ppm

peptide solution contains an anti-wrinkle octapeptide that is an elongation of the hexapeptide ARGIRELINE® peptide. Therefore, it reduces expression wrinkles through a Botulinum Toxin-like mechanism of action.

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for whole face with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect



3ml × 2vials/box

GANA peptides (AT5)

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collage formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA peptides AT5

- Acetyl tetrapeptide-5
- Pure peptides solution
- Packing unit : 3ml × 2vials/box
- Concentration : 500ppm

Anti eyebag peptide

Puffy eyebag treatment

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for both eyebag with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect



3ml × 2vials/box

GANA peptides (NP1)



3ml × 2vials/box

What is peptide?

The peptides are chain of amino acids
The each peptides has different amino acids sequence
Each peptides has different function

For example

- Whitening
- Collage formation
- Wrinkle removal
- Muscle relaxation like botulinum toxin

You can choose each peptides products according to your need

Products name : GANA peptides NP1

- Nonapeptide-1
- Pure peptides solution
- Packing unit : 3ml × 2vials/box
- Concentration : 500ppm

Description

Nonapeptide-1 can prevent melanin synthesis and unwanted pigmentation by preventing activation of the tyrosinase, thus allowing for a better control over skin tone.

Functions

Nonapeptide-1 is one skin whitening nonapeptide. It can be applied to various kinds of whitening or spot corrective products.

How to use

1. Clean face with alcohol
2. Apply peptide solution
 - one vial for whole face with 0.2 mm dermaroller or dermastamp to facilitate peptide absorption
3. Interval 2 weeks
4. Total 2-3 session will be needed

TIP

Each peptides has different function
So each peptides can be mixed with other peptides
To make peptides cocktail or other skin booster
For better effect

GANA AM

Main component

Astragalus Membranaceus Root Extract 100ppm



3ml × 2vials/box

What is Astragalus Membranaceus Root Extract used for?

TA-65 in Astragalus Membranaceus Root Extract increase.
Telomerase activity consequently increase length of telomere.

Telomere are the caps at the end of each strand of DNA that protect our chromosomes, like the plastic tips at the end of shoelaces. Without the coating, shoelaces become frayed until they can no longer do their job, just as without telomeres, DNA strands become damaged and our cells can't do their job.

Each time a cell divides, the telomere get shorter. When they get too short, the cell can no longer divide; It becomes inactive or "senescent" or it dies. This shortening process is associated with aging, cancer, and a higher risk of death. So telomere also have been compared with a bomb fuse.

Clinic care

- 84 GANA LIFT (Cone type lifting)
 - 86 GANA PDO (Cog threads)
 - 88 GANA PATCH-KFDA approved (Wound dressing)
-

We also supply the research and development service for customers. We develop and manufacture any products in dermal fillers. Mesotherapy and medical cosmetics field according to customer's need Our MOQ of R&D is reasonable with competitive price

www.ganarnd.co.kr

GANA LIFT

(Cone type lifting)

Currently not in production



What is Dr.S?

Dr.S is the advanced concept of the Lift thread which lifts the wrinkled face from aging and the body which lost its elasticity.

Dr.S improved demerits of existing lift thread and maximized the effect and the maintenance power and the efficiency.

Thread

USP No.1 PP (Polypropylene)

CONE

PLLA (Poly-L-Lactic-Acid) Screw Anchor type 4mm Cone, Biodegradation

What is biodegradable material



The term biodegradable is used to describe materials that decompose through the actions of bacteria, fungi, and other living organisms -simply put it has the ability to break down or decompose back into the natural environment without causing harm. Examples of well known biodegradable materials used in medical industries are PLLA, PLGA, PGA, PDO and PCL.

What is PLLA

Poly(L-lactide) is a biodegradable material with good bio-absorbability and bio-compatibility. After degradation in the body, PLLA products are excreted by metabolism and thus has no harmful side effects to the human body. It is for these reasons that PLLA products are currently widely used in the field of medicine, such as for one-time infusion equipment, medical surgical suture, drug delivery carriers, tissue engineering scaffolds, bone anchors, injectable microcapsules, microspheres, implants, elastomermaterials for organ support, etc.

Better benefits of PLLA bio material


Item	PLLA (Dr.SInnovation)	PDO (Mostin the market)
Tensile strength (MPa)	900	490
Elongation at break (%)	25	35
Modulus of elasticity (GPa)	8.5	2.1
Degradation time (Months)	24	6~7

PAST treatment**SMAS treatment****Comparison**

EZ ; COG




Silhouette




Treatment cost ; EZ < Silhouette
 Priod time ; EZ < Silhouette

Treadtreatment

Silhouette



Dr.S

**2 Types of Dr.S**

Maintenance : 3 year

Efficiency : High tensile lifting effect and reliable traction

Cone : Poly-L-Lactic-acid

Thread USP No.1 : Strong Lift effect

Screw Anchor type cone : 4mm Cone, Biodegradation

Ring : Ultrasonic welding method

The Operation

Dr.S Advantage

Strong Effect face and body : PP thread, Big size Screw Anchor type cone(4mm)



Dr. S Efficiency : High tensile lifting effect and reliable traction.

Biodegradable material safe cones in the human body.

Maintenance : 3 years

Not need to Incisionalface lifting methods

The lifting effect can be seen immediately after the procedure.

**How to procedure for hip****How to procedure for Face**

GANA PDO (Cog threads)

GANA PDO types

Type	Needle length	PDO length	Needle shape	Thread thickness	Needle gauge	Cog type	
GANA PDO 19G-100mm	100 mm	150 mm	W type cannula	1-0	19 gauge	Bidirectional cog	Face lifting
GANA PDO 18G-100mm	100 mm	150 mm	W type cannula	1	18 gauge	Bidirectional cog	Face lifting
GANA PDO 21G-90mm	90 mm	150 mm	L type cannula	2-0	21G	Bidirectional cog	Eyebrow lifting
GANA PDO two needles	100 mm	400 mm		1-0	19G	Bidirectional cog	

We only make 4 type of PDO threads

1. GANA PDO

- 1) 19G-100mm, bidirectional cog
W type
- 2) 18G-100mm, bidirectional cog
W type
- 3) 21G -90mm, bidirectional cog
L type for eyebrow lifting
- 4) Double arms
two needles 400mm
- This product is also for face lifting

GANA PDO

19G-100mm


W type

one pack include 20 threads



GANA PDO types

Type	Needle length	PDO length	Needle shape	Thread thickness	Needle guage	Cog type	
GANA PDO 19G-100mm	100 mm	150 mm	W type cannula	1-0	19 guage	Bidirectional cog	6D
GANA PDO 18G-100mm	100 mm	150 mm	W type cannula	1	18 G	Bidirectional cog	6D
GANA PDO two needles	100 mm	400 mm		1-0	19G	Directional cog	6D



GANA PDO
Double arm
Two needles 400mm
Two threads in one box



How to use double arm

GANA PATCH

KFDA approved
(Wound dressing)



GANA PATCH Package Contents
GANA PATCH sheet
(sheet size 15 × 10 cm)

KFDA and FDA approved product

**GANA PATCH is bio cellulose sheet
containing cross linked hyaluronic acid**

Bacterial cellulose

Main use of GANA PATCH is open wound covering.
GANA PATCH is completely non allergenic and do not cause
any inflammation even in body.

GANA PATCH

Wound covering materials

Development objective

It is impossible to cure patients who has widespread damage skin so far.

What is difference,

- As soon as applied to patient GANA PATCH can keep moisturizing from evaporation
- Prevent infection
- Mechanical-chemical protection
- Absorb exudate
- Reduce pain
- Components in GANA PATCH stimulate the collagen synthesis of soft tissue defect so that there will be no more exudate
- High concentration Cross linked hyaluronic acid gel absorb watery content from wound.
Stuff soft tissue defect immediately and keep moisturizing from evaporation.
- Do not need to change dressing every day.
Change GANA PATCH every week.

In case of major wound, accompanied by soft tissue loss and epithelium damage. GANA PATCH includes active ingredients for epithelial cell and soft tissues regeneration.

Not necessary of skin graft

After an immediate indication, immediate recovery of skin functions from burn injuries or skin loss.

Regeneration soft tissue and epithelium

Bio cellulose itself replaces soft tissue mechanically

Not necessary of wound dressing

Once apply to any wound you just keep watching until you see healed wound

Progress

Bio cellulose itself does not cause any immune side effect and any other response in the body. The epithelial cell migrates to GANA PATCH and then GANA PATCH may be positioned under the epidermis.

WHAT IS THE IDEAL wound dressing?

- Able to resist infection
- Able to prevent water loss
- Able to withstand the shear forces
- Cost effective
- Widely available
- Long shelf life and easy to store
- Lack of antigenicity
- Flexible in thickness
- Durable with long-term wound stability
- Can be conformed to irregular wound surfaces and Easy to be secured and applied

GANA PATCH Package Contents

GANA PATCH sheet
(sheet size 15 × 10 cm)



Bacterial cellulose


Bacterial cellulose is an organic compound with the formula $(C_6H_{10}O_5)_n$ produced by certain types of bacteria. While cellulose is a basic structural material of most plants, it is also produced by bacteria, principally of the genera *Acetobacter*, *Sarcina ventriculi* and *Agrobacterium*. Bacterial, or microbial, cellulose has different properties from plant cellulose and is characterized by high purity, strength, moldability and increased water holding ability. [1] In natural habitats, the majority of bacteria synthesize extracellular polysaccharides, such as cellulose, which form protective envelopes around the cells. While bacterial cellulose is produced in nature, many methods are currently being investigated to enhance cellulose growth from cultures in laboratories as a large-scale process. By controlling synthesis methods, the resulting microbial cellulose can be tailored to have specific desirable properties. For example, attention has been given to the bacteria *Acetobacter xylinum* due to its cellulose's unique mechanical properties and applications to biotechnology, microbiology, and materials science. Historically, bacterial cellulose has been limited to the manufacture of Nata de coco, a South-East Asian food product.[2] With advances in the ability to synthesize and characterize bacterial cellulose, the material is being used for a wide variety of commercial applications including textiles, cosmetics, and food products, as well as medical applications. Many patents have been issued in microbial cellulose applications and several active areas of research are attempting to better characterize microbial cellulose and utilize it in new areas.[1]

SEM findings of Bacterial cellulose

Bacterial cellulose aperture is so small
Any kinds of cells and bacteria can not pass through
only water can pass through
And surface area of bacterial cellulose is so wide
Biocellulose can hold a lot of water inside

In conclusion

Biocellulose can protect any kinds of infection
Absorb exudate from wound and evaporate water outward




- Hyaluronic acid (HA; conjugate base hyaluronate), also called hyaluronan, is an anionic, nonsulfated glycosaminoglycan distributed widely throughout connective, epithelial, and neural tissues. It is unique among glycosaminoglycans in that it is nonsulfated, forms in the plasma membrane instead of the Golgi apparatus, and can be very large, with its molecular weight often reaching the millions.[2] One of the chief components of the extracellular matrix, hyaluronan contributes significantly to cell proliferation and migration, and may also be involved in the progression of some malignant tumors.[3]
- The average 70kg (154lb) person has roughly 15grams of hyaluronan in the body, one-third of which is turned over (degraded and synthesized) every day.[4] Hyaluronic acid is also a component of the group A streptococcal extracellular capsule,[5] and is believed to play a role in virulence.[6][7]

How to use

1. Sterilize wound
2. Take out GANA PATCH sheet out of bottle. Use sterile technique
3. Cover GANA PATCH sheet on the wound
4. Cover GANA PATCH sheet with sterile gauze
5. Change GANA PATCH every week

So you do not have to clean the wound every day
just check wound whether there is infection



ANA R&D

 GANA COSMETIC

 Dr. PARK

ANA R&D MEDICAL DEVICES

We also supply the research and development service for customers. We develop and manufacture any products in dermal fillers. Mesotherapy and medical cosmetics field according to customer's need Our MOQ of R&D is reasonable with competitive price

Address

555, sunil technopia, Dunchon-daero, Jungwon-gu,
Seongnam-si, Gyeonggi-do, South Korea
Office # 1108, Factory 1 #103, Factory 2 #905
Factory 3 #902, Factory 4 # 812, Factory 5 # 909
Factory 6 #904, Factory 7 #1206
Zip Code : 13215

TEL : 82-31-732-0242

FAX : 82-2-6455-7512

E-mail : dydgh416@naver.com

Homepage : www.ganarnd.co.kr