

Representations of Lie Groups

A Mathematical Dissection of Quantum Spin

Part I: Mathematical Foundations

Gordon Chan
Supervisor: Assoc. Prof. François Gay-Balmaz

28 July 2026

Roadmap

- 1 Topological Spaces
- 2 Topological Manifolds
- 3 Smooth Manifolds
- 4 Tangent Spaces
- 5 Algebraic Structures
- 6 Lie Groups
- 7 Matrix Lie Groups
- 8 The Rotation Group $SO(3)$
- 9 The Special Unitary Group $SU(2)$
- 10 The Double Cover $SU(2) \rightarrow SO(3)$
- 11 Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$
- 12 Summary and Next Steps

Definition

A **topological space** is a pair $(X, \mathcal{T})$ where X is a set and $\mathcal{T} \subseteq \mathcal{P}(X)$ satisfies:

- ① (Trivial Bounds) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.
- ② (Arbitrary Unions) $\forall \{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T} : \bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}$.
- ③ (Finite Intersections) $\forall U, V \in \mathcal{T} : U \cap V \in \mathcal{T}$.

Definition

A **topological space** is a pair $(X, \mathcal{T})$ where X is a set and $\mathcal{T} \subseteq \mathcal{P}(X)$ satisfies:

- ① (Trivial Bounds) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.
- ② (Arbitrary Unions) $\forall \{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T} : \bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}$.
- ③ (Finite Intersections) $\forall U, V \in \mathcal{T} : U \cap V \in \mathcal{T}$.

Why this matters: A topology tells us what “closeness” means without needing a distance function. Open sets are the primitive notion of “nearby” — they generalize open intervals in $\mathbb{R}$.

Topological Spaces

Definition

A **topological space** is a pair $(X, \mathcal{T})$ where X is a set and $\mathcal{T} \subseteq \mathcal{P}(X)$ satisfies:

- ① (Trivial Bounds) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.
- ② (Arbitrary Unions) $\forall \{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T} : \bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}$.
- ③ (Finite Intersections) $\forall U, V \in \mathcal{T} : U \cap V \in \mathcal{T}$.

Why this matters: A topology tells us what “closeness” means without needing a distance function. Open sets are the primitive notion of “nearby” — they generalize open intervals in $\mathbb{R}$.

Key constructions: Product topology, subspace topology, metric topologies (e.g. Euclidean topology on $\mathbb{R}^n$).

Definition

A **topological manifold** is a topological space $(M, \mathcal{T})$ satisfying:

- ① (Hausdorff) Points can be separated by disjoint open sets.
- ② (Second-Countable) The topology has a countable basis.
- ③ (Locally Euclidean) Every point has a neighborhood homeomorphic to $\mathbb{R}^n$ for some fixed $n = \dim(M)$.

Definition

A **topological manifold** is a topological space $(M, \mathcal{T})$ satisfying:

- ① (Hausdorff) Points can be separated by disjoint open sets.
- ② (Second-Countable) The topology has a countable basis.
- ③ (Locally Euclidean) Every point has a neighborhood homeomorphic to $\mathbb{R}^n$ for some fixed $n = \dim(M)$.

Intuition: A manifold is a space that *locally looks like* $\mathbb{R}^n$, but may have a nontrivial global shape (e.g. a sphere S^2 or a torus T^2).

Topological Manifolds

Definition

A **topological manifold** is a topological space $(M, \mathcal{T})$ satisfying:

- ① (Hausdorff) Points can be separated by disjoint open sets.
- ② (Second-Countable) The topology has a countable basis.
- ③ (Locally Euclidean) Every point has a neighborhood homeomorphic to $\mathbb{R}^n$ for some fixed $n = \dim(M)$.

Intuition: A manifold is a space that *locally looks like* $\mathbb{R}^n$, but may have a nontrivial global shape (e.g. a sphere S^2 or a torus T^2).

Why Hausdorff + second-countable? These are technical conditions that prevent pathological spaces. They ensure partitions of unity exist.

Definition

A **smooth manifold** is a triple $(M, \mathcal{T}, \mathcal{A})$ where:

- 1 $(M, \mathcal{T})$ is a topological manifold.
- 2 $\mathcal{A} = \{(U_\alpha, \varphi_\alpha)\}_{\alpha \in \mathcal{I}}$ is a **smooth atlas**: each $(U_\alpha, \varphi_\alpha)$ is a chart, the charts cover M , and transition maps $\varphi_\beta \circ \varphi_\alpha^{-1}$ are C^∞ .
- 3 $\mathcal{A}$ is **maximal** w.r.t. smooth compatibility.

Definition

A **smooth manifold** is a triple $(M, \mathcal{T}, \mathcal{A})$ where:

- 1 $(M, \mathcal{T})$ is a topological manifold.
- 2 $\mathcal{A} = \{(U_\alpha, \varphi_\alpha)\}_{\alpha \in \mathcal{I}}$ is a **smooth atlas**: each $(U_\alpha, \varphi_\alpha)$ is a chart, the charts cover M , and transition maps $\varphi_\beta \circ \varphi_\alpha^{-1}$ are C^∞ .
- 3 $\mathcal{A}$ is **maximal** w.r.t. smooth compatibility.

Key idea: Charts are local coordinate systems. Smoothness means that when you switch between coordinate systems, the change-of-coordinates map is infinitely differentiable.

Smooth Maps and Diffeomorphisms

Smooth Map

A map $F : M \rightarrow N$ between smooth manifolds is **smooth** if its coordinate representation $\psi \circ F \circ \varphi^{-1}$ is C^∞ for all compatible charts.

Smooth Maps and Diffeomorphisms

Smooth Map

A map $F : M \rightarrow N$ between smooth manifolds is **smooth** if its coordinate representation $\psi \circ F \circ \varphi^{-1}$ is C^∞ for all compatible charts.

Diffeomorphism

$F : M \rightarrow N$ is a **diffeomorphism** if it is smooth, bijective, and F^{-1} is also smooth. Diffeomorphic manifolds are “the same” from the smooth perspective.

Smooth Maps and Diffeomorphisms

Smooth Map

A map $F : M \rightarrow N$ between smooth manifolds is **smooth** if its coordinate representation $\psi \circ F \circ \varphi^{-1}$ is C^∞ for all compatible charts.

Diffeomorphism

$F : M \rightarrow N$ is a **diffeomorphism** if it is smooth, bijective, and F^{-1} is also smooth. Diffeomorphic manifolds are “the same” from the smooth perspective.

Examples: $\mathbb{R}^n$ is diffeomorphic to the open unit ball $B_1(0)$. S^n is NOT diffeomorphic to $\mathbb{R}^n$ (compact vs. non-compact).

Tangent Spaces via Derivations

Motivation

How do we define tangent vectors to a smooth manifold without embedding it in $\mathbb{R}^N$? Answer: **derivations** — linear maps that satisfy the Leibniz rule.

Tangent Spaces via Derivations

Motivation

How do we define tangent vectors to a smooth manifold without embedding it in $\mathbb{R}^N$? Answer: **derivations** — linear maps that satisfy the Leibniz rule.

Definition

Let M be a smooth manifold, $p \in M$. The **stalk** $C_p^\infty M$ is the set of germs of smooth functions at p . The **tangent space** at p is

$$T_p M = \{ v : C_p^\infty M \rightarrow \mathbb{R} \mid v \text{ is a derivation at } p \}.$$

$T_p M$ is an $\mathbb{R}$ -vector space with $\dim(T_p M) = \dim(M)$.

Tangent Spaces via Derivations

Motivation

How do we define tangent vectors to a smooth manifold without embedding it in $\mathbb{R}^N$? Answer: **derivations** — linear maps that satisfy the Leibniz rule.

Definition

Let M be a smooth manifold, $p \in M$. The **stalk** $C_p^\infty M$ is the set of germs of smooth functions at p . The **tangent space** at p is

$$T_p M = \{ v : C_p^\infty M \rightarrow \mathbb{R} \mid v \text{ is a derivation at } p \}.$$

$T_p M$ is an $\mathbb{R}$ -vector space with $\dim(T_p M) = \dim(M)$.

Why derivations? For $\mathbb{R}^n$, a tangent vector v acts on functions by the directional derivative: $v(f) = \sum_i v^i \frac{\partial f}{\partial x^i} \Big|_p$. This algebraic characterization generalizes to any manifold.

The Differential

Definition

Let $F : M \rightarrow N$ be smooth, $p \in M$. The **differential of F at p** is the linear map $dF_p : T_p M \rightarrow T_{F(p)} N$ defined by

$$\forall v \in T_p M, \forall [f]_{F(p)} \in C_{F(p)}^\infty N : \quad (dF_p(v))([f]_{F(p)}) = v([f \circ F]_p).$$

The Differential

Definition

Let $F : M \rightarrow N$ be smooth, $p \in M$. The **differential of F at p** is the linear map $dF_p : T_p M \rightarrow T_{F(p)} N$ defined by

$$\forall v \in T_p M, \forall [f]_{F(p)} \in C_{F(p)}^\infty N : \quad (dF_p(v))([f]_{F(p)}) = v([f \circ F]_p).$$

Chain Rule

For smooth maps $F : M \rightarrow N$ and $G : N \rightarrow P$,

$$d(G \circ F)_p = dG_{F(p)} \circ dF_p.$$

The Differential

Definition

Let $F : M \rightarrow N$ be smooth, $p \in M$. The **differential of F at p** is the linear map $dF_p : T_p M \rightarrow T_{F(p)} N$ defined by

$$\forall v \in T_p M, \forall [f]_{F(p)} \in C_{F(p)}^\infty N : \quad (dF_p(v))([f]_{F(p)}) = v([f \circ F]_p).$$

Chain Rule

For smooth maps $F : M \rightarrow N$ and $G : N \rightarrow P$,

$$d(G \circ F)_p = dG_{F(p)} \circ dF_p.$$

Interpretation: The differential is the “best linear approximation” to a smooth map, generalizing the Jacobian matrix from multivariable calculus.

Groups, Fields, and Vector Spaces

Group

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Groups, Fields, and Vector Spaces

Group

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Field

A set $\mathbb{F}$ (e.g. $\mathbb{R}, \mathbb{C}$) with addition, multiplication, and all the usual arithmetic axioms — a domain where linear algebra works.

Groups, Fields, and Vector Spaces

Group

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Field

A set $\mathbb{F}$ (e.g. $\mathbb{R}, \mathbb{C}$) with addition, multiplication, and all the usual arithmetic axioms — a domain where linear algebra works.

Vector Space

An abelian group $(V, +)$ with scalar multiplication $\mathbb{F} \times V \rightarrow V$ satisfying distributivity and compatibility axioms.

Groups, Fields, and Vector Spaces

Group

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Field

A set $\mathbb{F}$ (e.g. $\mathbb{R}, \mathbb{C}$) with addition, multiplication, and all the usual arithmetic axioms — a domain where linear algebra works.

Vector Space

An abelian group $(V, +)$ with scalar multiplication $\mathbb{F} \times V \rightarrow V$ satisfying distributivity and compatibility axioms.

Why these matter: Lie groups combine the smooth structure of manifolds with the algebraic structure of groups. Lie algebras are vector spaces with an extra bracket operation.

$\mathbb{F}$ -Algebra

A vector space $\mathcal{A}$ over $\mathbb{F}$ together with a bilinear multiplication $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$.
(Examples: $\mathbb{R}^{n,n}$ with matrix multiplication, $C^\infty(M)$ with pointwise multiplication.)

Algebras and Lie Algebras

$\mathbb{F}$ -Algebra

A vector space $\mathcal{A}$ over $\mathbb{F}$ together with a bilinear multiplication $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$. (Examples: $\mathbb{R}^{n,n}$ with matrix multiplication, $C^\infty(M)$ with pointwise multiplication.)

Lie Algebra

An $\mathbb{F}$ -vector space $\mathfrak{g}$ with a bilinear bracket $[\bullet, \bullet] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying:

- 1 (Skew-symmetry) $[Y, X] = -[X, Y]$
- 2 (Jacobi identity) $[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$

Algebras and Lie Algebras

$\mathbb{F}$ -Algebra

A vector space $\mathcal{A}$ over $\mathbb{F}$ together with a bilinear multiplication $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$. (Examples: $\mathbb{R}^{n,n}$ with matrix multiplication, $C^\infty(M)$ with pointwise multiplication.)

Lie Algebra

An $\mathbb{F}$ -vector space $\mathfrak{g}$ with a bilinear bracket $[\bullet, \bullet] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying:

- ① (Skew-symmetry) $[Y, X] = -[X, Y]$
- ② (Jacobi identity) $[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$

Key example: $(\mathbb{R}^3, \times)$ — the cross product makes $\mathbb{R}^3$ a Lie algebra!
 $[v, w] = v \times w$ satisfies skew-symmetry and Jacobi.

Lie Groups

Definition

A **Lie group** is a smooth manifold $(G, \mathcal{T}, \mathcal{A})$ that is also a group $(G, \{\mu, \iota, e\})$ such that:

- 1 $\mu : G \times G \rightarrow G$ (multiplication) is smooth.
- 2 $\iota : G \rightarrow G$ (inversion) is smooth.

Lie Groups

Definition

A **Lie group** is a smooth manifold $(G, \mathcal{T}, \mathcal{A})$ that is also a group $(G, \{\mu, \iota, e\})$ such that:

- 1 $\mu : G \times G \rightarrow G$ (multiplication) is smooth.
- 2 $\iota : G \rightarrow G$ (inversion) is smooth.

The synthesis: A Lie group unifies geometry (smooth manifold) with algebra (group). This allows us to study continuous symmetries using both differential calculus and group theory.

Lie Groups

Definition

A **Lie group** is a smooth manifold $(G, \mathcal{T}, \mathcal{A})$ that is also a group $(G, \{\mu, \iota, e\})$ such that:

- 1 $\mu : G \times G \rightarrow G$ (multiplication) is smooth.
- 2 $\iota : G \rightarrow G$ (inversion) is smooth.

The synthesis: A Lie group unifies geometry (smooth manifold) with algebra (group). This allows us to study continuous symmetries using both differential calculus and group theory.

Examples:

- $(\mathbb{R}^n, +)$ — the translation group.
- $GL(n, \mathbb{R})$ — invertible $n \times n$ real matrices.
- $SO(n)$ — rotations in $\mathbb{R}^n$.
- $SU(n)$ — unitary transformations with determinant 1.

Left Translation and Left-Invariant Vector Fields

Left Translation

For $g \in G$, $L_g : G \rightarrow G$ is defined by $L_g(h) = gh$. L_g is a diffeomorphism.

Left Translation and Left-Invariant Vector Fields

Left Translation

For $g \in G$, $L_g : G \rightarrow G$ is defined by $L_g(h) = gh$. L_g is a diffeomorphism.

Left-Invariant Vector Field

A smooth vector field X on G is **left-invariant** if

$$\forall g \in G, \forall f \in C^\infty(G) : \quad X(f \circ L_g) = X(f) \circ L_g.$$

Left Translation and Left-Invariant Vector Fields

Left Translation

For $g \in G$, $L_g : G \rightarrow G$ is defined by $L_g(h) = gh$. L_g is a diffeomorphism.

Left-Invariant Vector Field

A smooth vector field X on G is **left-invariant** if

$$\forall g \in G, \forall f \in C^\infty(G) : \quad X(f \circ L_g) = X(f) \circ L_g.$$

Geometric meaning: A left-invariant vector field is completely determined by its value at the identity e . Translating the vector at e by L_g gives the vector at any point g .

The Lie Algebra of a Lie Group

Theorem

Let G be a Lie group. Define $\mathfrak{g} = T_e G$ (tangent space at identity). For each $v \in \mathfrak{g}$, there exists a **unique** left-invariant vector field X^v with $X_e^v = v$. Define the bracket on $\mathfrak{g}$ by

$$[v, w]_{\mathfrak{g}} = [X^v, X^w]_{\circ}|_e.$$

Then $(\mathfrak{g}, [\bullet, \bullet]_{\mathfrak{g}})$ is a Lie $\mathbb{R}$ -algebra with $\dim(\mathfrak{g}) = \dim(G)$.

The Lie Algebra of a Lie Group

Theorem

Let G be a Lie group. Define $\mathfrak{g} = T_e G$ (tangent space at identity). For each $v \in \mathfrak{g}$, there exists a **unique** left-invariant vector field X^v with $X_e^v = v$. Define the bracket on $\mathfrak{g}$ by

$$[v, w]_{\mathfrak{g}} = [X^v, X^w]_{\circ}|_e.$$

Then $(\mathfrak{g}, [\bullet, \bullet]_{\mathfrak{g}})$ is a Lie $\mathbb{R}$ -algebra with $\dim(\mathfrak{g}) = \dim(G)$.

The bridge: Every Lie group gives rise to a Lie algebra (its tangent space at the identity). The Lie algebra captures the **infinitesimal** structure of the group.

The Closed Subgroup Theorem (É. Cartan)

Theorem

Let G be a Lie group and $H \subseteq G$ a closed subgroup. Then H admits a unique smooth structure making it a Lie group such that the inclusion $H \hookrightarrow G$ is a smooth embedding.

The Closed Subgroup Theorem (É. Cartan)

Theorem

Let G be a Lie group and $H \subseteq G$ a closed subgroup. Then H admits a unique smooth structure making it a Lie group such that the inclusion $H \hookrightarrow G$ is a smooth embedding.

Powerful consequence: To show a subset of matrices is a Lie group, you only need to check it's a closed subgroup of some $GL(n, \mathbb{F})$. No need to construct charts explicitly!

The Closed Subgroup Theorem (É. Cartan)

Theorem

Let G be a Lie group and $H \subseteq G$ a closed subgroup. Then H admits a unique smooth structure making it a Lie group such that the inclusion $H \hookrightarrow G$ is a smooth embedding.

Powerful consequence: To show a subset of matrices is a Lie group, you only need to check it's a closed subgroup of some $GL(n, \mathbb{F})$. No need to construct charts explicitly!

Examples: $SO(n)$, $SU(n)$, $SL(n, \mathbb{R})$, $Sp(2n)$ are all closed subgroups of $GL(N, \mathbb{F})$ and therefore Lie groups.

The General Linear Group $GL(n, \mathbb{F})$

Definition

$$GL(n, \mathbb{F}) = \{ \mathbf{A} \in \mathbb{F}^{n,n} \mid \det(\mathbf{A}) \neq 0 \}$$

The General Linear Group $GL(n, \mathbb{F})$

Definition

$$GL(n, \mathbb{F}) = \{ \mathbf{A} \in \mathbb{F}^{n,n} \mid \det(\mathbf{A}) \neq 0 \}$$

- $GL(n, \mathbb{F})$ is an open subset of $\mathbb{F}^{n,n} \cong \mathbb{F}^{n^2}$, so it is a smooth manifold of dimension n^2 .

The General Linear Group $GL(n, \mathbb{F})$

Definition

$$GL(n, \mathbb{F}) = \{ \mathbf{A} \in \mathbb{F}^{n,n} \mid \det(\mathbf{A}) \neq 0 \}$$

- $GL(n, \mathbb{F})$ is an open subset of $\mathbb{F}^{n,n} \cong \mathbb{F}^{n^2}$, so it is a smooth manifold of dimension n^2 .
- Its Lie algebra is $\mathfrak{gl}(n, \mathbb{F}) = \mathbb{F}^{n,n}$ (all $n \times n$ matrices) with the commutator bracket:

$$[\mathbf{A}, \mathbf{B}] = \mathbf{AB} - \mathbf{BA}.$$

The General Linear Group $GL(n, \mathbb{F})$

Definition

$$GL(n, \mathbb{F}) = \{ \mathbf{A} \in \mathbb{F}^{n,n} \mid \det(\mathbf{A}) \neq 0 \}$$

- $GL(n, \mathbb{F})$ is an open subset of $\mathbb{F}^{n,n} \cong \mathbb{F}^{n^2}$, so it is a smooth manifold of dimension n^2 .
- Its Lie algebra is $\mathfrak{gl}(n, \mathbb{F}) = \mathbb{F}^{n,n}$ (all $n \times n$ matrices) with the commutator bracket:

$$[\mathbf{A}, \mathbf{B}] = \mathbf{AB} - \mathbf{BA}.$$

- Most matrix Lie groups arise as closed subgroups of $GL(n, \mathbb{F})$. Their Lie algebras are subspaces of $\mathfrak{gl}(n, \mathbb{F})$ closed under the commutator.

The Special Orthogonal Group $SO(3)$

Definition

$$SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} \mid \mathbf{R}^T \mathbf{R} = \mathbf{I}_3, \det(\mathbf{R}) = 1 \}.$$

The Special Orthogonal Group $SO(3)$

Definition

$$SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} \mid \mathbf{R}^T \mathbf{R} = \mathbf{I}_3, \det(\mathbf{R}) = 1 \}.$$

Lie Algebra $\mathfrak{so}(3)$

$$\mathfrak{so}(3) = T_{\mathbf{I}_3} SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} \mid \mathbf{R}^T = -\mathbf{R} \}.$$

$\dim(\mathfrak{so}(3)) = 3$. A basis $(\mathbf{L}_1, \mathbf{L}_2, \mathbf{L}_3)$:

$$\mathbf{L}_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \mathbf{L}_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \quad \mathbf{L}_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The Special Orthogonal Group $SO(3)$

Definition

$$SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} \mid \mathbf{R}^T \mathbf{R} = \mathbf{I}_3, \det(\mathbf{R}) = 1 \}.$$

Lie Algebra $\mathfrak{so}(3)$

$$\mathfrak{so}(3) = T_{\mathbf{I}_3} SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} \mid \mathbf{R}^T = -\mathbf{R} \}.$$

$\dim(\mathfrak{so}(3)) = 3$. A basis $(\mathbf{L}_1, \mathbf{L}_2, \mathbf{L}_3)$:

$$\mathbf{L}_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \mathbf{L}_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \quad \mathbf{L}_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Commutation Relations

$[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ — the same structure as the cross product on $\mathbb{R}^3$!

The Special Unitary Group $SU(2)$

Definition

$$SU(2) = \{ \mathbf{U} \in \mathbb{C}^{2,2} \mid \mathbf{U}^\dagger \mathbf{U} = \mathbf{I}_2, \det(\mathbf{U}) = 1 \}.$$

The Special Unitary Group $SU(2)$

Definition

$$SU(2) = \{ \mathbf{U} \in \mathbb{C}^{2,2} \mid \mathbf{U}^\dagger \mathbf{U} = \mathbf{I}_2, \det(\mathbf{U}) = 1 \}.$$

Lie Algebra $\mathfrak{su}(2)$

$$\mathfrak{su}(2) = T_{\mathbf{I}_2} SU(2) = \{ \mathbf{U} \in \mathbb{C}^{2,2} \mid \mathbf{U}^\dagger = -\mathbf{U}, \operatorname{tr}(\mathbf{U}) = 0 \}.$$

$\dim(\mathfrak{su}(2)) = 3$. A basis $(\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$:

$$\mathbf{T}_1 = \frac{1}{2} \begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix}, \quad \mathbf{T}_2 = \frac{1}{2} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \mathbf{T}_3 = \frac{1}{2} \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}.$$

The Special Unitary Group $SU(2)$

Definition

$$SU(2) = \{ \mathbf{U} \in \mathbb{C}^{2,2} \mid \mathbf{U}^\dagger \mathbf{U} = \mathbf{I}_2, \det(\mathbf{U}) = 1 \}.$$

Lie Algebra $\mathfrak{su}(2)$

$$\mathfrak{su}(2) = T_{\mathbf{I}_2} SU(2) = \{ \mathbf{U} \in \mathbb{C}^{2,2} \mid \mathbf{U}^\dagger = -\mathbf{U}, \operatorname{tr}(\mathbf{U}) = 0 \}.$$

$\dim(\mathfrak{su}(2)) = 3$. A basis $(\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$:

$$\mathbf{T}_1 = \frac{1}{2} \begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix}, \quad \mathbf{T}_2 = \frac{1}{2} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \mathbf{T}_3 = \frac{1}{2} \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}.$$

Commutation Relations

$[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$ — **identical** to $\mathfrak{so}(3)$! This is the first hint of the isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$.

The Double Cover $\Phi : \mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$

Construction

For $\mathbf{U} \in \mathrm{SU}(2)$, define $\Phi(\mathbf{U}) : \mathfrak{su}(2) \rightarrow \mathfrak{su}(2)$ by

$$\Phi(\mathbf{U})(\mathbf{X}) = \mathbf{U}\mathbf{X}\mathbf{U}^\dagger.$$

Using the basis $\beta = (\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$, identify $\mathfrak{su}(2) \cong \mathbb{R}^3$. Then the matrix $[\Phi(\mathbf{U})]_\beta^\beta$ lies in $\mathrm{SO}(3)$.

The Double Cover $\Phi : \text{SU}(2) \rightarrow \text{SO}(3)$

Construction

For $\mathbf{U} \in \text{SU}(2)$, define $\Phi(\mathbf{U}) : \mathfrak{su}(2) \rightarrow \mathfrak{su}(2)$ by

$$\Phi(\mathbf{U})(\mathbf{X}) = \mathbf{U}\mathbf{X}\mathbf{U}^\dagger.$$

Using the basis $\beta = (\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$, identify $\mathfrak{su}(2) \cong \mathbb{R}^3$. Then the matrix $[\Phi(\mathbf{U})]_\beta^\beta$ lies in $\text{SO}(3)$.

Theorem


$\Phi : \text{SU}(2) \rightarrow \text{SO}(3)$ is:

- 1 A smooth group homomorphism.
- 2 Surjective.
- 3 Has kernel $\ker(\Phi) = \{\pm \mathbf{I}_2\}$ (two elements).

Thus $\text{SU}(2)$ is a **double cover** of $\text{SO}(3)$.


The Double Cover: Geometric Picture

$$\mathrm{SU}(2) \cong S^3$$




$$\xrightarrow{\quad \Phi \quad}$$

$$\mathrm{SO}(3) \cong \mathbb{R}P^3$$



The Double Cover: Geometric Picture



Why two-to-one? Both $\mathbf{U}$ and $-\mathbf{U}$ give the same rotation. A rotation by angle θ about axis $\hat{\mathbf{n}}$ lifts to $\pm \cos(\theta/2)\mathbf{I}_2 \mp i \sin(\theta/2)(\hat{\mathbf{n}} \cdot \boldsymbol{\sigma})$. Increasing θ by 2π flips the sign!

Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$

Theorem

Define $\varphi : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2)$ by $\varphi(\mathbf{L}_i) = \mathbf{T}_i$ (and extend $\mathbb{R}$ -linearly). Then φ is a Lie algebra isomorphism.

Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$

Theorem

Define $\varphi : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2)$ by $\varphi(\mathbf{L}_i) = \mathbf{T}_i$ (and extend $\mathbb{R}$ -linearly). Then φ is a Lie algebra isomorphism.

Proof (essence)

- φ maps a basis to a basis $\implies$ linear isomorphism.
- Both satisfy $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ and $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$.
- Therefore $\varphi([\mathbf{L}_i, \mathbf{L}_j]) = \varphi(\varepsilon_{ijk} \mathbf{L}_k) = \varepsilon_{ijk} \mathbf{T}_k = [\mathbf{T}_i, \mathbf{T}_j] = [\varphi(\mathbf{L}_i), \varphi(\mathbf{L}_j)]$.

Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$

Theorem

Define $\varphi : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2)$ by $\varphi(\mathbf{L}_i) = \mathbf{T}_i$ (and extend $\mathbb{R}$ -linearly). Then φ is a Lie algebra isomorphism.

Proof (essence)

- φ maps a basis to a basis $\implies$ linear isomorphism.
- Both satisfy $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ and $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$.
- Therefore $\varphi([\mathbf{L}_i, \mathbf{L}_j]) = \varphi(\varepsilon_{ijk} \mathbf{L}_k) = \varepsilon_{ijk} \mathbf{T}_k = [\mathbf{T}_i, \mathbf{T}_j] = [\varphi(\mathbf{L}_i), \varphi(\mathbf{L}_j)]$.

Significance: Although $SU(2)$ and $SO(3)$ are different as groups (double cover vs. quotient), their Lie algebras are **identical**. Infinitesimal rotations are the same in both pictures.

What We've Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds**
— the geometric foundation.

What We've Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds**
— the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.

What We've Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- ③ **Lie groups** = smooth manifolds + groups — where geometry meets algebra.

What We've Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- ③ **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- ④ Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).

What We've Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- ③ **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- ④ Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).
- ⑤ The **Closed Subgroup Theorem** lets us recognize matrix Lie groups easily.

What We've Built So Far

- 1 **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- 2 **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- 3 **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- 4 Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).
- 5 The **Closed Subgroup Theorem** lets us recognize matrix Lie groups easily.
- 6 $SO(3)$ (rotations) and $SU(2)$ (unitary 2×2 matrices) are the central examples.

What We've Built So Far

- 1 **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- 2 **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- 3 **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- 4 Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).
- 5 The **Closed Subgroup Theorem** lets us recognize matrix Lie groups easily.
- 6 $SO(3)$ (rotations) and $SU(2)$ (unitary 2×2 matrices) are the central examples.
- 7 $SU(2)$ is a **double cover** of $SO(3)$. Their Lie algebras are **isomorphic**: $\mathfrak{so}(3) \cong \mathfrak{su}(2)$.

What Comes Next (Part II)

Representation Theory (next meeting)

- Representations of Lie algebras and Lie groups.
- Complexification $\mathfrak{su}(2)_{\mathbb{C}} \cong \mathfrak{sl}(2, \mathbb{C})$ and the Cartan–Weyl basis (H, E, F) .
- Classification of irreducible representations of $\mathfrak{su}(2)$: for each $j \in \{0, \frac{1}{2}, 1, \frac{3}{2}, \dots\}$, a unique irrep of dimension $2j + 1$.
- Integration: every $\mathfrak{su}(2)$ -representation exponentiates to an $SU(2)$ -representation.
- Projective representations and the descent to $SO(3)$: integer $j \rightarrow$ ordinary representations, half-integer $j \rightarrow$ spinor (projective) representations.
- Physical spin operators and the 4π periodicity of spinors.

- Hall, B. C. (2015). *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction* (2nd ed.). Springer.
- Lee, J. M. (2012). *Introduction to Smooth Manifolds* (2nd ed.). Springer.
- Steinacker, H. (2024). *Lie Groups and Lie Algebras for Physicists*. Lecture notes, University of Vienna.
- Full paper with proofs: github.com/ganddoffcha/quantum-spin

Thank you!

Questions?