

REPRESENTATIONS OF LIE GROUPS

A MATHEMATICAL DISSECTION OF QUANTUM SPIN

GORDON CHAN

ABSTRACT. Introductory quantum mechanics courses often present the concept of spin heuristically. In this project, we investigate the mathematical foundations of quantum mechanical spin. In particular, we explore the geometric and algebraic origins of spin, in the language of Lie groups, Lie algebras, and their unitary and irreducible representations. A central focus is the topological and algebraic relationship between the rotation group $\mathrm{SO}(3)$ and its universal double cover, the special unitary group $\mathrm{SU}(2)$, which is diffeomorphic to $\mathbb{S}^3$. By classifying the irreducible representations of $\mathrm{SU}(2)$, we present the formalism of how this abstract algebraic classification naturally gives rise to the physical phenomenon of integer and half-integer spin. Ultimately, we hope to bridge topology and abstract algebra with the physical ideas in quantum physics.

1. TOPOLOGICAL SPACES

Definition 1.1. A topological space is a pair $(X, \mathcal{T})$, where

- X is a set, and
- $\mathcal{T} \subseteq \mathcal{P}(X)$ is a subcollection of subsets of X ,

that satisfy the following three properties:

- (1) (Trivial Bounds) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.
- (2) (Arbitrary Unions) $\forall \{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T} : \bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}$.
- (3) (Finite Intersections) $\forall U, V \in \mathcal{T} : U \cap V \in \mathcal{T}$.

Definition 1.2. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces. The product topology on $X \times Y$, denoted by “ $\mathcal{T}_X \otimes \mathcal{T}_Y$ ”, is defined as:

$$\mathcal{T}_X \otimes \mathcal{T}_Y \stackrel{\text{df}}{=} \{W \subseteq X \times Y : \forall (x, y) \in W, \exists U \in \mathcal{T}_X, \exists V \in \mathcal{T}_Y : (x, y) \in U \times V \subseteq W\}.$$

Proposition 1.3 (Product Space is a Topological Space). *Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces. Then the pair $(X \times Y, \mathcal{T}_X \otimes \mathcal{T}_Y)$ is a topological space.*

Proof. $X \times Y$ is a set. By definition, $\mathcal{T}_X \otimes \mathcal{T}_Y \subseteq \mathcal{P}(X \times Y)$.

- (1) (Trivial Bounds) Vacuously, $\emptyset \in \mathcal{T}_X \otimes \mathcal{T}_Y$. For any $(x, y) \in X \times Y$, we have $X \in \mathcal{T}_X$, $Y \in \mathcal{T}_Y$, and $(x, y) \in X \times Y \subseteq X \times Y$, so $X \times Y \in \mathcal{T}_X \otimes \mathcal{T}_Y$.
- (2) (Arbitrary Unions) Let $\{W_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T}_X \otimes \mathcal{T}_Y$. Let $(x, y) \in \bigcup_{\alpha \in \mathcal{I}} W_\alpha$. We can find a particular $\alpha^* \in \mathcal{I}$ such that $(x, y) \in W_{\alpha^*} \in \mathcal{T}_X \otimes \mathcal{T}_Y$. By definition, we can find $U \in \mathcal{T}_X$ and $V \in \mathcal{T}_Y$ such that $(x, y) \in U \times V \subseteq W_{\alpha^*} \subseteq \bigcup_{\alpha \in \mathcal{I}} W_\alpha$. Thus, $\bigcup_{\alpha \in \mathcal{I}} W_\alpha \in \mathcal{T}_X \otimes \mathcal{T}_Y$.
- (3) (Finite Intersections) Let $W_1, W_2 \in \mathcal{T}_X \otimes \mathcal{T}_Y$. Let $(x, y) \in W_1 \cap W_2$. By definition, we can find $U_1 \times V_1, U_2 \times V_2 \in \mathcal{T}_X \otimes \mathcal{T}_Y$ such that $(x, y) \in U_1 \times V_1 \subseteq W_1$ and $(x, y) \in U_2 \times V_2 \subseteq W_2$. So,

$(x, y) \in (U_1 \times V_1) \cap (U_2 \times V_2) \subseteq W_1 \cap W_2$. Since $(U_1 \times V_1) \cap (U_2 \times V_2) = \underbrace{(U_1 \cap U_2)}_{\in \mathcal{T}_X} \times \underbrace{(V_1 \cap V_2)}_{\in \mathcal{T}_Y}$, we have $W_1 \cap W_2 \in \mathcal{T}_X \otimes \mathcal{T}_Y$, as desired. $\square$

Definition 1.4. Let $(X, \mathcal{T})$ be a topological space. Let $A \subseteq X$. The subspace topology on A , denoted by “ $\mathcal{T}|_A$ ”, is defined as:

$$\mathcal{T}|_A \stackrel{\text{df}}{=} \{A \cap U : U \in \mathcal{T}\}.$$

Proposition 1.5 (Subspace Topology is a Topology). *Let $(X, \mathcal{T})$ be a topological space. Let $A \subseteq X$. Then the pair $(A, \mathcal{T}|_A)$ is a topological space.*

Proof. A is a set. For any $A \cap U \in \mathcal{T}|_A$, $A \cap U \subseteq A$, so $A \cap U \in \mathcal{P}(A)$, yielding $\mathcal{T}|_A \subseteq \mathcal{P}(A)$.

(1) (Trivial Bounds) Since $\emptyset \in \mathcal{T}$, $\emptyset = A \cap \emptyset \in \mathcal{T}|_A$. Since $X \in \mathcal{T}$, $A = A \cap X \in \mathcal{T}|_A$.

(2) (Arbitrary Unions) Let $\{A \cap U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T}|_A$. We have $\bigcup_{\alpha \in \mathcal{I}} (A \cap U_\alpha) = A \cap \underbrace{\bigcup_{\alpha \in \mathcal{I}} U_\alpha}_{\in \mathcal{T}} \in \mathcal{T}|_A$.

(3) (Finite Intersections) Let $A \cap U, A \cap V \in \mathcal{T}|_A$. We have $(A \cap U) \cap (A \cap V) = A \cap \underbrace{(U \cap V)}_{\in \mathcal{T}} \in \mathcal{T}|_A$. $\square$

Proposition 1.6 (Iterated Subspace Topology). *Let $(X, \mathcal{T})$ be a topological space. Let $A, B \subseteq X$. Then*

$$(\mathcal{T}|_A)|_B = \mathcal{T}|_{A \cap B} = (\mathcal{T}|_B)|_A.$$

Proof.

$$\begin{aligned} (\mathcal{T}|_A)|_B &= \{B \cap U : U \in \mathcal{T}|_A\} \\ &= \{B \cap (A \cap U) : U \in \mathcal{T}\} \\ &= \{(A \cap B) \cap U : U \in \mathcal{T}\} = \mathcal{T}|_{A \cap B} \\ &= \{A \cap (B \cap U) : U \in \mathcal{T}\} \\ &= \{A \cap U : U \in \mathcal{T}|_B\} = (\mathcal{T}|_B)|_A. \end{aligned}$$

$\square$

Theorem 1.7 (Commutivity of Product and Subspace). *Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces. Let $U \in \mathcal{T}_X$ and $V \in \mathcal{T}_Y$. Then $\mathcal{T}_X|_U \otimes \mathcal{T}_Y|_V = (\mathcal{T}_X \otimes \mathcal{T}_Y)|_{U \times V}$.*

Proof. ($\subseteq$) Let $W \in \mathcal{T}_X|_U \otimes \mathcal{T}_Y|_V$. Let $(x, y) \in W$. By the definition of product topology, we can find $A \cap U \in \mathcal{T}_X|_U$ and $B \cap V \in \mathcal{T}_Y|_V$ with $A \in \mathcal{T}_X$ and $B \in \mathcal{T}_Y$, such that $(x, y) \in (A \cap U) \times (B \cap V) \subseteq W$. Now notice that $(A \cap U) \times (B \cap V) = (A \times B) \cap (U \times V)$. Since $A \times B \in \mathcal{T}_X \otimes \mathcal{T}_Y$, we have $(A \times B) \cap (U \times V) \in (\mathcal{T}_X \otimes \mathcal{T}_Y)|_{U \times V}$. So, for any $(x, y) \in W$, we can find a $\mathcal{O}_{(x, y)} \in (\mathcal{T}_X \otimes \mathcal{T}_Y)|_{U \times V}$, such that $(x, y) \in \mathcal{O}_{(x, y)} \subseteq W$. Notice the following trick:

$$W = \bigcup_{(x, y) \in W} \mathcal{O}_{(x, y)}.$$

Together with the Arbitrary Union property, we conclude $W \in (\mathcal{T}_X \otimes \mathcal{T}_Y)|_{U \times V}$.

($\supseteq$) Let $W \cap (U \times V) \in (\mathcal{T}_X \otimes \mathcal{T}_Y)|_{U \times V}$ with $W \in \mathcal{T}_X \otimes \mathcal{T}_Y$. Let $(x, y) \in W \cap (U \times V)$. This means $(x, y) \in W$ with $x \in U$ and $y \in V$. Since $W \in \mathcal{T}_X \otimes \mathcal{T}_Y$, we can find $A \in \mathcal{T}_X$ and $B \in \mathcal{T}_Y$ such that

$(x, y) \in A \times B \subseteq W$. Notice that $(x, y) \in \underbrace{(A \cap U)}_{\in \mathcal{T}_X|_U} \times \underbrace{(B \cap V)}_{\in \mathcal{T}_Y|_V} = (A \times B) \cap (U \times V) \subseteq W \cap (U \times V)$, which is precisely the definition for $W \cap (U \times V) \in \mathcal{T}_X|_U \otimes \mathcal{T}_Y|_V$. $\square$

Definition 1.8. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces. A map $f : X \rightarrow Y$ is said to be continuous w.r.t. $\mathcal{T}_X$ and $\mathcal{T}_Y$ iff:

$$\forall V \in \mathcal{T}_Y : f^{-1}(V) \in \mathcal{T}_X.$$

Definition 1.9. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces. A map $f : X \rightarrow Y$ is said to be a homeomorphism between $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ iff it satisfies the following three properties:

- (1) (Invertibility) $f : X \rightarrow Y$ is a bijection.
- (2) (Continuity) $f : X \rightarrow Y$ is continuous w.r.t. $\mathcal{T}_X$ and $\mathcal{T}_Y$.
- (3) (Continuity of the Inverse) $f^{-1} : Y \rightarrow X$ is continuous w.r.t. $\mathcal{T}_Y$ and $\mathcal{T}_X$.

Definition 1.10. A topological space $(X, \mathcal{T}_X)$ is said to be homeomorphic to another topological space $(Y, \mathcal{T}_Y)$, denoted by “ $(X, \mathcal{T}_X) \cong (Y, \mathcal{T}_Y)$ ”, iff there exists a homeomorphism between $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$.

Theorem 1.11 (Restriction of a homeomorphism to a subspace). *Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces. Let $(X, \mathcal{T}_X) \cong (Y, \mathcal{T}_Y)$. Let $A \subseteq X$. Let $f : X \rightarrow Y$ be a homeomorphism between $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$. Then $(A, \mathcal{T}_X|_A) \cong (f(A), \mathcal{T}_Y|_{f(A)})$.*

Proof. Consider the restriction $f|_A : A \rightarrow f(A)$. We show that it is a homeomorphism between $(A, \mathcal{T}_X|_A)$ and $(f(A), \mathcal{T}_Y|_{f(A)})$.

- (1) (Invertibility) The restriction $f^{-1}|_{f(A)} : f(A) \rightarrow A$ is the inverse.
- (2) (Continuity of $f|_A$) Let $V \cap f(A) \in \mathcal{T}|_{f(A)}$ with $V \in \mathcal{T}_Y$. We have the following:

$$\begin{aligned} \left[(f|_A)^{-1} \right] (V \cap f(A)) &= \{x \in A : f|_A(x) \in V \cap f(A)\} \\ &= \{x \in A : f|_A(x) \in V \text{ and } f|_A(x) \in f(A)\} \\ &= \{x \in A : f|_A(x) \in V\} \\ &= A \cap \underbrace{f^{-1}(V)}_{\in \mathcal{T}_X} \in \mathcal{T}_X|_A. \end{aligned}$$

- (3) (Continuity of $f^{-1}|_{f(A)}$) The same argument as (2), with f replaced by f^{-1} .

$\square$

Theorem 1.12 (Invariance of the Product under Homeomorphism). *Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$, $(Z, \mathcal{T}_Z)$ and $(W, \mathcal{T}_W)$ be topological spaces. Let $(X, \mathcal{T}_X) \cong (Y, \mathcal{T}_Y)$ and $(Z, \mathcal{T}_Z) \cong (W, \mathcal{T}_W)$. Then $(X \times Z, \mathcal{T}_X \otimes \mathcal{T}_Z) \cong (Y \times W, \mathcal{T}_Y \otimes \mathcal{T}_W)$.*

Proof. Let $f : X \rightarrow Y$ be a homeomorphism between $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$. Let $g : Z \rightarrow W$ be a homeomorphism between $(Z, \mathcal{T}_Z)$ and $(W, \mathcal{T}_W)$. Define a map $f \times g : X \times Z \rightarrow Y \times W$ as follows:

$$\forall (x, z) \in X \times Z : [f \times g](x, z) \stackrel{\text{df}}{=} (f(x), g(z)).$$

I claim that $f \times g$ is a homeomorphism between $(X \times Z, \mathcal{T}_X \otimes \mathcal{T}_Z)$ and $(Y \times W, \mathcal{T}_Y \otimes \mathcal{T}_W)$:

(1) (Invertibility) The map $(f \times g)^{-1} : Y \times W \rightarrow X \times Z$ defined by

$$\forall (y, w) \in Y \times W : [(f \times g)^{-1}](y, w) \stackrel{\text{df}}{=} (f^{-1}(y), g^{-1}(w))$$

is an inverse for $f \times g$. For any $(x, z) \in X \times Z$, we have

$$[(f \times g)^{-1} \circ (f \times g)](x, z) = [(f \times g)^{-1}](f(x), g(z)) = (f^{-1}(f(x)), g^{-1}(g(z))) = (x, z).$$

For any $(y, w) \in Y \times W$, we have

$$[(f \times g) \circ (f \times g)^{-1}](y, w) = [f \times g](f^{-1}(y), g^{-1}(w)) = (f(f^{-1}(y)), g(g^{-1}(w))) = (y, w).$$

(2) (Continuity) Let $K \in \mathcal{T}_Y \otimes \mathcal{T}_W$. Let $(x, z) \in [(f \times g)^{-1}](K)$. This means $(f(x), g(z)) \in K$. Since $K \in \mathcal{T}_Y \otimes \mathcal{T}_W$, we can find $U \in \mathcal{T}_Y$ and $V \in \mathcal{T}_W$ such that $(f(x), g(z)) \in U \times V \subseteq K$. Taking preimages, we have

$$(x, z) \in \underbrace{f^{-1}(U)}_{\in \mathcal{T}_X} \times \underbrace{g^{-1}(V)}_{\in \mathcal{T}_Z} \subseteq [(f \times g)^{-1}](U \times V) \subseteq [(f \times g)^{-1}](K).$$

In other words, for any $(x, z) \in [(f \times g)^{-1}](K)$, there is a $\mathcal{O}_{(x,z)} \in \mathcal{T}_X \otimes \mathcal{T}_Z$ satisfying $(x, z) \in \mathcal{O}_{(x,z)} \subseteq [(f \times g)^{-1}](K)$. We use the trick

$$[(f \times g)^{-1}](K) = \bigcup_{(x,z) \in [(f \times g)^{-1}](K)} \mathcal{O}_{(x,z)},$$

as well as the Arbitrary Union property, to conclude that $[(f \times g)^{-1}](K) \in \mathcal{T}_X \otimes \mathcal{T}_Z$.

(3) (Continuity of the Inverse) The same argument as (2), with $(f \times g)$ replaced by $(f \times g)^{-1}$. □

Definition 1.13. A metric space is a pair (X, d) , where

- X is a set, and
- $d : X \times X \rightarrow \mathbb{R}$,

that satisfy the following three properties:

- (1) (Identity of Indiscernibles) $\forall x, y \in X : d(x, y) = 0 \iff x = y$.
- (2) (Symmetry) $\forall x, y \in X : d(x, y) = d(y, x)$.
- (3) (Triangle Inequality) $\forall x, y, z \in X : d(x, y) + d(y, z) \geq d(x, z)$.

Lemma 1.14 (Non-Negativity of a Metric). *Let (X, d) be a metric space. Let $x, y \in X$. Then $d(x, y) \geq 0$.*

Proof.

$$d(x, y) = \frac{d(x, y) + d(x, y)}{2} \stackrel{(2)}{=} \frac{d(x, y) + d(y, x)}{2} \stackrel{(3)}{\geq} \frac{d(x, x)}{2} \stackrel{(1)}{=} 0.$$

□

Definition 1.15. Let (X, d) be a metric space. Let $x \in X$. The ϵ -ball centered at x , denoted by “ $B_d(x, \epsilon)$ ”, is defined as:

$$B_d(x, \epsilon) \stackrel{\text{df}}{=} \{y \in X : |d(x, y) < \epsilon\}.$$

The topology associated to d , denoted by “ $\mathcal{T}_d$ ”, is defined as:

$$\mathcal{T}_d \stackrel{\text{df}}{=} \{U \subseteq X : \forall x \in U, \exists \epsilon > 0 : B_d(x, \epsilon) \subseteq U\}.$$

Lemma 1.16 (Metric Topology is a Topology). *Let (X, d) be a metric space. Then $(X, \mathcal{T}_d)$ is a topological space.*

Proof. X is a set. By definition, for any $U \in \mathcal{T}_d$, $U \subseteq X$, meaning $\mathcal{T}_d \subseteq \mathcal{P}(X)$.

(1) (Trivial Bounds) Vacuously, $\emptyset \in \mathcal{T}_d$. For any $x \in X$, there exists $\epsilon = 67 > 0$ such that $B_d(x, \epsilon) \subseteq X$, so $X \in \mathcal{T}_d$.

(2) (Arbitrary Unions) Let $\{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T}_d$. We have

$$\begin{aligned} \forall \alpha \in \mathcal{I} : \quad U_\alpha \in \mathcal{T}_d; \\ \iff \forall \alpha \in \mathcal{I}, \forall x \in U_\alpha, \exists \epsilon > 0 : \quad B_d(x, \epsilon) \subseteq U_\alpha \end{aligned}$$

Let $x \in \bigcup_{\alpha \in \mathcal{I}} U_\alpha$. We can find a particular $\alpha^* \in \mathcal{I}$ such that $x \in U_{\alpha^*}$. So,

$$\exists \epsilon > 0 : \quad B_d(x, \epsilon) \subseteq U_{\alpha^*} \subseteq \bigcup_{\alpha \in \mathcal{I}} U_\alpha.$$

Thus, $\bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}_d$.

(3) (Finite Intersections) Let $U, V \in \mathcal{T}_d$. Let $x \in U \cap V$, so $x \in U$ and $x \in V$. We can find $\epsilon_1, \epsilon_2 > 0$ such that $B_d(x, \epsilon_1) \subseteq U$ and $B_d(x, \epsilon_2) \subseteq V$. Choose $\epsilon = \min(\{\epsilon_1, \epsilon_2\})$. By how the ϵ -balls are defined, we have $B_d(x, \epsilon) \subseteq B_d(x, \epsilon_1) \subseteq U$ and $B_d(x, \epsilon) \subseteq B_d(x, \epsilon_2) \subseteq V$, so $B_d(x, \epsilon) \subseteq U \cap V$. Finally, $U \cap V \in \mathcal{T}_d$, as desired.

□

Definition 1.17. The Pythagorean Distance Function for $\mathbb{R}^n$ is the function $\text{dist}_n : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n : \quad \text{dist}_n(\mathbf{x}, \mathbf{y}) \stackrel{\text{df}}{=} \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$$

Lemma 1.18 (Euclidean Space is a Metric Space). *The pair $(\mathbb{R}^n, \text{dist}_n)$ is a metric space. The pair $(\mathbb{R}^n, \mathcal{T}_{\text{dist}_n})$ is a topological space.*

Theorem 1.19 (Compatibility of Euclidean and Product Topologies). *Let $U \in \mathcal{T}_{\text{dist}_m}$. Let $V \in \mathcal{T}_{\text{dist}_n}$. Then $\mathcal{T}_{\text{dist}_m}|_U \otimes \mathcal{T}_{\text{dist}_n}|_V = \mathcal{T}_{\text{dist}_{m+n}}|_{U \times V}$.*

Proof. We shall first show that $\mathcal{T}_{\text{dist}_m} \otimes \mathcal{T}_{\text{dist}_n} = \mathcal{T}_{\text{dist}_{m+n}}$.

($\subseteq$) Let $W \in \mathcal{T}_{\text{dist}_m} \otimes \mathcal{T}_{\text{dist}_n}$. Let $(\mathbf{x}, \mathbf{y}) \in W$. By the definition of product topology, we can find $U \in \mathcal{T}_{\text{dist}_m}$ and $V \in \mathcal{T}_{\text{dist}_n}$ such that $(\mathbf{x}, \mathbf{y}) \in U \times V \subseteq W$. Furthermore, we can find $r, s > 0$ such that $B_{\text{dist}_m}(\mathbf{x}, r) \subseteq U$ and $B_{\text{dist}_n}(\mathbf{y}, s) \subseteq V$. Choose $\epsilon = \min(\{r, s\})$ and consider

$B_{\text{dist}_{m+n}}((\mathbf{x}, \mathbf{y}), \epsilon)$. Let $(\mathbf{x}', \mathbf{y}') \in B_{\text{dist}_{m+n}}((\mathbf{x}, \mathbf{y}), \epsilon)$. Notice the following trick:

$$\begin{aligned}
\text{dist}_m(\mathbf{x}, \mathbf{x}') &\leq \sqrt{\text{dist}_m(\mathbf{x}, \mathbf{x}')^2 + \text{dist}_n(\mathbf{y}, \mathbf{y}')^2} \\
&= \sqrt{\sum_{i=1}^m (x_i - x'_i)^2 + \sum_{i=1}^n (y_i - y'_i)^2} \\
&= \sqrt{\sum_{i=1}^{m+n} (z_i - z'_i)^2} \quad \text{where } \mathbf{z} = (\mathbf{x}, \mathbf{y}) \text{ and } \mathbf{z}' = (\mathbf{x}', \mathbf{y}') \\
&= \text{dist}_{m+n}((\mathbf{x}, \mathbf{y}), (\mathbf{x}', \mathbf{y}')) \\
&< \epsilon \\
&\leq r.
\end{aligned}$$

This means $\mathbf{x}' \in B_{\text{dist}_m}(\mathbf{x}, r) \subseteq U$. Similarly, we have $\mathbf{y}' \in B_{\text{dist}_n}(\mathbf{y}, s) \subseteq V$. In sum, $(\mathbf{x}', \mathbf{y}') \in U \times V$. So, $B_{\text{dist}_{m+n}}((\mathbf{x}, \mathbf{y}), \epsilon) \subseteq U \times V \subseteq W$. We thus have satisfied the condition for $W \in \mathcal{T}_{\text{dist}_{m+n}}$.

($\supseteq$) Let $W \in \mathcal{T}_{\text{dist}_{m+n}}$. Let $(\mathbf{x}, \mathbf{y}) \in W$ with $\mathbf{x} \in \mathbb{R}^m$ and $\mathbf{y} \in \mathbb{R}^n$. We can find a $\epsilon > 0$ such that $B_{\text{dist}_{m+n}}((\mathbf{x}, \mathbf{y}), \epsilon) \subseteq W$. Consider $X = B_{\text{dist}_m}(\mathbf{x}, \frac{\epsilon}{2})$ and $Y = B_{\text{dist}_n}(\mathbf{y}, \frac{\epsilon}{2})$. Clearly $\mathbf{x} \in X \in \mathcal{T}_{\text{dist}_m}$ and $\mathbf{y} \in Y \in \mathcal{T}_{\text{dist}_n}$. Let $(\mathbf{x}', \mathbf{y}') \in X \times Y$. Notice the following trick:

$$\text{dist}_{m+n}((\mathbf{x}, \mathbf{y}), (\mathbf{x}', \mathbf{y}')) = \sqrt{\text{dist}_m(\mathbf{x}, \mathbf{x}')^2 + \text{dist}_n(\mathbf{y}, \mathbf{y}')^2} \leq \sqrt{\left(\frac{\epsilon}{2}\right)^2 + \left(\frac{\epsilon}{2}\right)^2} < \epsilon.$$

This means, $(\mathbf{x}', \mathbf{y}') \in B_{\text{dist}_{m+n}}((\mathbf{x}, \mathbf{y}), \epsilon)$, meaning $X \times Y \subseteq B_{\text{dist}_{m+n}}((\mathbf{x}, \mathbf{y}), \epsilon) \subseteq W$. So, $(\mathbf{x}, \mathbf{y}) \in X \times Y \subseteq W$, yielding $W \in \mathcal{T}_{\text{dist}_m} \otimes \mathcal{T}_{\text{dist}_n}$.

At last, the original proposition can be proven by citing 1.7: Let $U \in \mathcal{T}_{\text{dist}_m}$ and $V \in \mathcal{T}_{\text{dist}_n}$. Since $(\mathbb{R}^m, \mathcal{T}_{\text{dist}_m})$ and $(\mathbb{R}^n, \mathcal{T}_{\text{dist}_n})$ are topological spaces, we have

$$\mathcal{T}_{\text{dist}_m}|_U \otimes \mathcal{T}_{\text{dist}_n}|_V = (\mathcal{T}_{\text{dist}_m} \otimes \mathcal{T}_{\text{dist}_n})|_{U \times V} = \mathcal{T}_{\text{dist}_{m+n}}|_{U \times V}.$$

□

2. TOPOLOGICAL MANIFOLDS

Definition 2.1. A topological manifold of dimension n is a pair $(M, \mathcal{T})$, where

- $(M, \mathcal{T})$ is a topological space

that satisfies the following three properties:

- (1) (Hausdorffness) $\forall p \in M, \forall q \in M \setminus \{p\}, \exists U, V \in \mathcal{T} : p \in U \text{ and } q \in V \text{ and } U \cap V = \emptyset$.
- (2) (Second-Countability) $\exists \{B_i\}_{i \in \mathbb{N}} \subseteq \mathcal{T}, \forall U \in \mathcal{T}, \forall p \in U, \exists i \in \mathbb{N} : p \in B_i \subseteq U$.
- (3) (Local Euclideaness of Dimension n)

$$\forall p \in M, \exists U \in \{U \in \mathcal{T} : p \in U\}, \exists V \in \mathcal{T}_{\text{dist}_n} : (U, \mathcal{T}|_U) \cong (V, \mathcal{T}_{\text{dist}_n}|_V).$$

Proposition 2.2 (Product of Topological Manifolds). *Let $(M, \mathcal{T}_M)$ and $(N, \mathcal{T}_N)$ be topological manifolds of dimensions m and n respectively. Then $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)$ is a topological manifold of dimension $m + n$.*

Proof. By 1.3, we know that $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)$ is a topological space. It remains to check that it satisfies the three properties.

- (1) (Hausdorffness) Let $(p_1, p_2) \in M \times N$. Let $(q_1, q_2) \in M \times N \setminus \{(p_1, p_2)\}$. WLoG, assume $p_1 \neq q_1$. We can find $U, V \in \mathcal{T}_M$ such that $p_1 \in U$, $q_1 \in V$, and $U \cap V = \emptyset$. Consequently, we have found $U \times N, V \times N \in \mathcal{T}_M \otimes \mathcal{T}_N$ satisfying $(p_1, p_2) \in U \times N$, $(q_1, q_2) \in V \times N$, and $(U \times N) \cap (V \times N) = (U \cap V) \times (N \cap N) = \emptyset \times N = \emptyset$.
- (2) (Second-Countability) Let $\{B_i\}_{i \in \mathbb{N}} \subseteq \mathcal{T}_M$ satisfy for any $U \in \mathcal{T}_M$ and for any $p \in U$, there is an $i \in \mathbb{N}$ such that $p \in B_i \subseteq U$. Let $\{C_j\}_{j \in \mathbb{N}} \subseteq \mathcal{T}_N$ satisfy for any $V \in \mathcal{T}_N$ and for any $q \in V$, there is a $j \in \mathbb{N}$ such that $q \in C_j \subseteq V$. Consider the collection $\{B_i \times C_j\}_{i,j \in \mathbb{N}} \subseteq \mathcal{T}_M \otimes \mathcal{T}_N$. Let $W \in \mathcal{T}_M \otimes \mathcal{T}_N$. Let $(p, q) \in W$. Recall from 1.2, we can find $U \in \mathcal{T}_M$ and $V \in \mathcal{T}_N$ satisfying $(p, q) \in U \times V \subseteq W$. By above, we can further find $i, j \in \mathbb{N}$ such that $p \in B_i \subseteq U$ and $q \in C_j \subseteq V$. So, $(p, q) \in B_i \times C_j \subseteq U \times V \subseteq W$.
- (3) (Local Euclideaness of Dimension $m + n$) Let $(p, q) \in M \times N$. Since $p \in M$, we can find a $U_1 \in \mathcal{T}_M$ with $p \in U_1$ and a $V_1 \in \mathcal{T}_{\text{dist}_m}$ such that $(U_1, \mathcal{T}_M|_{U_1}) \cong (V_1, \mathcal{T}_{\text{dist}_m}|_{V_1})$. Since $q \in N$, we can find a $U_2 \in \mathcal{T}_N$ with $q \in U_2$ and a $V_2 \in \mathcal{T}_{\text{dist}_n}$ such that $(U_2, \mathcal{T}_N|_{U_2}) \cong (V_2, \mathcal{T}_{\text{dist}_n}|_{V_2})$. By 1.12, we know that

$$(U_1 \times U_2, \mathcal{T}_M|_{U_1} \otimes \mathcal{T}_N|_{U_2}) \cong (V_1 \times V_2, \mathcal{T}_{\text{dist}_m}|_{V_1} \otimes \mathcal{T}_{\text{dist}_n}|_{V_2}).$$

By 1.7 on the LHS and 1.19 on the RHS, this is equivalent to:

$$(U_1 \times U_2, (\mathcal{T}_M \otimes \mathcal{T}_N)|_{U_1 \times U_2}) \cong (V_1 \times V_2, \mathcal{T}_{\text{dist}_{m+n}}|_{V_1 \times V_2}).$$

□

Proposition 2.3 (Open Subspace of a Topological Manifold). *Let $(M, \mathcal{T})$ be a topological manifold of dimension n . Let $W \in \mathcal{T}$. Then $(W, \mathcal{T}|_W)$ is a topological manifold of dimension n .*

Proof. By 1.5, we know that $(W, \mathcal{T}|_W)$ is a topological space. It remains to check that it satisfies the three properties.

- (1) (Hausdorffness) Let $p \in W$. Let $q \in W \setminus \{p\}$. This means $p \in M$ and $q \in M \setminus \{p\}$. We can find $U, V \in \mathcal{T}$ such that $p \in U$, $q \in V$ and $U \cap V = \emptyset$. Note that $p \in U \cap W$ and $q \in V \cap W$. At the end, $(U \cap W) \cap (V \cap W) = (U \cap V) \cap (W \cap W) = \emptyset \cap W = \emptyset$, and we are done.
- (2) (Second-Countability) Let $\{B_i\}_{i \in \mathbb{N}} \subseteq \mathcal{T}$ satisfy for any $U \in \mathcal{T}$ and for any $p \in U$, there is an $i \in \mathbb{N}$ such that $p \in B_i \subseteq U$. Let $U \cap W \in \mathcal{T}|_W$ for some $U \in \mathcal{T}$. Let $p \in U \cap W$. Remember that $W \in \mathcal{T}$, so $U \cap W \in \mathcal{T}$. We can find an $i \in \mathbb{N}$ such that $p \in B_i \subseteq U \cap W$. I can also say that $p \in B_i \cap W \subseteq U \cap W$. So, the collection $\{B_i \cap W\}_{i \in \mathbb{N}}$ is a countable basis for $(W, \mathcal{T}|_W)$.
- (3) (Local Euclideaness of Dimension n) Let $p \in W$. So $p \in M$. We can find $U \in \mathcal{T}$ with $p \in U$ and $V \in \mathcal{T}_{\text{dist}_n}$ such that $(U, \mathcal{T}|_U) \cong (V, \mathcal{T}_{\text{dist}_n}|_V)$. Let $\varphi : U \rightarrow V$ be a homeomorphism between $(U, \mathcal{T}|_U)$ and $(V, \mathcal{T}_{\text{dist}_n}|_V)$. By 1.11, since $U \cap W \subseteq U$, we have $(U \cap W, (\mathcal{T}|_U)|_{U \cap W}) \cong (\varphi(U \cap W), (\mathcal{T}_{\text{dist}_n}|_V)|_{\varphi(U \cap W)})$. By 1.6, we have the following chain of equivalences:

$$\begin{aligned} (U \cap W, (\mathcal{T}|_U)|_{U \cap W}) &\cong (\varphi(U \cap W), (\mathcal{T}_{\text{dist}_n}|_V)|_{\varphi(U \cap W)}) \\ (U \cap W, \mathcal{T}|_{U \cap W}) &\cong (\varphi(U \cap W), \mathcal{T}_{\text{dist}_n}|_{V \cap \varphi(U \cap W)}) \\ (U \cap W, \mathcal{T}|_{U \cap W}) &\cong (\varphi(U \cap W), \mathcal{T}_{\text{dist}_n}|_{\varphi(U \cap W)}) \\ (U \cap W, (\mathcal{T}|_U)|_W) &\cong (\varphi(U \cap W), \mathcal{T}_{\text{dist}_n}|_{\varphi(U \cap W)}). \end{aligned}$$

Thus $U \cap W \in \mathcal{T}|_W$ with $p \in U \cap W$, and $\varphi(U \cap W) \in \mathcal{T}_{\text{dist}_n}$, since $U \cap W \in \mathcal{T}$ and φ is a homeomorphism that maps open to open.

□

3. SMOOTH MANIFOLDS

Definition 3.1. Let $U \in \mathcal{T}_{\text{dist}_m}$. A map $\mathbf{f} : U \rightarrow \mathbb{R}^n$ is said to be $\underline{C^\infty}$ iff:

$$\forall k_1, \dots, k_m \in \mathbb{Z}_{\geq 0}, \forall i \in \{1, \dots, n\} : \quad \frac{\partial^{k_1+\dots+k_m} f_i}{\partial x_1^{k_1} \dots \partial x_m^{k_m}} \text{ exists.}$$

Lemma 3.2 (Restriction of a C^∞ Map). *Let $\mathbf{f}$ be C^∞ . Let $U \subseteq \text{Dom}(\mathbf{f})$ and $U \in \mathcal{T}_{\text{dist}_m}$. Then $\mathbf{f}|_U$ is C^∞ .*

Definition 3.3. Let $(M, \mathcal{T})$ be a topological manifold of dimension n . Let $U \in \mathcal{T}$ be an open set in M . Let $\varphi : U \rightarrow \mathbb{R}^n$ be a coordinate map for U . The pair (U, φ) is said to be an $\underline{(M, \mathcal{T})}$ -chart iff $\varphi : U \rightarrow \varphi(U)$ is a homeomorphism between $(U, \mathcal{T}|_U)$ and $(\varphi(U), \mathcal{T}_{\text{dist}_n}|_{\varphi(U)})$.

Definition 3.4. A smooth manifold of dimension n is a triplet $(M, \mathcal{T}, \mathcal{A})$, where

- $(M, \mathcal{T})$ is a topological manifold of dimension n , and
- $\mathcal{A} \stackrel{\text{df}}{=} \{(U_\alpha, \varphi_\alpha)\}_{\alpha \in \mathcal{I}}$ is called a smooth structure on $(M, \mathcal{T})$,

that satisfy the following three properties:

- (1) (Atlas) $\forall (U, \varphi) \in \mathcal{A} : (U, \varphi)$ is an $(M, \mathcal{T})$ -chart and $\bigcup_{(U, \varphi) \in \mathcal{A}} U = M$.
- (2) (Smoothness) $\forall (U, \varphi), (V, \psi) \in \mathcal{A} : (\psi \circ \varphi^{-1})|_{\varphi(U \cap V)}$ is C^∞ .
- (3) (Maximality) $\left\{ (U, \varphi) : \begin{array}{l} (U, \varphi) \text{ is an } (M, \mathcal{T})\text{-chart} \\ \forall (V, \psi) \in \mathcal{A} : \\ (\psi \circ \varphi^{-1})|_{\varphi(U \cap V)} \text{ and } (\varphi \circ \psi^{-1})|_{\psi(U \cap V)} \text{ are } C^\infty \end{array} \right\} \subseteq \mathcal{A}$.

Definition 3.5. Let $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$ be smooth manifolds. The product smooth structure on $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)$, denoted by “ $\mathcal{A}_M \otimes \mathcal{A}_N$ ”, is defined as:

$$\mathcal{A}_M \otimes \mathcal{A}_N \stackrel{\text{df}}{=} \mathcal{P} \cup \mathcal{Q}$$

where

$$\mathcal{P} = \{(U \times V, \varphi \times \psi) : (U, \varphi) \in \mathcal{A}_M \text{ and } (V, \psi) \in \mathcal{A}_N\}$$

and

$$\mathcal{Q} = \left\{ (W, \chi) : \begin{array}{l} (W, \chi) \text{ is an } (M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)\text{-chart} \\ (W, \chi) \notin \mathcal{P} \\ \forall (U \times V, \varphi \times \psi) \in \mathcal{P} : \\ \chi \circ (\varphi \times \psi)^{-1} : [\varphi \times \psi]((U \times V) \cap W) \rightarrow \chi((U \times V) \cap W) \text{ is } C^\infty \\ (\varphi \times \psi) \circ \chi^{-1} : \chi((U \times V) \cap W) \rightarrow [\varphi \times \psi]((U \times V) \cap W) \text{ is } C^\infty \end{array} \right\}.$$

Theorem 3.6 (Product of Smooth Manifolds). *Let $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$ be smooth manifolds of dimensions m and n respectively. Then $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N, \mathcal{A}_M \otimes \mathcal{A}_N)$ is a smooth manifold of dimension $m + n$.*

Proof. By 2.2, we know that $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)$ is a topological manifold of dimension $m + n$. It remains to check that $\mathcal{A}_M \otimes \mathcal{A}_N$ satisfy the three properties.

- (1) (Atlas) By design, any $(W, \chi) \in \mathcal{Q} \subseteq \mathcal{A}_M \otimes \mathcal{A}_N$ is an $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)$ -chart. Otherwise, let $(U \times V, \varphi \times \psi) \in \mathcal{P} \subseteq \mathcal{A}_M \otimes \mathcal{A}_N$. We know that (U, φ) is an $(M, \mathcal{T}_M)$ -chart and (V, ψ) is an $(N, \mathcal{T}_N)$ -chart. So, $\varphi : U \rightarrow \varphi(U)$ is a homeomorphism between $(U, \mathcal{T}|_U)$ and

$(\varphi(U), \mathcal{T}_{\text{dist}_m}|_{\varphi(U)})$. Similarly, $\psi : V \rightarrow \psi(V)$ is a homeomorphism between $(V, \mathcal{T}|_V)$ and $(\psi(V), \mathcal{T}_{\text{dist}_n}|_{\psi(V)})$. By the same argument used in the proof for 1.12, $\varphi \times \psi : U \times V \rightarrow \varphi(U) \times \psi(V) = [\varphi \times \psi](U \times V) \subseteq \mathbb{R}^{m+n}$ is a homeomorphism between $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)$ and $(\varphi(U) \times \psi(V), \mathcal{T}_{\text{dist}_m}|_{\varphi(U)} \otimes \mathcal{T}_{\text{dist}_n}|_{\psi(V)})$. Now notice the following:

$$\begin{aligned} (\varphi(U) \times \psi(V), \mathcal{T}_{\text{dist}_m}|_{\varphi(U)} \otimes \mathcal{T}_{\text{dist}_n}|_{\psi(V)}) &= (\varphi(U) \times \psi(V), \mathcal{T}_{\text{dist}_{m+n}}|_{\varphi(U) \times \psi(V)}) \\ &= ([\varphi \times \psi](U \times V), \mathcal{T}_{\text{dist}_{m+n}}|_{[\varphi \times \psi](U \times V)}). \end{aligned}$$

Indeed, $(U \times V, \varphi \times \psi)$ is an $(M \times N, \mathcal{T}_M \otimes \mathcal{T}_N)$ -chart. Also, we have

$$\bigcup_{(\alpha, \beta) \in \mathcal{I} \times \mathcal{J}} U_\alpha \times V_\beta = \bigcup_{\alpha \in \mathcal{I}} U_\alpha \times \bigcup_{\beta \in \mathcal{J}} V_\beta = M \times N.$$

(2) (Smoothness) There are three cases:

(a) (Both charts in $\mathcal{P}$) Let $(U \times V, \varphi \times \psi), (U' \times V', \varphi' \times \psi') \in \mathcal{P}$. We have the following:

$$\begin{aligned} (\varphi' \times \psi') \circ (\varphi \times \psi)^{-1} &= (\varphi' \times \psi') \circ (\varphi^{-1} \times \psi^{-1}) \\ &= \underbrace{(\varphi' \circ \varphi^{-1})}_{C^\infty} \times \underbrace{(\psi' \circ \psi^{-1})}_{C^\infty}. \end{aligned}$$

(b) (One chart in $\mathcal{P}$ and one chart in $\mathcal{Q}$) The transition map is C^∞ by design.

(c) (Both charts in $\mathcal{Q}$) Let $(W, \chi), (W', \chi') \in \mathcal{Q}$. Let $p \in W \cap W'$. Since $\mathcal{P}$ covers $M \times N$ as shown in (1), we can find a chart $(U \times V, \varphi \times \psi) \in \mathcal{P}$ with $p \in U \times V$. On the neighborhood $\chi((U \times V) \cap W \cap W')$, we have the following:

$$\chi' \circ \chi^{-1} = \underbrace{(\chi' \circ (\varphi \times \psi)^{-1})}_{C^\infty} \circ \underbrace{((\varphi \times \psi) \circ \chi^{-1})}_{C^\infty}.$$

This argument could be repeated for any point $p \in W \cap W'$. So, $\chi' \circ \chi^{-1} : \chi(W \cap W') \rightarrow \chi'(W \cap W')$ is C^∞ .

(3) (Maximality) This is by design. □

Definition 3.7. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $W \in \mathcal{T}$. The restricted smooth structure on W , denoted by $\mathcal{A}|_W$, is defined as:

$$\mathcal{A}|_W \stackrel{\text{df}}{=} \{(U, \varphi) \in \mathcal{A} : U \subseteq W\}.$$

Lemma 3.8 (Restriction of a Chart). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $W \in \mathcal{T}$. Let $(U, \varphi) \in \mathcal{A}$. Then $(U \cap W, \varphi|_{U \cap W}) \in \mathcal{A}$. Consequently, $(U \cap W, \varphi|_{U \cap W}) \in \mathcal{A}|_W$.*

Proof. (1) $((U \cap W, \varphi|_{U \cap W})$ is an $(M, \mathcal{T})$ -chart) Since $U, W \in \mathcal{T}$, $U \cap W \in \mathcal{T}$. Moreover, by the same reasoning used in the proof for 1.11, it can be seen that $\varphi|_{U \cap W} : U \cap W \rightarrow \varphi(U \cap W)$ is a homeomorphism between $(U \cap W, (\mathcal{T}|_U)|_{U \cap W})$ and $(\varphi(U \cap W), (\mathcal{T}_{\text{dist}_n}|_{\varphi(U)})|_{\varphi(U \cap W)})$. By 1.6 and the fact that φ is injective, $\varphi|_{U \cap W} : U \cap W \rightarrow \varphi(U \cap W)$ is a homeomorphism between $(U \cap W, \mathcal{T}|_{U \cap W})$ and $(\varphi(U \cap W), \mathcal{T}_{\text{dist}_n}|_{\varphi(U \cap W)})$.

- (2) $((U \cap W, \varphi|_{U \cap W})$ is smoothly compatible with every chart in $\mathcal{A}$) Let $(V, \psi) \in \mathcal{A}$. We have the following:

$$\begin{aligned}\psi \circ (\varphi|_{U \cap W})^{-1} &= \psi \circ \varphi^{-1}|_{\varphi(U \cap W)} = \underbrace{(\psi \circ \varphi^{-1})}_{C^\infty}|_{\varphi(U \cap V \cap W)}; \\ \varphi|_{U \cap W} \circ \psi^{-1} &= \underbrace{(\varphi \circ \psi^{-1})}_{C^\infty}|_{\psi(U \cap V \cap W)}.\end{aligned}$$

By the maximality of $\mathcal{A}$, we have $(U \cap W, \varphi|_{U \cap W}) \in \mathcal{A}$. Consequently, since $U \in \mathcal{T}$, $U \cap W \in \mathcal{T}|_W$, so $(U \cap W, \varphi|_{U \cap W}) \in \mathcal{A}|_W$. $\square$

Theorem 3.9 (Open Subspace of a Smooth Manifold). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold of dimension n . Let $W \in \mathcal{T}$. Then $(W, \mathcal{T}|_W, \mathcal{A}|_W)$ is a smooth manifold of dimension n .*

Proof. By 2.3, we know that $(W, \mathcal{T}|_W)$ is a topological manifold of dimension n . It remains to check that $\mathcal{A}|_W$ is a smooth structure on $(W, \mathcal{T}|_W)$.

- (1) (Atlas) Let $(U, \varphi) \in \mathcal{A}|_W$. Since $U \subseteq W \in \mathcal{T}$, we have $U = U \cap W \in \mathcal{T}|_W$. Also, $\varphi : U \rightarrow \varphi(U)$ is by definition a homeomorphism between $(U, \mathcal{T}|_U)$ and $(\varphi(U), \mathcal{T}_{\text{dist}_n}|_{\varphi(U)})$. Thus, (U, φ) is a $(W, \mathcal{T}|_W)$ -chart. The coverage is done as follows:

$$\bigcup_{(U, \varphi) \in \mathcal{A}|_W} U = \bigcup_{U \subseteq W} U = W.$$

- (2) (Smoothness) Smooth compatibility is inherited from $\mathcal{A}$, since $\mathcal{A}|_W \subseteq \mathcal{A}$.

- (3) (Maximality) Let (U, φ) be a $(W, \mathcal{T}|_W)$ -chart that is smoothly compatible with every chart in $\mathcal{A}|_W$. Let $(V, \psi) \in \mathcal{A}$. By 3.8, since $W \in \mathcal{T}$, we have $(V \cap W, \psi|_{V \cap W}) \in \mathcal{A}|_W$. So, (U, φ) and $(V \cap W, \psi|_{V \cap W})$ are smoothly compatible. Now, noticing that $U \cap V = U \cap V \cap W$ because $U \subseteq W$, we have

$$\begin{aligned}(\psi \circ \varphi^{-1})|_{\varphi(U \cap V)} &= (\psi \circ \varphi^{-1})|_{\varphi(U \cap V \cap W)} = \underbrace{(\psi|_{V \cap W} \circ \varphi^{-1})}_{C^\infty}|_{\varphi(U \cap (V \cap W))}; \\ (\varphi \circ \psi^{-1})|_{\psi(U \cap V)} &= (\varphi \circ \psi^{-1})|_{\psi(U \cap V \cap W)} = \underbrace{(\varphi \circ (\psi|_{V \cap W})^{-1})}_{C^\infty}|_{\psi(U \cap (V \cap W))}.\end{aligned}$$

This means (U, φ) is smoothly compatible with every chart in $\mathcal{A}$. Since $U \in \mathcal{T}|_W$ and $W \in \mathcal{T}$, $U \in \mathcal{T}$ and (U, φ) is an $(M, \mathcal{T})$ -chart. By the maximality of $\mathcal{A}$, $(U, \varphi) \in \mathcal{A}$. Since $U \subseteq W$, $(U, \varphi) \in \mathcal{A}|_W$. $\square$

Definition 3.10. Let $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$ be smooth manifolds. A map $F : M \rightarrow N$ is said to be smooth w.r.t. $\mathcal{A}_M$ and $\mathcal{A}_N$ iff:

$$\begin{aligned}\forall p \in M, \forall (V, \psi) \in \{(V, \psi) \in \mathcal{A}_N : F(p) \in V\}, \\ \exists (U, \varphi) \in \{(U, \varphi) \in \mathcal{A}_M : p \in U \text{ and } F(U) \subseteq V\} : \\ \psi \circ F \circ \varphi^{-1} : \varphi(U) \rightarrow \psi(V) \text{ is } C^\infty.\end{aligned}$$

Lemma 3.11 (Simpler Smoothness Criterion). *Let $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$ be smooth manifolds. Let $F : M \rightarrow N$. Then F is smooth w.r.t. $\mathcal{A}_M$ and $\mathcal{A}_N$ iff*

$$\begin{aligned}\forall p \in M, \exists (U, \varphi) \in \mathcal{A}_M, \exists (V, \psi) \in \mathcal{A}_N : \\ p \in U \text{ and } F(U) \subseteq V \text{ and } \psi \circ F \circ \varphi^{-1} : \varphi(U) \rightarrow \psi(V) \text{ is } C^\infty.\end{aligned}$$

Proof. The forward direction follows directly from the definition. We prove the converse.

($\Leftarrow$) Let $p \in M$. Let $(W, \chi) \in \mathcal{A}_N$ with $F(p) \in W$. Since $p \in M$, by assumption, we can find $(U, \varphi) \in \mathcal{A}_M$ and $(V, \psi) \in \mathcal{A}_N$ such that $p \in U$, $F(U) \subseteq V$ and $\psi \circ F \circ \varphi^{-1}$ is C^∞ . Notice that $F(p) \in V \cap W$. Because φ and ψ are homeomorphisms, $F|_U$ is continuous w.r.t $\mathcal{T}_M|_U$ and $\mathcal{T}_N|_{F(U)}$. By 3.8, $(U \cap F^{-1}(V \cap W), \varphi|_{U \cap F^{-1}(V \cap W)}) \in \mathcal{A}_M$, with $p \in U \cap F^{-1}(V \cap W)$, and $F(U \cap F^{-1}(V \cap W)) = F(U) \cap V \cap W \subseteq W$. We also have

$$\chi \circ F \circ \left(\varphi|_{U \cap F^{-1}(V \cap W)} \right)^{-1} = \underbrace{(\chi \circ \psi^{-1})}_{C^\infty} \circ \underbrace{(\psi \circ F \circ \varphi^{-1})}_{C^\infty} \Big|_{\varphi(U \cap F^{-1}(V \cap W))}.$$

□

Definition 3.12. Let $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$ be smooth manifolds. A map $F : M \rightarrow N$ is said to be a diffeomorphism between $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$ iff it satisfies the following three properties:

- (1) (Invertibility) $F : M \rightarrow N$ is a bijection.
- (2) (Smoothness) $F : M \rightarrow N$ is smooth w.r.t. $\mathcal{A}_M$ and $\mathcal{A}_N$.
- (3) (Smoothness of the Inverse) $F^{-1} : N \rightarrow M$ is smooth w.r.t. $\mathcal{A}_N$ and $\mathcal{A}_M$.

Definition 3.13. A smooth manifold $(M, \mathcal{T}_M, \mathcal{A}_M)$ is said to be diffeomorphic to another smooth manifold $(N, \mathcal{T}_N, \mathcal{A}_N)$, denoted by “ $(M, \mathcal{T}_M, \mathcal{A}_M) \simeq (N, \mathcal{T}_N, \mathcal{A}_N)$ ”, iff there exists a diffeomorphism between $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$.

4. ALGEBRAIC STRUCTURES

Definition 4.1. An algebraic signature is a pair $(\mathcal{I}, \text{ar})$, where

- $\mathcal{I}$ is an index set, and
- $\text{ar} : \mathcal{I} \rightarrow \mathbb{Z}_{\geq 0}$ assigns each index to a nonnegative integer.

Definition 4.2. Let $\sigma = (\mathcal{I}, \text{ar})$ be an algebraic signature. A σ -structure is a pair $(A, \{\mu_i\}_{i \in \mathcal{I}})$, where

- A is a set, and
- $\forall i \in \mathcal{I} : \mu_i : A^{\text{ar}(i)} \rightarrow A$ is an $\text{ar}(i)$ -ary operation.

Definition 4.3. Let $(A, \{\mu_i^A\}_{i \in \mathcal{I}})$ and $(B, \{\mu_i^B\}_{i \in \mathcal{I}})$ be σ -structures. A map $\varphi : A \rightarrow B$ is said to be a σ -homomorphism iff

$$\forall i \in \mathcal{I}, \forall a_1, \dots, a_{\text{ar}(i)} \in A : \varphi(\mu_i^A(a_1, \dots, a_{\text{ar}(i)})) = \mu_i^B(\varphi(a_1), \dots, \varphi(a_{\text{ar}(i)})).$$

Definition 4.4. Let $(A, \{\mu_i^A\}_{i \in \mathcal{I}})$ and $(B, \{\mu_i^B\}_{i \in \mathcal{I}})$ be σ -structures. A map $\varphi : A \rightarrow B$ is said to be a σ -isomorphism iff it satisfies the following three properties:

- (1) $\varphi : A \rightarrow B$ is a bijection.
- (2) $\varphi : A \rightarrow B$ is a σ -homomorphism.
- (3) $\varphi^{-1} : B \rightarrow A$ is a σ -homomorphism.

Theorem 4.5 (a Concise Characterization for Isomorphisms). *Let*

$$\left(A, \{\mu_i^A\}_{i \in \mathcal{I}}\right) \quad \text{and} \quad \left(B, \{\mu_i^B\}_{i \in \mathcal{I}}\right)$$

be σ -structures. A map $\varphi : A \rightarrow B$ is a σ -isomorphism iff the following two statements hold:

- (1) $\varphi : A \rightarrow B$ is a bijection.
- (2) $\varphi : A \rightarrow B$ is a σ -homomorphism.

Proof. ($\implies$) The forward direction is logically trivial.

($\impliedby$) We want to show that $\varphi^{-1} : B \rightarrow A$ is a σ -homomorphism. Let $i \in \mathcal{I}$. Let $b_1, \dots, b_{\text{ar}(i)} \in B$. Since $\varphi : A \rightarrow B$ is a bijection, we can find uniquely $a_1, \dots, a_{\text{ar}(i)} \in A$ such that $\varphi(a_k) = b_k$ for every $k \in \{1, \dots, \text{ar}(i)\}$. We thus have the following:

$$\begin{aligned} \varphi^{-1}(\mu_i^B(b_1, \dots, b_{\text{ar}(i)})) &= \varphi^{-1}(\mu_i^B(\varphi(a_1), \dots, \varphi(a_{\text{ar}(i)}))) \\ &= \varphi^{-1}(\varphi(\mu_i^A(a_1, \dots, a_{\text{ar}(i)}))) \\ &= \mu_i^A(a_1, \dots, a_{\text{ar}(i)}) \\ &= \mu_i^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{\text{ar}(i)})) \end{aligned}$$

□

Lemma 4.6 (Transport of Structures). *Let $\left(A, \{\mu_i^A\}_{i \in \mathcal{I}}\right)$ be a σ -structure. Let $\{k_j\}_{j \in \mathcal{J}} \subseteq \mathbb{Z}_{\geq 0}$. For each $j \in \mathcal{J}$, define maps $s_j^A, t_j^A : A^{k_j} \rightarrow A$ satisfying:*

$$\forall a_1, \dots, a_{k_j} \in A : \quad s_j^A(a_1, \dots, a_{k_j}) = t_j^A(a_1, \dots, a_{k_j}).$$

Let B be a set. Let $\varphi : A \rightarrow B$ be a bijection. For each $i \in \mathcal{I}$, define an $\text{ar}(i)$ -ary operation $\mu_i^B : B^{\text{ar}(i)} \rightarrow B$ by:

$$\forall b_1, \dots, b_{\text{ar}(i)} \in B : \quad \mu_i^B(b_1, \dots, b_{\text{ar}(i)}) \stackrel{\text{df}}{=} \varphi(\mu_i^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{\text{ar}(i)}))).$$

Further, for each $j \in \mathcal{J}$, define maps $s_j^B, t_j^B : B^{k_j} \rightarrow B$ by:

$$\forall b_1, \dots, b_{k_j} \in B : \quad \begin{cases} s_j^B(b_1, \dots, b_{k_j}) \stackrel{\text{df}}{=} \varphi(s_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j}))); \\ t_j^B(b_1, \dots, b_{k_j}) \stackrel{\text{df}}{=} \varphi(t_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j}))). \end{cases}$$

Then the following three statements hold:

- (1) $\left(B, \{\mu_i^B\}_{i \in \mathcal{I}}\right)$ is a σ -structure.
- (2) $\varphi : A \rightarrow B$ is a σ -isomorphism.
- (3) $\forall j \in \mathcal{J}, \forall b_1, \dots, b_{k_j} \in B : \quad s_j^B(b_1, \dots, b_{k_j}) = t_j^B(b_1, \dots, b_{k_j}).$

Proof. (1) B is a set by assumption. For each $i \in \mathcal{I}$, $\mu_i^B : B^{\text{ar}(i)} \rightarrow B$ has the correct type.

(2) Since $\varphi : A \rightarrow B$ is a bijection, by 4.5, we just need to show that $\varphi : A \rightarrow B$ is a σ -homomorphism. Let $i \in \mathcal{I}$. Let $a_1, \dots, a_{\text{ar}(i)} \in A$. We can find $b_1, \dots, b_{\text{ar}(i)} \in B$ such that $b_k = \varphi(a_k)$ for every $k \in \{1, \dots, \text{ar}(i)\}$. We then have the following:

$$\begin{aligned} \varphi(\mu_i^A(a_1, \dots, a_{\text{ar}(i)})) &= \varphi(\mu_i^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{\text{ar}(i)}))) \\ &= \mu_i^B(b_1, \dots, b_{\text{ar}(i)}) \\ &= \mu_i^B(\varphi(a_1), \dots, \varphi(a_{\text{ar}(i)})). \end{aligned}$$

- (3) Let $j \in \mathcal{J}$. Let $b_1, \dots, b_{k_j} \in B$. Since $\varphi : A \rightarrow B$ is a bijection, we can find $a_1, \dots, a_{k_j} \in A$ such that $b_i = \varphi(a_i)$ for every $i \in \{1, \dots, k_j\}$. We have the following:

$$\begin{aligned} s_j^A(a_1, \dots, a_{k_j}) &= t_j^A(a_1, \dots, a_{k_j}) \\ s_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j})) &= t_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j})) \\ \varphi(s_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j}))) &= \varphi(t_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j}))) \\ s_j^B(b_1, \dots, b_{k_j}) &= t_j^B(b_1, \dots, b_{k_j}). \end{aligned}$$

□

Corollary 4.7 (Partial Transport). *Let $(A, \{\mu_i^A\}_{i \in \mathcal{I}})$ be a σ -structure. Let $\{k_j\}_{j \in \mathcal{J}} \subseteq \mathbb{Z}_{\geq 0}$. For each $j \in \mathcal{J}$, define maps $s_j^A, t_j^A : A^{k_j} \rightarrow A$ satisfying:*

$$\forall a_1, \dots, a_{k_j} \in A : s_j^A(a_1, \dots, a_{k_j}) = t_j^A(a_1, \dots, a_{k_j}).$$

Let $\mathcal{K} \subseteq \mathcal{I}$. Let $\tau = (\mathcal{K}, \text{ar}|_{\mathcal{K}})$ be a signature. Let $(B, \{\mu_k^B\}_{k \in \mathcal{K}})$ be a τ -structure. Let $\varphi : A \rightarrow B$ be a bijective τ -homomorphism. For each $i \in \mathcal{I} \setminus \mathcal{K}$, define an $\text{ar}(i)$ -ary operation $\mu_i^B : B^{\text{ar}(i)} \rightarrow B$ by:

$$\forall b_1, \dots, b_{\text{ar}(i)} \in B : \mu_i^B(b_1, \dots, b_{\text{ar}(i)}) \stackrel{\text{df}}{=} \varphi(\mu_i^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{\text{ar}(i)}))).$$

Further, for each $j \in \mathcal{J}$, define maps $s_j^B, t_j^B : B^{k_j} \rightarrow B$ by:

$$\forall b_1, \dots, b_{k_j} \in B : \begin{cases} s_j^B(b_1, \dots, b_{k_j}) \stackrel{\text{df}}{=} \varphi(s_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j}))); \\ t_j^B(b_1, \dots, b_{k_j}) \stackrel{\text{df}}{=} \varphi(t_j^A(\varphi^{-1}(b_1), \dots, \varphi^{-1}(b_{k_j}))). \end{cases}$$

Then the following three statements hold:

- (1) $(B, \{\mu_i^B\}_{i \in \mathcal{I}})$ is a σ -structure.
- (2) $\varphi : A \rightarrow B$ is a σ -isomorphism.
- (3) $\forall j \in \mathcal{J}, \forall b_1, \dots, b_{k_j} \in B : s_j^B(b_1, \dots, b_{k_j}) = t_j^B(b_1, \dots, b_{k_j})$.

Definition 4.8. A group is a pair $(G, \{\mu, \iota, e\})$, where

- G is a set,
- $\mu : G \times G \rightarrow G$ is a binary operation on G ,
- $\iota : G \rightarrow G$ is the inversion map, and
- $e \in G$ is the identity element,

that satisfy the following three properties:

- (1) (Associativity) $\forall x, y, z \in G : \mu(\mu(x, y), z) = \mu(x, \mu(y, z))$.
- (2) (Identity) $\forall x \in G : \mu(e, x) = x = \mu(x, e)$.
- (3) (Inverse) $\forall x \in G : \mu(\iota(x), x) = e = \mu(x, \iota(x))$.

Definition 4.9. Let $(G, \{\mu_G, \iota_G, e_G\})$ and $(H, \{\mu_H, \iota_H, e_H\})$ be groups. A map $\varphi : G \rightarrow H$ is said to be a group homomorphism iff $\forall x, y \in G : \varphi(\mu_G(x, y)) = \mu_H(\varphi(x), \varphi(y))$. The kernel of φ , denoted by “ $\ker(\varphi)$ ”, is defined as:

$$\ker(\varphi) \stackrel{\text{df}}{=} \{g \in G : \varphi(g) = e_H\}.$$

Definition 4.10. An abelian group is a pair $(G, \{\mu, \iota, e\})$, where

- $(G, \{\mu, \iota, e\})$ is a group,

that satisfies the following property:

$$(1) \text{ (Commutativity) } \forall x, y \in G : \quad \mu(x, y) = \mu(y, x).$$

Definition 4.11. A field is a pair $(\mathbb{F}, \{+_{\mathbb{F}}, -_{\mathbb{F}}, 0_{\mathbb{F}}, \times_{\mathbb{F}}, ^{-1}, 1_{\mathbb{F}}\})$, where

- $(\mathbb{F}, \{+_{\mathbb{F}}, -_{\mathbb{F}}, 0_{\mathbb{F}}\})$ is an abelian group,
- $\times_{\mathbb{F}} : \mathbb{F} \times \mathbb{F} \rightarrow \mathbb{F}$ is the field multiplication, and
- $(\mathbb{F}^{\times}, \{\times_{\mathbb{F}}|_{\mathbb{F}^{\times}}, ^{-1}, 1_{\mathbb{F}}\})$ is an abelian group,

that satisfy the following property:

$$(1) \text{ (Left Distributivity of Field Multiplication over Field Addition)}$$

$$\forall a, b, c \in \mathbb{F} : \quad a \times_{\mathbb{F}} (b +_{\mathbb{F}} c) = (a \times_{\mathbb{F}} b) +_{\mathbb{F}} (a \times_{\mathbb{F}} c).$$

Definition 4.12. An $\mathbb{F}$ -vector space is a pair $(V, \{+_V, -_V, \mathbf{0}_V, \cdot_V\})$, where

- $(V, \{+_V, -_V, \mathbf{0}_V\})$ is an abelian group, and
- $\cdot_V : \mathbb{F} \times V \rightarrow V$ is the scalar multiplication,

that satisfy the following four properties:

$$(1) \text{ (Left Distributivity of Scalar Multiplication over Vector Addition)}$$

$$\forall c \in \mathbb{F}, \forall \mathbf{u}, \mathbf{v} \in V : \quad c \cdot_V (\mathbf{u} +_V \mathbf{v}) = (c \cdot_V \mathbf{u}) +_V (c \cdot_V \mathbf{v}).$$

$$(2) \text{ (Right Distributivity of Scalar Multiplication over field Addition)}$$

$$\forall c, d \in \mathbb{F}, \forall \mathbf{u} \in V : \quad (c +_{\mathbb{F}} d) \cdot_V \mathbf{u} = (c \cdot_V \mathbf{u}) +_V (d \cdot_V \mathbf{u}).$$

$$(3) \text{ (Compatibility of Scalar Multiplication with Field Multiplication)}$$

$$\forall c, d \in \mathbb{F}, \forall \mathbf{u} \in V : \quad c \cdot_V (d \cdot_V \mathbf{u}) = (c \times_{\mathbb{F}} d) \cdot_V \mathbf{u}.$$

$$(4) \text{ (Existence of an Identity for Scalar Multiplication)}$$

$$\forall \mathbf{u} \in V : \quad 1_{\mathbb{F}} \cdot_V \mathbf{u} = \mathbf{u}.$$

Theorem 4.13 (Subspace Test). *Let $(V, \{+, -, \mathbf{0}, \cdot\})$ be an $\mathbb{F}$ -vector space. Let $W \subseteq V$. Then $(W, \{+, -, \mathbf{0}, \cdot\})$ is an $\mathbb{F}$ -vector space iff the following two statements hold:*

$$(1) \text{ (Non-Emptiness) } \mathbf{0} \in W.$$

$$(2) \text{ (Closure under Vector Addition and Scalar Multiplication)}$$

$$\forall \mathbf{u}, \mathbf{v} \in W, \forall \lambda \in \mathbb{F} : \quad \mathbf{u} + \lambda \cdot \mathbf{v} \in W.$$

Theorem 4.14 (Maps between a Set and a Vector Space Form a Vector Space). *Let $(V, \{+, -, \mathbf{0}, \cdot\})$ be an $\mathbb{F}$ -vector space. Let X be a set. Define the operations*

- $\dot{+} : V^X \times V^X \rightarrow V^X$, called “pointwise addition”,
- $\dot{-} : V^X \rightarrow V^X$, called “pointwise additive inversion”,
- $0_{X \rightarrow V} \in V^X$, called “zero map”, and
- $\dot{\cdot} : \mathbb{F} \times V^X \rightarrow V^X$, called “pointwise scalar multiplication”,

as follows:

$$(1) \text{ (Pointwise Addition) } \forall f, g \in V^X, \forall x \in X : \quad [f \dot{+} g](x) \stackrel{\text{df}}{=} f(x) + g(x).$$

$$(2) \text{ (Pointwise Additive Inversion) } \forall f \in V^X, \forall x \in X : \quad [\dot{-} f](x) \stackrel{\text{df}}{=} -f(x).$$

(3) (Zero Transformation) The map $\mathbf{0}_{X \rightarrow V} : X \rightarrow V$ is defined as:

$$\forall x \in X : \quad \mathbf{0}_{X \rightarrow V}(x) \stackrel{\text{df}}{=} \mathbf{0}.$$

(4) (Pointwise Scalar Multiplication) $\forall f \in V^X, \forall \lambda \in \mathbb{F}, \forall x \in X \quad [\lambda \cdot f](x) \stackrel{\text{df}}{=} \lambda \cdot f(x).$

Then $(V^X, \{\dot{+}, \dot{-}, \mathbf{0}_{X \rightarrow V}, \dot{\cdot}\})$ is an $\mathbb{F}$ -vector space.

Definition 4.15. Let $(V, \{+_V, -_V, \mathbf{0}_V, \cdot_V\})$ and $(W, \{+_W, -_W, \mathbf{0}_W, \cdot_W\})$ be $\mathbb{F}$ -vector spaces. A map $T : V \rightarrow W$ is said to be a linear transformation from V to W iff it satisfies the following property:

(1) (Linearity) $\forall \mathbf{u}, \mathbf{v} \in V, \forall \lambda \in \mathbb{F} : \quad T(\mathbf{u} +_V (\lambda \cdot_V \mathbf{v})) = T(\mathbf{u}) +_W (\lambda \cdot_W T(\mathbf{v})).$

A linear transformation from V to itself is called an operator on V . The set of linear transformations from V to W , denoted by " $\mathcal{L}(V, W)$ ", is defined as:

$$\mathcal{L}(V, W) \stackrel{\text{df}}{=} \{T : V \rightarrow W \quad | \quad T \text{ is a linear transformation}\}.$$

In the case where $V = W$, we simply denote $\mathcal{L}(V, W) = \mathcal{L}(V, V)$ by $\mathcal{L}(V)$.

Definition 4.16. Let $(V, \{+_V, -_V, \mathbf{0}_V, \cdot_V\})$ and $(W, \{+_W, -_W, \mathbf{0}_W, \cdot_W\})$ be $\mathbb{F}$ -vector spaces. Let $T \in \mathcal{L}(V, W)$. The null space of T , denoted by " $\text{Null}(T)$ ", is defined as the following:

$$\text{Null}(T) \stackrel{\text{df}}{=} \{\mathbf{v} \in V \quad | \quad T(\mathbf{v}) = \mathbf{0}_W\}.$$

The range of T , denoted by " $\text{Range}(T)$ ", is defined as the following:

$$\text{Range}(T) \stackrel{\text{df}}{=} \{T(\mathbf{v}) \quad | \quad \mathbf{v} \in V\}.$$

Definition 4.17. Let $(V, \{+_V, -_V, \mathbf{0}_V, \cdot_V\})$ and $(W, \{+_W, -_W, \mathbf{0}_W, \cdot_W\})$ be $\mathbb{F}$ -vector spaces. A map $T \in \mathcal{L}(V, W)$ is said to be a linear isomorphism iff it is bijective. (The inverse of a bijective linear map is automatically linear.)

Definition 4.18. $(V, \{+_V, -_V, \mathbf{0}_V, \cdot_V\})$ is said to be isomorphic to $(W, \{+_W, -_W, \mathbf{0}_W, \cdot_W\})$, denoted by " $V \cong W$ ", iff there exists a linear isomorphism between them.

Theorem 4.19 (linear transformations between vector spaces form a vector space). *Let $(V, \{+_V, -_V, \mathbf{0}_V, \cdot_V\})$ and $(W, \{+_W, -_W, \mathbf{0}_W, \cdot_W\})$ be $\mathbb{F}$ -vector spaces. Then $(\mathcal{L}(V, W), \{\dot{+}, \dot{-}, \mathbf{0}_{V \rightarrow W}, \dot{\cdot}\})$, is an $\mathbb{F}$ -vector space.*

Definition 4.20. An $\mathbb{F}$ -algebra is a pair $(A, \{+, -, \mathbf{0}, \cdot, \times\})$, where

- $(A, \{+, -, \mathbf{0}, \cdot\})$ is an $\mathbb{F}$ -vector space, and
- $\times : A \times A \rightarrow A$ is the algebra multiplication,

that satisfy the following property:

(1) (Bilinearity of the Algebra Multiplication) $\forall a, b, c \in A, \forall \lambda \in \mathbb{F} :$

$$a \times (b + \lambda \cdot c) = (a \times b) + \lambda \cdot (a \times c) \text{ and } (a + \lambda \cdot b) \times c = (a \times c) + \lambda \cdot (b \times c).$$

Definition 4.21. An $\mathbb{F}$ -algebra $(A, \{+, -, \mathbf{0}, \cdot, \times\})$ is said to be:

- (1) associative iff $\forall a, b, c \in A : \quad (a \times b) \times c = a \times (b \times c).$
- (2) commutative iff $\forall a, b \in A : \quad a \times b = b \times a.$

Definition 4.22. An $\mathbb{F}$ -algebra $(A, \{+, -, \mathbf{0}, \cdot, \times\})$ is said to be unital iff there exists an element $1 \in A$ such that $\forall a \in A : 1 \times a = a = a \times 1$. In this case the tuple expands to a pair $(A, \{+, -, \mathbf{0}, \cdot, \times, 1\})$.

Theorem 4.23 (Subalgebra Test). *Let $(A, \{+, -, \mathbf{0}, \cdot, \times\})$ be an $\mathbb{F}$ -algebra. Let $B \subseteq A$. Then $(B, \{+, -, \mathbf{0}, \cdot, \times\})$ is an $\mathbb{F}$ -algebra iff the following three statements hold:*

- (1) (Non-Emptiness) $\mathbf{0} \in B$.
- (2) (Closure under Vector Addition and Scalar Multiplication) $\forall a, b \in B, \forall \lambda \in \mathbb{F} : a + \lambda \cdot b \in B$.
- (3) (Closure under Algebra Multiplication) $\forall a, b \in B : a \times b \in B$.

Remark. Since associativity and commutativity are universally quantified equations, they are inherited by subsets automatically. Consequently, if $B \subseteq A$ satisfies the subalgebra test:

- if A is associative then B is associative.
- if A is commutative then B is commutative.
- if A is unital then B is unital iff $1 \in B$.

Theorem 4.24 (Maps between a Set and an Algebra Form an Algebra). *Let $(A, \{+, -, \mathbf{0}, \cdot, \times, 1\})$ be an associative unital $\mathbb{F}$ -algebra. Let X be a set. Define the pointwise algebra multiplication $\dot{\times} : A^X \times A^X \rightarrow A^X$ and unit $\dot{1} \in A^X$ by:*

$$\forall f, g \in A^X, \forall x \in X : [f \dot{\times} g](x) \stackrel{\text{df}}{=} f(x) \times g(x) \text{ and } \dot{1}(x) \stackrel{\text{df}}{=} 1.$$

Then $(A^X, \{\dot{+}, \dot{-}, 0_{X \rightarrow A}, \dot{\cdot}, \dot{\times}, \dot{1}\})$ is an associative unital $\mathbb{F}$ -algebra.

Proposition 4.25 (Associative Unital Algebra on $\mathcal{L}(V)$). *Let $(V, \{+, -, \mathbf{0}, \cdot\})$ be a $\mathbb{F}$ -vector space. Then $(\mathcal{L}(V), \{\dot{+}, \dot{-}, 0_{V \rightarrow V}, \dot{\cdot}, \circ, I_V\})$ is an associative unital algebra.*

Definition 4.26. A Lie $\mathbb{F}$ -algebra is a pair $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]\})$, where

- $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]\})$ is an $\mathbb{F}$ -algebra,

that satisfy the following two properties:

- (1) (Skew-symmetry) $\forall X, Y \in \mathfrak{g} : [Y, X] = -[X, Y]$.
- (2) (Jacobi Identity) $\forall X, Y, Z \in \mathfrak{g} : [X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = \mathbf{0}$.

Theorem 4.27 (Lie Subalgebra Test). *Let $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]\})$ be a Lie $\mathbb{F}$ -algebra. Let $\mathfrak{h} \subseteq \mathfrak{g}$. Then $(\mathfrak{h}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]\})$ is a Lie $\mathbb{F}$ -algebra iff the following three statements hold:*

- (1) (Non-Emptiness) $\mathbf{0} \in \mathfrak{h}$.
- (2) (Closure under Vector Addition and Scalar Multiplication) $\forall X, Y \in \mathfrak{h}, \forall \lambda \in \mathbb{F} : X + \lambda \cdot Y \in \mathfrak{h}$.
- (3) (Closure under the Lie Bracket) $\forall X, Y \in \mathfrak{h} : [X, Y] \in \mathfrak{h}$.

Theorem 4.28 (Lie Algebra Associated to an Associative Algebra). *Let $(A, \{+, -, \mathbf{0}, \cdot, \times\})$ be an associative $\mathbb{F}$ -algebra. Define the commutator $[\bullet, \bullet]_{\times} : A \times A \rightarrow A$ by*

$$\forall a, b \in A : [a, b]_{\times} \stackrel{\text{df}}{=} a \times b - b \times a.$$

Then $(A, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\times}\})$ is a Lie $\mathbb{F}$ -algebra.

Proof. $(A, \{+, -, \mathbf{0}, \cdot\})$ is a vector space (since A is an associative algebra). Also, $[\bullet, \bullet]_{\times} : A \times A \rightarrow A$ has the correct type. We just have to check the three properties:

(1) (Bilinearity) Let $\lambda \in \mathbb{F}$. Let $a, b, c \in A$. We have the following:

$$\begin{aligned}
 [a + \lambda \cdot b, c]_{\times} &= (a + \lambda \cdot b) \times c - c \times (a + \lambda \cdot b) \\
 &= a \times c + (\lambda \cdot b) \times c - c \times a - c \times (\lambda \cdot b) \\
 &= a \times c + \lambda \cdot (b \times c) - c \times a - \lambda \cdot (c \times b) \\
 &= a \times c - c \times a + \lambda \cdot (b \times c - c \times b) \\
 &= [a, c]_{\times} + \lambda \cdot [b, c]_{\times}; \\
 [a, b + \lambda \cdot c]_{\times} &= a \times (b + \lambda \cdot c) - (b + \lambda \cdot c) \times a \\
 &= a \times b + a \times (\lambda \cdot c) - b \times a - (\lambda \cdot c) \times a \\
 &= a \times b + \lambda \cdot (a \times c) - b \times a - \lambda \cdot (c \times a) \\
 &= a \times b - b \times a + \lambda \cdot (a \times c - c \times a) \\
 &= [a, b]_{\times} + \lambda \cdot [a, c]_{\times}.
 \end{aligned}$$

(2) (Skew-symmetry) Let $a, b \in A$. We have the following:

$$[a, b]_{\times} = b \times a - a \times b = -(a \times b - b \times a) = -[a, b]_{\times}.$$

(3) (Jacobi Identity) Let $a, b, c \in A$. We have the following:

$$\begin{aligned}
 [a, [b, c]_{\times}]_{\times} &= a \times [b, c]_{\times} - [b, c]_{\times} \times a \\
 &= a \times (b \times c - c \times b) - (b \times c - c \times b) \times a \\
 &= a \times b \times c - a \times c \times b - b \times c \times a + c \times b \times a.
 \end{aligned}$$

By the symmetry of the expressions, we also have:

$$\begin{aligned}
 [b, [c, a]_{\times}]_{\times} &= b \times c \times a - b \times a \times c - c \times a \times b + a \times c \times b; \\
 [c, [a, b]_{\times}]_{\times} &= c \times a \times b - c \times b \times a - a \times b \times c + b \times a \times c.
 \end{aligned}$$

Summing the three equations, we obtain:

$$[a, [b, c]_{\times}]_{\times} + [b, [c, a]_{\times}]_{\times} + [c, [a, b]_{\times}]_{\times} = \mathbf{0}.$$

□

Corollary 4.29 (Lie Subalgebra Criterion for Associative Algebras). *Let $(A, \{+, -, \mathbf{0}, \cdot, \times\})$ be an associative $\mathbb{F}$ -algebra. Let $B \subseteq A$. Then $(B, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\times}\})$ is a Lie $\mathbb{F}$ -algebra iff the following three statements hold:*

(1) (Non-Emptiness) $\mathbf{0} \in B$.

(2) (Closure under Vector Addition and Scalar Multiplication) $\forall a, b \in B, \forall \lambda \in \mathbb{F} : a + \lambda \cdot b \in B$.

(3) (Closure under the Commutator) $\forall a, b \in B : [a, b]_{\times} \in B$.

Definition 4.30. Let $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{g}}\})$ and $(\mathfrak{h}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{h}}\})$ be Lie $\mathbb{F}$ -algebras. A map $\varphi : \mathfrak{g} \rightarrow \mathfrak{h}$ is said to be a Lie algebra homomorphism between $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{g}}\})$ and $(\mathfrak{h}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{h}}\})$ iff it satisfies the following two properties:

(1) (Linearity) φ is a linear map.

(2) (Preservation of the Bracket) $\forall X, Y \in \mathfrak{g} : \varphi([X, Y]_{\mathfrak{g}}) = [\varphi(X), \varphi(Y)]_{\mathfrak{h}}$.

Definition 4.31. A Lie algebra homomorphism is called a Lie algebra isomorphism iff it is bijective. (The inverse of a bijective Lie algebra homomorphism is automatically a Lie algebra homomorphism.)

Definition 4.32. $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{g}}\})$ is said to be isomorphic to $(\mathfrak{h}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{h}}\})$, denoted by “ $\mathfrak{g} \cong \mathfrak{h}$ ”, iff there exists a Lie algebra isomorphism between them.

Definition 4.33. Let V be an $\mathbb{F}$ -vector space. The general linear Lie algebra of V , denoted by “ $\mathfrak{gl}(V)$ ”, is defined as

$$\mathfrak{gl}(V) \stackrel{\text{df}}{=} (\mathcal{L}(V), \{\dot{+}, \dot{-}, \mathbf{0}_{V \rightarrow V}, \dot{\cdot}, [\bullet, \bullet]_{\circ}\}).$$

5. DIFFERENTIAL CALCULUS ON SMOOTH MANIFOLDS

5.1. Local Constructions.

Definition 5.1. The standard smooth structure on $\mathbb{R}$, denoted by “ $\mathcal{A}_{\mathbb{R}}$ ”, is the smooth structure on $(\mathbb{R}, \mathcal{T}_{\text{dist}_1})$ that contains the identity chart $(\mathbb{R}, \text{id}_{\mathbb{R}})$.

Definition 5.2. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. The set of local smooth representatives at p , denoted by “ $\text{Rep}_p(M, \mathcal{T}, \mathcal{A})$ ”, is defined as:

$$\text{Rep}_p(M, \mathcal{T}, \mathcal{A}) \stackrel{\text{df}}{=} \{(U, f) : p \in U \in \mathcal{T} \text{ and } f : U \rightarrow \mathbb{R} \text{ is smooth w.r.t. } \mathcal{A}|_U \text{ and } \mathcal{A}_{\mathbb{R}}\}.$$

Definition 5.3. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$.

Let $(U, f), (V, g) \in \text{Rep}_p(M, \mathcal{T}, \mathcal{A})$. (U, f) and (V, g) are said to be germ equivalent at p , denoted by “ $(U, f) \sim_p (V, g)$ ”, iff

$$\exists W \in \mathcal{T} : p \in W \subseteq U \cap V \text{ and } f|_W = g|_W.$$

Definition 5.4. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. Let $(U, f) \in \text{Rep}_p(M, \mathcal{T}, \mathcal{A})$. The (U, f) -germ at p , denoted by “ $[(U, f)]_p$ ”, is defined as:

$$[(U, f)]_p \stackrel{\text{df}}{=} \{(V, g) \in \text{Rep}_p(M, \mathcal{T}, \mathcal{A}) : (U, f) \sim_p (V, g)\}.$$

The stalk of smooth functions at p , denoted by “ $C_p^{\infty}(M, \mathcal{T}, \mathcal{A})$ ”, is defined as:

$$C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) \stackrel{\text{df}}{=} \text{Rep}_p(M, \mathcal{T}, \mathcal{A}) / \sim_p = \left\{ [(U, f)]_p : (U, f) \in \text{Rep}_p(M, \mathcal{T}, \mathcal{A}) \right\}.$$

Theorem 5.5 (Stalk is a Commutative Associative Unital Algebra). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. Define the operations*

- $+$: $C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) \times C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) \rightarrow C_p^{\infty}(M, \mathcal{T}, \mathcal{A})$, the germ addition,
- $-$: $C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) \rightarrow C_p^{\infty}(M, \mathcal{T}, \mathcal{A})$, the additive germ inversion,
- $[(M, 0_{M \rightarrow \mathbb{R}})]_p \in C_p^{\infty}(M, \mathcal{T}, \mathcal{A})$, the zero germ,
- $\cdot$: $\mathbb{R} \times C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) \rightarrow C_p^{\infty}(M, \mathcal{T}, \mathcal{A})$, the scalar germ multiplication,
- $\times$: $C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) \times C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) \rightarrow C_p^{\infty}(M, \mathcal{T}, \mathcal{A})$, the algebra germ multiplication,
- $[(M, 1_{M \rightarrow \mathbb{R}})]_p \in C_p^{\infty}(M, \mathcal{T}, \mathcal{A})$, the unit germ,

on germs (via representatives) as follows:

(1) (Germ Addition)

$$\forall [(U, f)]_p, [(V, g)]_p \in C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) : [(U, f)]_p + [(V, g)]_p \stackrel{\text{df}}{=} [(U \cap V, f|_{U \cap V} + g|_{U \cap V})]_p.$$

(2) (Additive Germ Inversion)

$$\forall [(U, f)]_p \in C_p^{\infty}(M, \mathcal{T}, \mathcal{A}) : -[(U, f)]_p \stackrel{\text{df}}{=} [(U, -f)]_p.$$

(3) (Zero Germ)

$$[(M, 0_{M \rightarrow \mathbb{R}})]_p, \quad \forall q \in M : 0_{M \rightarrow \mathbb{R}}(q) \stackrel{\text{df}}{=} 0.$$

(4) (Scalar Germ Multiplication)

$$\forall [(U, f)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A}), \forall \lambda \in \mathbb{R} : \quad \lambda \cdot [(U, f)]_p \stackrel{\text{df}}{=} [(U, \lambda \cdot f)]_p.$$

(5) (Algebra Germ Multiplication)

$$\forall [(U, f)]_p, [(V, g)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A}) : \quad [(U, f)]_p \times [(V, g)]_p \stackrel{\text{df}}{=} [(U \cap V, (f \cdot g)|_{U \cap V})]_p.$$

(6) (Unit Germ)

$$[(M, 1_{M \rightarrow \mathbb{R}})]_p, \quad \forall q \in M : 1_{M \rightarrow \mathbb{R}}(q) \stackrel{\text{df}}{=} 1.$$

These operations are well-defined and

$$\left(C_p^\infty(M, \mathcal{T}, \mathcal{A}), \left\{ +, -, [(M, 0_{M \rightarrow \mathbb{R}})]_p, \cdot, \times, [(M, 1_{M \rightarrow \mathbb{R}})]_p \right\} \right)$$

is a commutative associative unital $\mathbb{R}$ -algebra.

Definition 5.6. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. A map $v : C_p^\infty(M, \mathcal{T}, \mathcal{A}) \rightarrow \mathbb{R}$ is said to be a derivation at p iff:

(1) ($\mathbb{R}$ -Linearity) $\forall \lambda \in \mathbb{R}, \forall [(U, f)]_p, [(V, g)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A}) :$

$$v\left([(U, f)]_p + \lambda \cdot [(V, g)]_p\right) = v\left([(U, f)]_p\right) + \lambda \cdot v\left([(V, g)]_p\right).$$

(2) (Leibniz Rule) $\forall [(U, f)]_p, [(V, g)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A}) :$

$$v\left([(U, f)]_p \times [(V, g)]_p\right) = v\left([(U, f)]_p\right) \cdot g(p) + f(p) \cdot v\left([(V, g)]_p\right).$$

The tangent space of $(M, \mathcal{T}, \mathcal{A})$ at p , denoted by “ $T_p(M, \mathcal{T}, \mathcal{A})$ ”, is defined as:

$$T_p(M, \mathcal{T}, \mathcal{A}) \stackrel{\text{df}}{=} \{v : C_p^\infty(M, \mathcal{T}, \mathcal{A}) \rightarrow \mathbb{R} : | v \text{ is a derivation at } p\}.$$

Theorem 5.7 (Tangent Space is a Vector Space). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. Then $(T_p(M, \mathcal{T}, \mathcal{A}), \{\dot{+}, \dot{-}, 0_{T_p(M)}, \dot{\cdot}\})$ is an $\mathbb{R}$ -vector space.*

Proof.

$$\mathbb{R}^{C_p^\infty(M, \mathcal{T}, \mathcal{A})}$$

is an $\mathbb{R}$ -vector space under pointwise operations (Theorem 4.14). $T_p(M, \mathcal{T}, \mathcal{A}) \subseteq \mathbb{R}^{C_p^\infty(M, \mathcal{T}, \mathcal{A})}$. Every derivation is linear, so $T_p(M, \mathcal{T}, \mathcal{A})$ is closed under $\dot{+}$ and $\dot{\cdot}$, and the zero map is a derivation. By the Subspace Test (Theorem 4.13),

$$(T_p(M, \mathcal{T}, \mathcal{A}), \{\dot{+}, \dot{-}, 0_{T_p(M)}, \dot{\cdot}\})$$

is an $\mathbb{R}$ -vector space. □

Theorem 5.8 (The Differential). *Let $(M, \mathcal{T}_M, \mathcal{A}_M)$ and $(N, \mathcal{T}_N, \mathcal{A}_N)$ be smooth manifolds. Let $F : M \rightarrow N$ be smooth w.r.t. $\mathcal{A}_M$ and $\mathcal{A}_N$. Let $p \in M$. The differential of F at p , denoted by “ dF_p ”, is the map $dF_p : T_p(M, \mathcal{T}_M, \mathcal{A}_M) \rightarrow T_{F(p)}(N, \mathcal{T}_N, \mathcal{A}_N)$ defined by:*

$\forall v \in T_p(M, \mathcal{T}_M, \mathcal{A}_M), \forall [(V, g)]_{F(p)} \in C_{F(p)}^\infty(N, \mathcal{T}_N, \mathcal{A}_N) :$

$$[dF_p(v)]\left([(V, g)]_{F(p)}\right) \stackrel{\text{df}}{=} v\left([(F^{-1}(V), g \circ F)]_p\right).$$

Then dF_p is a well-defined linear transformation.

Proof. (1) (Well-definedness — $dF_p(v)$ is a derivation at $F(p)$) Let $v \in T_p(M, \mathcal{T}_M, \mathcal{A}_M)$. We verify that $dF_p(v) : C_{F(p)}^\infty(N, \mathcal{T}_N, \mathcal{A}_N) \rightarrow \mathbb{R}$ satisfies the two defining properties of a derivation.

(a) ($\mathbb{R}$ -Linearity of $dF_p(v)$) Let $\lambda \in \mathbb{R}$ and $[(V, g)]_{F(p)}, [(W, h)]_{F(p)} \in C_{F(p)}^\infty(N, \mathcal{T}_N, \mathcal{A}_N)$. Then

$$\begin{aligned} [dF_p(v)]\left([(V, g)]_{F(p)} + \lambda \cdot [(W, h)]_{F(p)}\right) &= [dF_p(v)]\left([(V \cap W, g + \lambda \cdot h)]_{F(p)}\right) \\ &= v\left([(F^{-1}(V \cap W), (g + \lambda \cdot h) \circ F)]_p\right) \\ &= v\left([(F^{-1}(V) \cap F^{-1}(W), g \circ F + \lambda \cdot (h \circ F))]_p\right) \\ &= v\left([(F^{-1}(V), g \circ F)]_p + \lambda \cdot [(F^{-1}(W), h \circ F)]_p\right) \\ &= v\left([(F^{-1}(V), g \circ F)]_p\right) + \lambda \cdot v\left([(F^{-1}(W), h \circ F)]_p\right) \\ &= [dF_p(v)]\left([(V, g)]_{F(p)}\right) + \lambda \cdot [dF_p(v)]\left([(W, h)]_{F(p)}\right). \end{aligned}$$

(b) (Leibniz Rule of $dF_p(v)$) Let $[(V, g)]_{F(p)}, [(W, h)]_{F(p)} \in C_{F(p)}^\infty(N, \mathcal{T}_N, \mathcal{A}_N)$. Then

$$\begin{aligned} [dF_p(v)]\left([(V, g)]_{F(p)} \times [(W, h)]_{F(p)}\right) &= [dF_p(v)]\left([(V \cap W, g \cdot h)]_{F(p)}\right) \\ &= v\left([(F^{-1}(V \cap W), (g \cdot h) \circ F)]_p\right) \\ &= v\left([(F^{-1}(V) \cap F^{-1}(W), (g \circ F) \cdot (h \circ F))]_p\right) \\ &= v\left([(F^{-1}(V), g \circ F)]_p \times [(F^{-1}(W), h \circ F)]_p\right) \\ &= v\left([(F^{-1}(V), g \circ F)]_p\right) \cdot (h \circ F)(p) + (g \circ F)(p) \cdot v\left([(F^{-1}(W), h \circ F)]_p\right) \\ &= [dF_p(v)]\left([(V, g)]_{F(p)}\right) \cdot h(F(p)) + g(F(p)) \cdot [dF_p(v)]\left([(W, h)]_{F(p)}\right). \end{aligned}$$

Thus $dF_p(v) \in T_{F(p)}(N, \mathcal{T}_N, \mathcal{A}_N)$, so the codomain is correct.

(2) (Linearity of dF_p) Let $v, w \in T_p(M, \mathcal{T}_M, \mathcal{A}_M)$ and $\lambda \in \mathbb{R}$. For any germ $[(V, g)]_{F(p)} \in C_{F(p)}^\infty(N, \mathcal{T}_N, \mathcal{A}_N)$,

$$\begin{aligned} [dF_p(v + \lambda : w)]\left([(V, g)]_{F(p)}\right) &= (v + \lambda : w)\left([(F^{-1}(V), g \circ F)]_p\right) \\ &= v\left([(F^{-1}(V), g \circ F)]_p\right) + \lambda \cdot w\left([(F^{-1}(V), g \circ F)]_p\right) \\ &= [dF_p(v)]\left([(V, g)]_{F(p)}\right) + \lambda \cdot [dF_p(w)]\left([(V, g)]_{F(p)}\right) \\ &= [dF_p(v) + \lambda \cdot dF_p(w)]\left([(V, g)]_{F(p)}\right). \end{aligned}$$

Since the germ was arbitrary, $dF_p(v + \lambda : w) = dF_p(v) + \lambda : dF_p(w)$, establishing linearity. $\square$

Theorem 5.9 (Chain Rule). *Let $(M, \mathcal{T}_M, \mathcal{A}_M)$, $(N, \mathcal{T}_N, \mathcal{A}_N)$ and $(P, \mathcal{T}_P, \mathcal{A}_P)$ be smooth manifolds. Let $U \in \mathcal{T}_M$ and $V \in \mathcal{T}_N$. If*

- $F : U \rightarrow N$ is smooth w.r.t. $\mathcal{A}_M|_U$ and $\mathcal{A}_N$,
- $G : V \rightarrow P$ is smooth w.r.t. $\mathcal{A}_N|_V$ and $\mathcal{A}_P$, and
- $F(U) \subseteq V$,

then $G \circ F : U \rightarrow P$ is smooth w.r.t. $\mathcal{A}_M|_U$ and $\mathcal{A}_P$, and, for each $p \in U$,

$$d(G \circ F)_p = dG_{F(p)} \circ dF_p.$$

Proof. (1) (Smoothness of $G \circ F$) Let $p \in U$. Let $(W, \chi) \in \mathcal{A}_P$ with $[G \circ F](p) \in W$. By assumption $F(U) \subseteq V$, so $F(p) \in V$. Also, $[G \circ F](p) = G(F(p))$. Since $F(p) \in V$ and $(W, \chi) \in \mathcal{A}_P$ with $G(F(p)) \in W$, by the smoothness of G , we can find $(V_0, \psi) \in \mathcal{A}_N|_V$ with $F(p) \in V_0$ and $G(V_0) \subseteq W$, such that $\chi \circ G \circ \psi^{-1} : \psi(V_0) \rightarrow \chi(W)$ is C^∞ . Since $p \in U$ and $(V_0, \psi) \in \mathcal{A}_N|_V$ with $F(p) \in V_0$, by the smoothness of F , we can find $(U_0, \varphi) \in \mathcal{A}_M|_U$ with $p \in U_0$ and $F(U_0) \subseteq V_0$, such that $\psi \circ F \circ \varphi^{-1} : \varphi(U_0) \rightarrow \psi(V_0)$ is C^∞ . Notice the following:

$$\chi \circ (G \circ F) \circ \varphi^{-1} = \underbrace{(\chi \circ G \circ \psi^{-1})}_{C^\infty} \circ \underbrace{(\psi \circ F \circ \varphi^{-1})}_{C^\infty}.$$

Thus, we have found a $(U_0, \varphi) \in \mathcal{A}_M|_U$ with $p \in U_0$ and $[G \circ F](U_0) = G(F(U_0)) \subseteq G(V_0) \subseteq W$, such that $\chi \circ (G \circ F) \circ \varphi^{-1} : \varphi(U_0) \rightarrow \chi(W)$ is C^∞ .

(2) (the Formula) Let $p \in U$. Let $v \in T_p(M, \mathcal{T}_M, \mathcal{A}_M)$. Let $[(W, h)]_{[G \circ F](p)} \in C_{[G \circ F](p)}^\infty(P, \mathcal{T}_P, \mathcal{A}_P)$. From RHS to LHS:

$$\begin{aligned} [(dG_{F(p)} \circ dF_p)(v)] \left([(W, h)]_{[G \circ F](p)} \right) &= [dG_{F(p)}(dF_p(v))] \left([(W, h)]_{G(F(p))} \right) \\ &= [dF_p(v)] \left([(G^{-1}(W), h \circ G)]_{F(p)} \right) \\ &= v \left([(F^{-1}(G^{-1}(W)), (h \circ G) \circ F)]_p \right) \\ &= v \left(\left[\left[(G \circ F)^{-1} \right] (W), h \circ (G \circ F) \right]_p \right) \\ &= [d(G \circ F)_p(v)] \left([(W, h)]_{[G \circ F](p)} \right) \end{aligned}$$

□

5.2. Global Constructions.

Definition 5.10. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. The set of smooth functions on $(M, \mathcal{T}, \mathcal{A})$, denoted by " $C^\infty(M, \mathcal{T}, \mathcal{A})$ ", is defined as:

$$C^\infty(M, \mathcal{T}, \mathcal{A}) \stackrel{\text{df}}{=} \{f : M \rightarrow \mathbb{R} \mid f \text{ is smooth w.r.t. } \mathcal{A} \text{ and } \mathcal{A}_\mathbb{R}\}.$$

Theorem 5.11 ($C^\infty(M)$ is a Commutative Associative Unital Algebra). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. The set $C^\infty(M, \mathcal{T}, \mathcal{A})$ carries the pointwise operations (Definition 4.14 with $X = M$ and $A = \mathbb{R}$). Then $(C^\infty(M, \mathcal{T}, \mathcal{A}), \{+, -, 0_{M \rightarrow \mathbb{R}}, \cdot, \times, 1_{M \rightarrow \mathbb{R}}\})$ is a commutative associative unital $\mathbb{R}$ -algebra. Consequently, $(\mathcal{L}(C^\infty(M, \mathcal{T}, \mathcal{A})), \{\dot{+}, \dot{-}, \mathbf{0}, \dot{\cdot}, \circ, I\})$ is an associative unital $\mathbb{R}$ -algebra.*

Definition 5.12. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. A map $X \in \mathcal{L}(C^\infty(M, \mathcal{T}, \mathcal{A}))$ is said to be a smooth vector field iff it satisfies the following property:

(1) (Leibniz Rule) $\forall f, g \in C^\infty(M, \mathcal{T}, \mathcal{A}) : X(f \times g) = X(f) \times g + f \times X(g)$.

The set of smooth vector fields, denoted by " $\mathfrak{X}(M, \mathcal{T}, \mathcal{A})$ ", is defined as:

$$\mathfrak{X}(M, \mathcal{T}, \mathcal{A}) \stackrel{\text{df}}{=} \{X \in \mathcal{L}(C^\infty(M, \mathcal{T}, \mathcal{A})) \mid X \text{ is a smooth vector field}\}.$$

Lemma 5.13 ($\mathfrak{X}(M)$ is a Vector Space). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Then $\mathfrak{X}(M, \mathcal{T}, \mathcal{A}) \subseteq \mathcal{L}(C^\infty(M, \mathcal{T}, \mathcal{A}))$ and $(\mathfrak{X}(M, \mathcal{T}, \mathcal{A}), \{\dot{+}, \dot{-}, \mathbf{0}, \dot{\cdot}\})$ is an $\mathbb{R}$ -vector space.*

5.3. Local-to-Global Correspondences.

Lemma 5.14 (Germ Extension). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. Let $(U, f) \in \text{Rep}_p(M, \mathcal{T}, \mathcal{A})$. Then there exists $\tilde{f} \in C^\infty(M, \mathcal{T}, \mathcal{A})$ such that $(U, f) \sim_p (M, \tilde{f})$. Consequently, $[(U, f)]_p = [(M, \tilde{f})]_p$.*

Proof. Since $U \in \mathcal{T}$ and $p \in U$, we can choose an open neighborhood V of p with $\bar{V} \subseteq U$ (e.g., take a coordinate ball around p contained in U). Let $\rho : M \rightarrow [0, 1]$ be a smooth bump function with $\rho \equiv 1$ on V and $\text{supp}(\rho) \subseteq U$. Such a function exists on any smooth manifold.

Define $\tilde{f} : M \rightarrow \mathbb{R}$ by

$$\tilde{f}(q) \stackrel{\text{df}}{=} \begin{cases} \rho(q) \cdot f(q), & q \in U, \\ 0, & q \in M \setminus U. \end{cases}$$

On U , $\tilde{f} = \rho \cdot f$ is smooth because both ρ and f are smooth on U . On $M \setminus \text{supp}(\rho)$, we have $\rho \equiv 0$, so $\tilde{f} \equiv 0$, which is smooth. Since the two definitions agree on the overlap (where ρ transitions to 0), $\tilde{f}$ is smooth on all of M . Thus $\tilde{f} \in C^\infty(M, \mathcal{T}, \mathcal{A})$.

Finally, for any $q \in V$ we have $\rho(q) = 1$, so $\tilde{f}(q) = f(q)$. Hence V is an open neighborhood of p on which $\tilde{f}$ and f agree, i.e., $(U, f) \sim_p (M, \tilde{f})$. The equality of germs follows directly. $\square$

Lemma 5.15 (Locality of a Smooth Vector Field). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. Let $X \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$. Let $h \in C^\infty(M, \mathcal{T}, \mathcal{A})$. If*

$$\exists U \in \{U \in \mathcal{T} : p \in U\}, \forall q \in U : h(q) = 0,$$

then $[X(h)](p) = 0$.

Proof. Since $U \in \mathcal{T}$ and $p \in U$, we can find a smooth bump function $\rho \in C^\infty(M, \mathcal{T}, \mathcal{A})$ such that $\rho(p) = 0$ and $\rho \equiv 1$ on $M \setminus U$. (Such a function exists because on a smooth manifold one can construct a smooth function that vanishes at a point and equals 1 outside any prescribed open neighborhood of that point.)

Now observe that $h = \rho \dot{\times} h$ globally:

- If $q \in U$, then $h(q) = 0$ by hypothesis, so $[\rho \dot{\times} h](q) = \rho(q) \cdot 0 = 0 = h(q)$.
- If $q \notin U$, then $\rho(q) = 1$, so $[\rho \dot{\times} h](q) = 1 \cdot h(q) = h(q)$.

Thus $h = \rho \dot{\times} h$. Applying X to both sides and using the Leibniz rule for smooth vector fields, we obtain

$$X(h) = X(\rho \dot{\times} h) = X(\rho) \dot{\times} h + \rho \dot{\times} X(h).$$

Evaluating at p and using the pointwise definitions (Definition 4.14) gives

$$[X(h)](p) = [X(\rho)](p) \cdot h(p) + \rho(p) \cdot [X(h)](p).$$

Since $p \in U$, we have $h(p) = 0$ by hypothesis, and by construction $\rho(p) = 0$. Hence

$$[X(h)](p) = [X(\rho)](p) \cdot 0 + 0 \cdot [X(h)](p) = 0,$$

which completes the proof. $\square$

Definition 5.16. Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. Let $X \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$. The value of X at p , “denoted by $X|_p$ ”, is the map $X|_p : C_p^\infty(M, \mathcal{T}, \mathcal{A}) \rightarrow \mathbb{R}$ defined as the following:

$$\forall [(U, f)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A}) : X|_p \left([(U, f)]_p \right) \stackrel{\text{df}}{=} [X(\tilde{f})]_p(p).$$

Lemma 5.17 (Value of a Vector Field is a Derivation). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Let $p \in M$. Let $X \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$. Then $X|_p \in T_p(M, \mathcal{T}, \mathcal{A})$.*

Proof. $X|_p : C_p^\infty(M, \mathcal{T}, \mathcal{A}) \rightarrow \mathbb{R}$ already has the correct type. We just need to check that it is a derivation at p .

(1) (Linearity) Let $\lambda \in \mathbb{R}$. Let $[(U, f)]_p, [(V, g)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A})$. We have the following:

$$\begin{aligned}
 X|_p\left([(U, f)]_p + \lambda \cdot [(V, g)]_p\right) &= X|_p\left([(U, f)]_p + [(V, \lambda \cdot g)]_p\right) \\
 &= X|_p\left([(U \cap V, f + \lambda \cdot g)]_p\right) \\
 &= \left[X(\widetilde{f + \lambda \cdot g})\right](p) \\
 &= \left[X(\tilde{f} + \lambda \cdot \tilde{g})\right](p) \\
 &= \left[X(\tilde{f}) + \lambda \cdot X(\tilde{g})\right](p) \\
 &= \left[X(\tilde{f}) + (\lambda \cdot X)(\tilde{g})\right](p) \\
 &= \left[X(\tilde{f})\right](p) + [(\lambda \cdot X)(\tilde{g})](p) \\
 &= \left[X(\tilde{f})\right](p) + [\lambda \cdot X(\tilde{g})](p) \\
 &= \left[X(\tilde{f})\right](p) + \lambda \cdot [X(\tilde{g})](p) \\
 &= X|_p\left([(U, f)]_p\right) + \lambda \cdot X|_p\left([(V, g)]_p\right)
 \end{aligned}$$

(2) (Leibniz Rule) Let $[(U, f)]_p, [(V, g)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A})$. We have the following:

$$\begin{aligned}
 X|_p\left([(U, f)]_p \times [(V, g)]_p\right) &= X|_p\left([(U \cap V, (f \cdot g)|_{U \cap V})]_p\right) \\
 &= \left[X(\widetilde{(f \cdot g)|_{U \cap V}})\right](p) \\
 &= \left[X(\tilde{f} \times \tilde{g})\right](p) \\
 &= \left[X(\tilde{f}) \times \tilde{g} + \tilde{f} \times X(\tilde{g})\right](p) \\
 &= \left[X(\tilde{f}) \times \tilde{g}\right](p) + \left[\tilde{f} \times X(\tilde{g})\right](p) \\
 &= \left[X(\tilde{f})\right](p) \cdot \tilde{g}(p) + \tilde{f}(p) \cdot [X(\tilde{g})](p) \\
 &= X|_p\left([(U, f)]_p\right) \cdot g(p) + f(p) \cdot X|_p\left([(V, g)]_p\right).
 \end{aligned}$$

□

Definition 5.18 (General Evaluation Map). Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold and $p \in M$. The evaluation map at p is the map

$$\text{ev}_p : \mathfrak{X}(M, \mathcal{T}, \mathcal{A}) \rightarrow T_p(M, \mathcal{T}, \mathcal{A}), \quad \text{ev}_p(X) \stackrel{\text{df}}{=} X|_p,$$

where $X|_p$ is the value of the smooth vector field X at p . Lemma 5.17 guarantees that $X|_p \in T_p(M, \mathcal{T}, \mathcal{A})$, so the codomain is correct.

Lemma 5.19 (ev_p is Linear). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold and $p \in M$. Then the evaluation map $\text{ev}_p : \mathfrak{X}(M, \mathcal{T}, \mathcal{A}) \rightarrow T_p(M, \mathcal{T}, \mathcal{A})$ is a linear transformation.*

Proof. Let $X, Y \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$ and $\lambda \in \mathbb{R}$. For any germ $[(U, f)]_p \in C_p^\infty(M, \mathcal{T}, \mathcal{A})$, let $\tilde{f} \in C^\infty(M, \mathcal{T}, \mathcal{A})$ be a global representative (Lemma 5.14). Then

$$\begin{aligned} \text{ev}_p(X \dot{+} \lambda : Y) \left([(U, f)]_p \right) &= \left[(X \dot{+} \lambda : Y) \left(\tilde{f} \right) \right] (p) && \text{(definition of } X|_p) \\ &= \left[X \left(\tilde{f} \right) + \lambda \cdot Y \left(\tilde{f} \right) \right] (p) && \text{(pointwise operations)} \\ &= \left[X \left(\tilde{f} \right) \right] (p) + \lambda \cdot \left[Y \left(\tilde{f} \right) \right] (p) \\ &= \text{ev}_p(X) \left([(U, f)]_p \right) + \lambda \cdot \text{ev}_p(Y) \left([(U, f)]_p \right). \end{aligned}$$

Since the germ was arbitrary, $\text{ev}_p(X \dot{+} \lambda : Y) = \text{ev}_p(X) \dot{+} \lambda : \text{ev}_p(Y)$, establishing linearity. $\square$

Theorem 5.20 (Vector Fields form a Lie Algebra). *Let $(M, \mathcal{T}, \mathcal{A})$ be a smooth manifold. Define the commutator of $X, Y \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$ by $[X, Y]_\circ \stackrel{\text{df}}{=} X \circ Y - Y \circ X$, i.e.,*

$$\forall f \in C^\infty(M, \mathcal{T}, \mathcal{A}) : ([X, Y]_\circ)(f) = X(Y(f)) - Y(X(f)).$$

Then $[X, Y]_\circ \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$ and $(\mathfrak{X}(M, \mathcal{T}, \mathcal{A}), \{+, -, 0_{\mathfrak{X}(M)}, \cdot, [\bullet, \bullet]_\circ\})$ is a Lie $\mathbb{R}$ -algebra.

Proof. First, we verify that $[X, Y]_\circ$ is a smooth vector field.

$$[X, Y]_\circ : C^\infty(M, \mathcal{T}, \mathcal{A}) \rightarrow C^\infty(M, \mathcal{T}, \mathcal{A})$$

already has the correct type. We check the Leibniz Rule:

(1) (Linearity) Let $\lambda \in \mathbb{R}$. Let $f, g \in C^\infty(M, \mathcal{T}, \mathcal{A})$.

$$\begin{aligned} [X, Y]_\circ(f + \lambda \cdot g) &= X(Y(f + \lambda \cdot g)) - Y(X(f + \lambda \cdot g)) \\ &= X(Y(f) + \lambda \cdot Y(g)) - Y(X(f) + \lambda \cdot X(g)) \\ &= X(Y(f)) + \lambda \cdot X(Y(g)) - (Y(X(f)) + \lambda \cdot Y(X(g))) \\ &= X(Y(f)) + \lambda \cdot X(Y(g)) - Y(X(f)) - \lambda \cdot Y(X(g)) \\ &= X(Y(f)) - Y(X(f)) + \lambda \cdot X(Y(g)) - \lambda \cdot Y(X(g)) \\ &= X(Y(f)) - Y(X(f)) + \lambda \cdot (X(Y(g)) - Y(X(g))) \\ &= [X, Y]_\circ(f) + \lambda \cdot [X, Y]_\circ(g). \end{aligned}$$

(2) (Leibniz Rule) Let $f, g \in C^\infty(M, \mathcal{T}, \mathcal{A})$.

$$\begin{aligned} [X, Y]_\circ(f \times g) &= X(Y(f \times g)) - Y(X(f \times g)) \\ &= X(Y(f) \times g + f \times Y(g)) - Y(X(f) \times g + f \times X(g)) \\ &= X(Y(f) \times g) + X(f \times Y(g)) - Y(X(f) \times g) - Y(f \times X(g)) \\ &= X(Y(f)) \times g + Y(f) \times X(g) + X(f) \times Y(g) + f \times X(Y(g)) \\ &\quad - Y(X(f)) \times g - X(f) \times Y(g) - Y(f) \times X(g) - f \times Y(X(g)) \\ &= X(Y(f)) \times g - Y(X(f)) \times g + f \times X(Y(g)) - f \times Y(X(g)) \\ &= (X(Y(f)) - Y(X(f))) \times g + f \times (X(Y(g)) - Y(X(g))) \\ &= [X, Y]_\circ(f) \times g + f \times [X, Y]_\circ(g). \end{aligned}$$

Thus $[X, Y]_\circ \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$.

Now, $(\mathcal{L}(C^\infty(M, \mathcal{T}, \mathcal{A})), \{\dot{+}, \dot{-}, 0, \dot{\cdot}, \circ\})$ is an associative algebra, $\mathfrak{X}(M, \mathcal{T}, \mathcal{A}) \subseteq \mathcal{L}(C^\infty(M, \mathcal{T}, \mathcal{A}))$, $(\mathfrak{X}(M, \mathcal{T}, \mathcal{A}), \{\dot{+}, \dot{-}, 0, \dot{\cdot}\})$ is a vector space, and $\forall X, Y \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A}) : [X, Y]_\circ \in \mathfrak{X}(M, \mathcal{T}, \mathcal{A})$. By 4.29, $(\mathfrak{X}(M, \mathcal{T}, \mathcal{A}), \{\dot{+}, \dot{-}, 0, \dot{\cdot}, [\bullet, \bullet]_\circ\})$ is a Lie $\mathbb{R}$ -algebra. $\square$

6. LIE GROUPS AND LIE ALGEBRAS

Remark. Replacing “smooth manifold” with “topological space” and “smooth” with “continuous” in the definition below yields the notion of a topological group.

Definition 6.1. A Lie group is a 4-tuple $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$, where

- $(G, \mathcal{T}, \mathcal{A})$ is a smooth manifold, and
- $(G, \{\mu, \iota, e\})$ is a group,

that satisfy the following two properties:

- (1) (Smoothness of the Group Operation)
 $\mu : G \times G \rightarrow G$ is smooth w.r.t. $\mathcal{A} \otimes \mathcal{A}$ and $\mathcal{A}$.
- (2) (Smoothness of the Inversion Map) $\iota : G \rightarrow G$ is smooth w.r.t. $\mathcal{A}$ and $\mathcal{A}$.

We would adopt the common notation $\forall g, h \in G : gh \stackrel{\text{df}}{=} \mu(g, h)$ and $g^{-1} \stackrel{\text{df}}{=} \iota(g)$.

Lemma 6.2 (Left Inclusion is C^∞). *Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. The left inclusion map $i_g : G \rightarrow G \times G$ defined by*

$$\forall h \in G : i_g(h) = (g, h)$$

is smooth w.r.t. $\mathcal{A}$ and $\mathcal{A} \otimes \mathcal{A}$.

Proof. We would use the simpler smoothness criterion (Lemma 3.11). Let $h \in G$. Choose $(U, \varphi) \in \mathcal{A}$ with $g \in U$ and $(V, \psi) \in \mathcal{A}$ with $h \in V$. Consider $(U \times V, \varphi \times \psi) \in \mathcal{A} \otimes \mathcal{A}$. We have

$$i_g(V) = \{g\} \times V \subseteq U \times V.$$

Moreover,

$$(\varphi \times \psi) \circ i_g \circ \psi^{-1} = \left(\underbrace{c_{\varphi(g)}}_{C^\infty}, \underbrace{\text{id}_{\mathbb{R}}}_{C^\infty} \right).$$

□

Definition 6.3. Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. Let $g \in G$. The left translation at g , denoted by “ L_g ”, is the map $L_g : G \rightarrow G$ defined as the following:

$$\forall h \in G : L_g(h) \stackrel{\text{df}}{=} [\mu \circ i_g](h) = gh.$$

Theorem 6.4 (Left Translation is a Diffeomorphism). *Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. Let $g \in G$. Then $L_g : G \rightarrow G$ is a diffeomorphism between $(G, \mathcal{T}, \mathcal{A})$ and itself.*

Proof. (1) (Invertibility) The map $L_{g^{-1}} : G \rightarrow G$ defined by

$$\forall h \in G : L_{g^{-1}}(h) = g^{-1}h$$

is an inverse for L_g . For any $h \in G$, we have

$$\begin{aligned} [L_{g^{-1}} \circ L_g](h) &= L_{g^{-1}}(L_g(h)) = L_{g^{-1}}(gh) = g^{-1}(gh) = (g^{-1}g)h = eh = h; \\ [L_g \circ L_{g^{-1}}](h) &= L_g(L_{g^{-1}}(h)) = L_g(g^{-1}h) = g(g^{-1}h) = (gg^{-1})h = eh = h. \end{aligned}$$

- (2) (Smoothness of L_g) Since we have a Lie group, $\mu : G \times G \rightarrow G$ is smooth w.r.t. $\mathcal{A} \otimes \mathcal{A}$ and $\mathcal{A}$. Also, $i_g : G \rightarrow G \times G$ is smooth w.r.t. $\mathcal{A}$ and $\mathcal{A} \otimes \mathcal{A}$. By the argument used in the proof for the Chain Rule, the composition $\mu \circ i_g$ is smooth w.r.t. $\mathcal{A}$ and $\mathcal{A}$.

- (3) (Smoothness of $L_{g^{-1}}$) The same argument as (2), with g replaced by g^{-1} .

□

Definition 6.5. Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. Let $X \in \mathfrak{X}(G, \mathcal{T}, \mathcal{A})$. The smooth vector field $X : C^\infty(G, \mathcal{T}, \mathcal{A}) \rightarrow C^\infty(G, \mathcal{T}, \mathcal{A})$ is said to be μ -left-invariant iff it satisfy the following property:

$$\forall g \in G, \forall f \in C^\infty(G, \mathcal{T}, \mathcal{A}) : \quad X(f \circ L_g) = X(f) \circ L_g.$$

Lemma 6.6 (Equivalent Characterization of Left-Invariance). *Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. Let $X \in \mathfrak{X}(G, \mathcal{T}, \mathcal{A})$. The smooth vector field $X : C^\infty(G, \mathcal{T}, \mathcal{A}) \rightarrow C^\infty(G, \mathcal{T}, \mathcal{A})$ is μ -left-invariant iff*

$$\forall g, h \in G, \forall [(U, f)]_{gh} \in C_{gh}^\infty(G, \mathcal{T}, \mathcal{A}) : \quad X|_{gh} \left([(U, f)]_{gh} \right) = X|_h \left([(L_g^{-1}(U), f \circ L_g)]_h \right).$$

Proof. ($\implies$) Let $g, h \in G$. Let $[(U, f)]_{gh} \in C_{gh}^\infty(G, \mathcal{T}, \mathcal{A})$. Since $gh \in G$ and $(U, f) \in \text{Rep}_{gh}(G, \mathcal{T}, \mathcal{A})$, by the Germ Extension Lemma, we can find $\tilde{f} \in C^\infty(G, \mathcal{T}, \mathcal{A})$ such that $[(U, f)]_{gh} = \left[(G, \tilde{f}) \right]_{gh}$. On the LHS, we have

$$X|_{gh} \left([(U, f)]_{gh} \right) = X|_{gh} \left(\left[(G, \tilde{f}) \right]_{gh} \right) = \left[X(\tilde{f}) \right](gh).$$

Notice that

$$(U, f) \sim_{gh} (G, \tilde{f}) \implies (L_g^{-1}(U), f \circ L_g) \sim_h (L_g^{-1}(G), \tilde{f} \circ L_g) = (G, \tilde{f} \circ L_g).$$

On the RHS, we have

$$X|_h \left([(L_g^{-1}(U), f \circ L_g)]_h \right) = X|_h \left(\left[(G, \tilde{f} \circ L_g) \right]_h \right) = \left[X(\tilde{f} \circ L_g) \right](h).$$

By assumption, we have

$$\left[X(\tilde{f} \circ L_g) \right](h) = \left[X(\tilde{f}) \circ L_g \right](h) = \left[X(\tilde{f}) \right](L_g(h)) = \left[X(\tilde{f}) \right](gh).$$

($\impliedby$) Let $g \in G$. Let $f \in C^\infty(G, \mathcal{T}, \mathcal{A})$. Let $h \in G$. Since $gh \in G$ and $[(G, f)]_{gh} \in C^\infty(G, \mathcal{T}, \mathcal{A})$, by assumption, we have

$$\begin{aligned} X|_{gh} \left([(G, f)]_{gh} \right) &= X|_h \left([(L_g^{-1}(G), f \circ L_g)]_h \right) \\ X|_{gh} \left([(G, f)]_{gh} \right) &= X|_h \left([(G, f \circ L_g)]_h \right) \\ [X(f)](gh) &= [X(f \circ L_g)](h) \\ [X(f)](L_g(h)) &= [X(f \circ L_g)](h) \\ [X(f) \circ L_g](h) &= [X(f \circ L_g)](h) \end{aligned}$$

So, $X(f \circ L_g) = X(f) \circ L_g$, as desired. $\square$

Definition 6.7. Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. The set of left-invariant vector fields on G , denoted by " $\mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ ", is defined as:

$$\mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\}) \stackrel{\text{df}}{=} \{X \in \mathfrak{X}(G, \mathcal{T}, \mathcal{A}) : X \text{ is } \mu\text{-left-invariant}\}.$$

Theorem 6.8 (Left-Invariant Fields form a Lie Algebra). *Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. Then $(\mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\}), \dot{+}, \dot{-}, 0_{\mathfrak{X}(G)}, \dot{\cdot}, [\bullet, \bullet]_{\circ})$ is a Lie $\mathbb{R}$ -algebra.*

Proof. We apply the Lie Subalgebra Test (Theorem 4.27) to $\mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\}) \subseteq \mathfrak{X}(G, \mathcal{T}, \mathcal{A})$, where $\mathfrak{X}(G, \mathcal{T}, \mathcal{A})$ is a Lie $\mathbb{R}$ -algebra with the commutator bracket (Theorem 5.20). We verify the three conditions.

- (1) (Non-Emptiness) The zero smooth vector field $0_{\mathfrak{X}(G)} \in \mathfrak{X}(G, \mathcal{T}, \mathcal{A})$ is μ -left-invariant, because for any $g \in G$ and $f \in C^\infty(G, \mathcal{T}, \mathcal{A})$,

$$0_{\mathfrak{X}(G)}(f \circ L_g) = 0_{\mathfrak{X}(G)} = 0_{G \rightarrow \mathbb{R}} = 0_{G \rightarrow \mathbb{R}} \circ L_g = 0_{\mathfrak{X}(G)}(f) \circ L_g.$$

Hence $0_{\mathfrak{X}(G)} \in \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$.

- (2) (Closure under Vector Addition and Scalar Multiplication) Let $X, Y \in \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ and $\lambda \in \mathbb{R}$. For any $g \in G$ and $f \in C^\infty(G, \mathcal{T}, \mathcal{A})$,

$$\begin{aligned} [X + \lambda \cdot Y](f \circ L_g) &= X(f \circ L_g) + \lambda \cdot Y(f \circ L_g) && \text{(pointwise operations)} \\ &= X(f) \circ L_g + \lambda \cdot (Y(f) \circ L_g) && (X, Y \text{ are left-invariant}) \\ &= (X(f) + \lambda \cdot Y(f)) \circ L_g \\ &= [X + \lambda \cdot Y](f) \circ L_g. \end{aligned}$$

Thus $X + \lambda \cdot Y$ is μ -left-invariant.

- (3) (Closure under the Lie Bracket) Let $X, Y \in \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$. For any $g \in G$ and $f \in C^\infty(G, \mathcal{T}, \mathcal{A})$,

$$\begin{aligned} [X, Y]_\circ(f \circ L_g) &= X(Y(f \circ L_g)) - Y(X(f \circ L_g)) \\ &= X(Y(f) \circ L_g) - Y(X(f) \circ L_g) && (X \text{ and } Y \text{ are } \mu\text{-left-invariant}) \\ &= X(Y(f)) \circ L_g - Y(X(f)) \circ L_g \\ &= (X(Y(f)) - Y(X(f))) \circ L_g \\ &= [X, Y]_\circ(f) \circ L_g. \end{aligned}$$

Thus $[X, Y]_\circ$ is μ -left-invariant, i.e., $[X, Y]_\circ \in \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$.

All three conditions of Theorem 4.27 are satisfied, so $(\mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\}), \dot{+}, \dot{-}, 0_{\mathfrak{X}(G)}, \dot{\cdot}, [\bullet, \bullet]_\circ)$ is a Lie $\mathbb{R}$ -algebra. $\square$

Theorem 6.9 (Evaluation at the Identity is a Linear Isomorphism). *Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. The map*

$$\text{ev}_e : \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\}) \rightarrow T_e(G, \mathcal{T}, \mathcal{A}), \quad \text{ev}_e(X) \stackrel{\text{df}}{=} X|_e,$$

is a linear isomorphism.

Proof. (1) (Linearity) The evaluation map ev_p on the full space $\mathfrak{X}(M, \mathcal{T}, \mathcal{A})$ is linear (a consequence of the pointwise operations on $\mathcal{L}(C^\infty(M, \mathcal{T}, \mathcal{A}))$). Restricting to the subspace $\mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ preserves linearity.

- (2) (Injectivity) Let $X \in \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ and suppose $\text{ev}_e(X) = X|_e = 0$. Let $g \in G$ and let $[(U, f)]_g \in C_g^\infty(G)$ be an arbitrary germ at g . Using the equivalent characterisation of left-invariance (Lemma 6.6) with $h = e$, we obtain

$$X_g([(U, f)]_g) = X|_e([(L_g^{-1}(U), f \circ L_g)]_e) = 0([(L_g^{-1}(U), f \circ L_g)]_e) = 0.$$

Since the germ at g was arbitrary, $X_g = 0$ for every $g \in G$. Hence $X = 0_{\mathfrak{X}(G)}$, the zero smooth vector field. This proves ev_e is injective.

- (3) (Surjectivity) Let $v \in T_e(G, \mathcal{T}, \mathcal{A})$ be an arbitrary tangent vector at the identity. Define a map X^v on germs by

$$\forall g \in G, \forall [(U, f)]_g \in C_g^\infty(G) : \quad X_g^v([(U, f)]_g) \stackrel{\text{df}}{=} v([(L_g^{-1}(U), f \circ L_g)]_e).$$

We verify that X^v is a well-defined left-invariant smooth vector field and that $\text{ev}_e(X^v) = v$.

(a) (Well-definedness on germs) If $[(U, f)]_g = [(V, \tilde{f})]_g$, then f and $\tilde{f}$ agree on an open neighborhood W of g . Then $f \circ L_g$ and $\tilde{f} \circ L_g$ agree on $L_g^{-1}(W)$, which is an open neighborhood of e . Hence the germs at e coincide, and the value $v\left([(L_g^{-1}(U), f \circ L_g)]_e\right)$ depends only on the germ $[(U, f)]_g$.

(b) (X_g^v is a derivation at g) Let $\lambda \in \mathbb{R}$ and $[(U, f)]_g, [(V, \tilde{f})]_g \in C_g^\infty(G)$. For linearity:

$$\begin{aligned} X_g^v\left([(U, f)]_g + \lambda \cdot [(V, \tilde{f})]_g\right) &= v\left([(L_g^{-1}(U \cap V), (f + \lambda \cdot \tilde{f}) \circ L_g)]_e\right) \\ &= v\left([(L_g^{-1}(U), f \circ L_g)]_e + \lambda \cdot [(L_g^{-1}(V), \tilde{f} \circ L_g)]_e\right) \\ &= X_g^v\left([(U, f)]_g\right) + \lambda \cdot X_g^v\left([(V, \tilde{f})]_g\right). \end{aligned}$$

For the Leibniz rule:

$$\begin{aligned} X_g^v\left([(U, f)]_g \times [(V, \tilde{f})]_g\right) &= v\left([(L_g^{-1}(U \cap V), (f \cdot \tilde{f}) \circ L_g)]_e\right) \\ &= v\left([(L_g^{-1}(U), f \circ L_g)]_e \times [(L_g^{-1}(V), \tilde{f} \circ L_g)]_e\right) \\ &= v\left([(L_g^{-1}(U), f \circ L_g)]_e\right) \cdot \tilde{f}(g) + f(g) \cdot v\left([(L_g^{-1}(V), \tilde{f} \circ L_g)]_e\right) \\ &= X_g^v\left([(U, f)]_g\right) \cdot \tilde{f}(g) + f(g) \cdot X_g^v\left([(V, \tilde{f})]_g\right), \end{aligned}$$

where we used that $[(\tilde{f} \circ L_g)]_e = \tilde{f}(g)$ and $[(f \circ L_g)]_e = f(g)$.

(c) (Smoothness of X^v) Let $f \in C^\infty(G, \mathcal{T}, \mathcal{A})$. The map

$$g \mapsto X_g^v\left([(G, f)]_g\right) = v\left([(G, f \circ L_g)]_e\right)$$

is smooth because v is a tangent vector (a differential operator) acting on the smooth map $(g, h) \mapsto f(gh)$, which is smooth since the group multiplication is smooth. Consequently $X^v \in \mathfrak{X}(G, \mathcal{T}, \mathcal{A})$.

(d) (Left-invariance of X^v) For any $g, h \in G$ and any germ $[(U, f)]_{gh}$ at gh ,

$$\begin{aligned} X_{gh}^v\left([(U, f)]_{gh}\right) &= v\left([(L_{gh}^{-1}(U), f \circ L_{gh})]_e\right) \\ &= v\left([(L_h^{-1}(L_g^{-1}(U)), (f \circ L_g) \circ L_h)]_e\right) \\ &= X_h^v\left([(L_g^{-1}(U), f \circ L_g)]_h\right), \end{aligned}$$

which, by Lemma 6.6, means X^v is μ -left-invariant. Hence $X^v \in \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$.

(e) (Evaluation at the identity) For any germ $[(U, f)]_e$ at e ,

$$[\text{ev}_e(X^v)]([(U, f)]_e) = X_e^v([(U, f)]_e) = v\left([(L_e^{-1}(U), f \circ L_e)]_e\right) = v([(U, f)]_e),$$

because $L_e = \text{id}_G$ and $f \circ L_e = f$. Thus $\text{ev}_e(X^v) = v$, establishing surjectivity. $\square$

Theorem 6.10 (Lie Algebra of a Lie Group). *Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group. Let $\mathfrak{g} = T_e(G, \mathcal{T}, \mathcal{A})$. For every $v \in \mathfrak{g}$, let $X^v \in \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ denote the unique left-invariant smooth vector field satisfying $X^v|_e = v$. (Existence and uniqueness follows from*

$$\text{ev}_e : \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\}) \xrightarrow{\cong} T_e(G, \mathcal{T}, \mathcal{A})$$

being an isomorphism.) Define a map $[\bullet, \bullet]_{\mathfrak{g}} : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ by

$$\forall v, w \in \mathfrak{g} : [v, w]_{\mathfrak{g}} \stackrel{\text{df}}{=} [X^v, X^w]_{\circ}|_e.$$

Then

$$\left(\mathfrak{g}, \left\{ \dot{+}, \dot{-}, \mathbf{0}_{T_e(G)}, \dot{\cdot}, [\bullet, \bullet]_{\mathfrak{g}} \right\} \right)$$

is a Lie $\mathbb{R}$ -algebra, and $\text{ev}_e : \mathfrak{X}_L(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\}) \rightarrow \mathfrak{g}$ is an isomorphism of Lie algebras.

Proof. From 6.8, we know that $(\mathfrak{X}_L(G), [\bullet, \bullet]_{\circ})$ is a Lie $\mathbb{R}$ -algebra. We have also proven in 6.9 that $\text{ev}_e : \mathfrak{X}_L(G) \rightarrow \mathfrak{g}$ is a linear isomorphism. So it is a bijective vector space homomorphism. Since the Lie algebra axioms are equational, by 4.7, defining the bracket as above allows for a partial transport of Lie structure onto $\mathfrak{g}$, making $(\mathfrak{g}, [\bullet, \bullet]_{\mathfrak{g}})$ a Lie $\mathbb{R}$ -algebra and the map $\text{ev}_e : \mathfrak{X}_L(G) \rightarrow \mathfrak{g}$ a Lie isomorphism. $\square$

Definition 6.11. Let $\mathfrak{g}$ be an Lie $\mathbb{F}$ -algebra. Let $\pi_1 : \mathfrak{g} \rightarrow \mathfrak{gl}(V_1)$ and $\pi_2 : \mathfrak{g} \rightarrow \mathfrak{gl}(V_2)$ be representations. π_1 and π_2 are said to be equivalent iff there exists an linear isomorphism $T : V_1 \rightarrow V_2$ such that:

$$\forall X \in \mathfrak{g} : T \circ \pi_1(X) = \pi_2(X) \circ T.$$

7. MATRIX LIE GROUPS AND LIE ALGEBRAS

Theorem 7.1 (Closed Subgroup Theorem – É. Cartan). *Let $(G, \mathcal{T}, \mathcal{A}, \{\mu, \iota, e\})$ be a Lie group and let $H \subseteq G$ be a subgroup with $G \setminus H \in \mathcal{T}$. Then there exists a unique smooth structure $\mathcal{A}_H$ satisfying the following two properties:*

- (1) $\iota_H : H \hookrightarrow G$ is a smooth embedding.
- (2) $(H, \mathcal{T}|_H, \mathcal{A}_H, \{\mu|_{H \times H}, \iota|_H, e\})$ is a Lie group.

(The proof is very long.)

Definition 7.2. The General $\mathbb{F}$ -Linear set of degree n , denoted by “ $\text{GL}(n; \mathbb{F})$ ”, is defined as:

$$\text{GL}(n; \mathbb{F}) \stackrel{\text{df}}{=} \{\mathbf{A} \in \mathbb{F}^{n,n} : \det(\mathbf{A}) \neq 0\}.$$

Definition 7.3. The standard smooth structure on $\mathbb{R}^{n,n}$, denoted by “ $\mathcal{A}_{\mathbb{R}^{n,n}}$ ”, is defined as the unique maximal smooth atlas on $\mathbb{R}^{n,n}$ that contains the $(\mathbb{R}^{n,n}, \mathcal{T}_{\mathbb{R}^{n^2}})$ -chart $(\mathbb{R}^{n,n}, \text{id}_{\mathbb{R}^{n,n}})$.

Definition 7.4. The General $\mathbb{R}$ -Linear Lie group of degree n , denoted by “ $\text{GL}(n; \mathbb{R})$ ”, is defined as the following:

$$\text{GL}(n; \mathbb{R}) \stackrel{\text{df}}{=} \left(\text{GL}(n; \mathbb{R}), \mathcal{T}_{\mathbb{R}^{n^2}}|_{\text{GL}(n; \mathbb{R})}, \mathcal{A}_{\mathbb{R}^{n,n}}|_{\text{GL}(n; \mathbb{R})}, \{\bullet\bullet, \bullet^{-1}, \mathbf{I}_n\} \right).$$

Definition 7.5. The Special Orthogonal Lie group of degree n , denoted by “ $\text{SO}(n)$ ”, is defined as the following:

$$\text{SO}(n) \stackrel{\text{df}}{=} \left(\text{SO}(n), \mathcal{T}_{\mathbb{R}^{n^2}}|_{\text{SO}(n)}, \mathcal{A}_{\text{SO}(n)}, \{\bullet\bullet, \bullet^{\top}, \mathbf{I}_n\} \right),$$

where

$$\text{SO}(n) \stackrel{\text{df}}{=} \{\mathbf{R} \in \mathbb{R}^{n,n} : \mathbf{R}^{\top} \mathbf{R} = \mathbf{I}_n \text{ and } \det(\mathbf{R}) = 1\},$$

and $\mathcal{A}_{\text{SO}(n)}$ is the unique smooth structure as per 7.1 when considering $\text{SO}(n)$ as a closed subgroup of $\text{GL}(n; \mathbb{R})$.

Definition 7.6. The standard smooth structure on $\mathbb{C}^{n,n}$, denoted by “ $\mathcal{A}_{\mathbb{C}^{n,n}}$ ”, is the unique smooth structure on $\mathbb{C}^{n,n}$ that contains the $(\mathbb{C}^{n,n}, \mathcal{T}_{\mathbb{R}^{2n^2}})$ -chart $(\mathbb{C}^{n,n}, \text{id}_{\mathbb{C}^{n,n}})$, where $\mathbb{C}^{n,n}$ is identified with $\mathbb{R}^{2n^2}$ via the natural map $(a_{ij}) \mapsto (\Re(a_{11}), \Im(a_{11}), \dots, \Re(a_{nn}), \Im(a_{nn}))$.

Definition 7.7. The General $\mathbb{C}$ -Linear Lie group of degree n , denoted by “ $\mathrm{GL}(n; \mathbb{C})$ ”, is defined as the following:

$$\mathrm{GL}(n; \mathbb{C}) \stackrel{\mathrm{df}}{=} \left(\mathrm{GL}(n; \mathbb{C}), \mathcal{T}_{\mathbb{R}^{2n^2}}|_{\mathrm{GL}(n; \mathbb{C})}, \mathcal{A}_{\mathbb{C}^{n,n}}|_{\mathrm{GL}(n; \mathbb{C})}, \{\bullet\bullet, \bullet^{-1}, \mathbf{I}_n\} \right).$$

Definition 7.8. The Special Unitary Lie group of degree n , denoted by “ $\mathrm{SU}(n)$ ”, is defined as the following:

$$\mathrm{SU}(n) \stackrel{\mathrm{df}}{=} \left(\mathrm{SU}(n), \mathcal{T}_{\mathbb{R}^{2n^2}}|_{\mathrm{SU}(n)}, \mathcal{A}_{\mathrm{SU}(n)}, \{\bullet\bullet, \bullet^\dagger, \mathbf{I}_n\} \right),$$

where

$$\mathrm{SU}(n) \stackrel{\mathrm{df}}{=} \left\{ \mathbf{U} \in \mathbb{C}^{n,n} : \mathbf{U}^\dagger \mathbf{U} = \mathbf{I}_n \text{ and } \det(\mathbf{U}) = 1 \right\}.$$

and $\mathcal{A}_{\mathrm{SU}(n)}$ is the unique smooth structure as per 7.1 when considering $\mathrm{SU}(n)$ as a closed subgroup of $\mathrm{GL}(n; \mathbb{C})$.

Definition 7.9. The General $\mathbb{R}$ -Linear Lie algebra of degree n , denoted by “ $\mathfrak{gl}(n; \mathbb{R})$ ”, is defined as the following:

$$\mathfrak{gl}(n; \mathbb{R}) \stackrel{\mathrm{df}}{=} (\mathbb{R}^{n,n}, \{+, -, \mathbf{0}_n, \cdot_{\mathbb{R}}, [\bullet, \bullet]\}).$$

Definition 7.10. The Special Orthogonal Lie algebra of degree n , denoted by “ $\mathfrak{so}(n)$ ”, is defined as the following:

$$\mathfrak{so}(n) \stackrel{\mathrm{df}}{=} (\mathfrak{so}(n), \{+, -, \mathbf{0}_n, \cdot_{\mathbb{R}}, [\bullet, \bullet]\}),$$

where

$$\mathfrak{so}(n) \stackrel{\mathrm{df}}{=} \{ \mathbf{R} \in \mathbb{R}^{n,n} : \mathbf{R}^\top + \mathbf{R} = \mathbf{0}_n \}.$$

Definition 7.11. The General $\mathbb{C}$ -Linear Lie algebra of degree n , denoted by “ $\mathfrak{gl}(n; \mathbb{C})$ ”, is defined as the following:

$$\mathfrak{gl}(n; \mathbb{C}) \stackrel{\mathrm{df}}{=} (\mathbb{C}^{n,n}, \{+, -, \mathbf{0}_n, \cdot_{\mathbb{C}}, [\bullet, \bullet]\}).$$

Definition 7.12. The Special Unitary Lie algebra of degree n , denoted by “ $\mathfrak{su}(n)$ ”, is defined as the following:

$$\mathfrak{su}(n) \stackrel{\mathrm{df}}{=} (\mathfrak{su}(n), \{+, -, \mathbf{0}_n, \cdot_{\mathbb{C}}, [\bullet, \bullet]_{\circ}\}),$$

where

$$\mathfrak{su}(n) \stackrel{\mathrm{df}}{=} \left\{ \mathbf{U} \in \mathbb{C}^{n,n} : \mathbf{U}^\dagger + \mathbf{U} = \mathbf{0}_n \text{ and } \mathrm{tr}(\mathbf{U}) = 0 \right\}.$$

Lemma 7.13 (Hadamard). *Let $U \in \mathcal{T}_{\mathbb{R}^n}$ be a convex set. Let $p \in U$. Let $f : U \rightarrow \mathbb{R}$ be smooth w.r.t. $\mathcal{A}_{\mathbb{R}^n}$. Then there exist smooth functions $g_1, \dots, g_n : U \rightarrow \mathbb{R}$ such that*

$$f(x) = f(p) + \sum_{i=1}^n (x^i - p^i) g_i(x), \quad g_i(p) = \frac{\partial f}{\partial x^i}(p).$$

Lemma 7.14 (Linear Identification of the Tangent Space at the Identity). *Let $G = \mathrm{GL}(n; \mathbb{R})$ and $e = \mathbf{I}_n$. Define $\Phi : T_e G \rightarrow \mathbb{R}^{n,n}$ by*

$$\forall v \in T_e G : \quad \Phi(v) \stackrel{\mathrm{df}}{=} (v(x_{ij}))_{i,j=1}^n,$$

where $x_{ij} : \mathrm{GL}(n; \mathbb{R}) \rightarrow \mathbb{R}$ are the standard coordinate functions. Then Φ is a linear isomorphism $T_e G \cong \mathbb{R}^{n,n}$.

Proof. Because $\mathrm{GL}(n; \mathbb{R})$ is open in $\mathbb{R}^{n,n} \cong \mathbb{R}^{n^2}$, the identity map restricts to a global chart $(\mathrm{GL}(n; \mathbb{R}), \mathrm{id}_{\mathrm{GL}(n; \mathbb{R})})$ in the smooth structure of G . Hence every derivation at e is determined by its action on the x_{ij} .

(1) (Linearity) Each $v \mapsto v(x_{ij})$ is linear, so Φ is linear.

- (2) (Injectivity) Suppose $\Phi(v) = 0$, i.e. $v(x_{ij}) = 0$ for all i, j . For any $f \in C^\infty(G)$, restrict to a convex neighborhood of e . By Hadamard's Lemma (Lemma 7.13), there exist smooth g_{ij} such that

$$f(\mathbf{A}) = f(\mathbf{I}_n) + \sum_{i,j} (a_{ij} - \delta_{ij}) g_{ij}(\mathbf{A}).$$

Applying v , the constant term yields 0. For each summand:

$$v((x_{ij} - \delta_{ij})g_{ij}) = (x_{ij}(e) - \delta_{ij})v(g_{ij}) + g_{ij}(e)v(x_{ij} - \delta_{ij}) = 0,$$

since $x_{ij}(e) = \delta_{ij}$ and $v(x_{ij}) = 0$. Thus $v(f) = 0$ for all f , so $v = 0$.

- (3) (Surjectivity) For $\xi \in \mathbb{R}^{n,n}$, define $v_\xi : C^\infty(G) \rightarrow \mathbb{R}$ by

$$\forall f \in C^\infty(G) : \quad v_\xi(f) \stackrel{\text{df}}{=} \left. \frac{d}{dt} f(\mathbf{I}_n + t\xi) \right|_{t=0}.$$

For small t , $\det(\mathbf{I}_n + t\xi) \neq 0$, so $\mathbf{I}_n + t\xi \in G$. Then $v_\xi(x_{ij}) = \xi_{ij}$, so $\Phi(v_\xi) = \xi$. Thus Φ is surjective.

Therefore $\Phi : T_e G \rightarrow \mathbb{R}^{n,n}$ is a linear isomorphism. $\square$

Lemma 7.15 (Left-Invariant Vector Fields on Matrix Lie Groups). *Let $G = \text{GL}(n; \mathbb{R})$ and $e = \mathbf{I}_n$. Let Φ be the isomorphism from Lemma 7.14. For any $\mathbf{A} \in \mathbb{R}^{n,n}$, define $X^{\mathbf{A}} : C^\infty(G) \rightarrow C^\infty(G)$ by*

$$\forall g \in G, \forall f \in C^\infty(G) : \quad [X^{\mathbf{A}}(f)](g) \stackrel{\text{df}}{=} \left. \frac{d}{dt} f(g(\mathbf{I}_n + t\mathbf{A})) \right|_{t=0}.$$

Then $X^{\mathbf{A}} \in \mathfrak{X}_L(G)$ and $\text{ev}_e(X^{\mathbf{A}})$ is the unique tangent vector satisfying $\Phi(\text{ev}_e(X^{\mathbf{A}})) = \mathbf{A}$, i.e. $X^{\mathbf{A}}$ is the unique left-invariant field corresponding to $\Phi^{-1}(\mathbf{A})$.

Proof. The map $t \mapsto g(\mathbf{I}_n + t\mathbf{A})$ is a smooth curve in G for small t , so the derivative exists and defines a smooth function $X^{\mathbf{A}}f$. Left-invariance: for any $h \in G$,

$$\begin{aligned} [X^{\mathbf{A}}(f \circ L_h)](g) &= \left. \frac{d}{dt} f(hg(\mathbf{I}_n + t\mathbf{A})) \right|_{t=0} \\ &= [X^{\mathbf{A}}(f)](hg) = [(X^{\mathbf{A}}f) \circ L_h](g), \end{aligned}$$

so $X^{\mathbf{A}} \in \mathfrak{X}_L(G)$. Evaluating at the identity on the coordinate functions:

$$[\text{ev}_e(X^{\mathbf{A}})](x_{ij}) = \left. \frac{d}{dt} (\delta_{ij} + tA_{ij}) \right|_{t=0} = A_{ij},$$

hence $\Phi(\text{ev}_e(X^{\mathbf{A}})) = \mathbf{A}$. $\square$

Lemma 7.16 (Commutator of Left-Invariant Fields on Matrix Lie Groups). *Let $G = \text{GL}(n; \mathbb{R})$. For $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{n,n}$, let $X^{\mathbf{A}}, X^{\mathbf{B}}$ be the left-invariant vector fields from Lemma 7.15. Then*

$$[X^{\mathbf{A}}, X^{\mathbf{B}}] = X^{[\mathbf{A}, \mathbf{B}]}, \quad [\mathbf{A}, \mathbf{B}] \stackrel{\text{df}}{=} \mathbf{AB} - \mathbf{BA}.$$

Proof. (1) (Second derivative decomposition) For $f \in C^\infty(G)$ and $g \in G$,

$$[X^{\mathbf{A}}(X^{\mathbf{B}}f)](g) = \left. \frac{\partial^2}{\partial t \partial s} f(g(\mathbf{I}_n + t\mathbf{A})(\mathbf{I}_n + s\mathbf{B})) \right|_{(t,s)=(0,0)}.$$

Expand the matrix product:

$$(\mathbf{I}_n + t\mathbf{A})(\mathbf{I}_n + s\mathbf{B}) = \mathbf{I}_n + t\mathbf{A} + s\mathbf{B} + ts\mathbf{AB}.$$

Hence

$$[X^{\mathbf{A}}(X^{\mathbf{B}}f)](g) = \left. \frac{\partial^2}{\partial t \partial s} f(g(\mathbf{I}_n + t\mathbf{A} + s\mathbf{B} + ts\mathbf{AB})) \right|_{(t,s)=(0,0)}.$$

- (2) (Chain rule expansion) Let $\mathbf{H}(t, s) \stackrel{\text{df}}{=} \mathbf{I}_n + t\mathbf{A} + s\mathbf{B} + ts\mathbf{AB}$. By the chain rule for partial derivatives,

$$\frac{\partial^2}{\partial t \partial s} f(g\mathbf{H}(t, s)) = \sum_{p,q,r,\ell=1}^n \frac{\partial^2 f}{\partial x_{pq} \partial x_{r\ell}}(g\mathbf{H}) \cdot \frac{\partial}{\partial t}(g\mathbf{H})_{pq} \cdot \frac{\partial}{\partial s}(g\mathbf{H})_{r\ell} + \sum_{p,q=1}^n \frac{\partial f}{\partial x_{pq}}(g\mathbf{H}) \cdot \frac{\partial^2}{\partial t \partial s}(g\mathbf{H})_{pq}.$$

Evaluate at $(t, s) = (0, 0)$. Since $\mathbf{H}(0, 0) = \mathbf{I}_n$,

$$\begin{aligned} \left. \frac{\partial}{\partial t}(g\mathbf{H})_{pq} \right|_{(t,s)=(0,0)} &= (g\mathbf{A})_{pq}, \\ \left. \frac{\partial}{\partial s}(g\mathbf{H})_{r\ell} \right|_{(t,s)=(0,0)} &= (g\mathbf{B})_{r\ell}, \\ \left. \frac{\partial^2}{\partial t \partial s}(g\mathbf{H})_{pq} \right|_{(t,s)=(0,0)} &= (g\mathbf{AB})_{pq}. \end{aligned}$$

- (3) (Symmetrize and subtract) The mixed second derivative $\frac{\partial^2}{\partial t \partial s}$ is symmetric in t, s , so the first sum (the quadratic form) is symmetric in $\mathbf{A}, \mathbf{B}$. Therefore

$$[X^{\mathbf{A}}(X^{\mathbf{B}}f)](g) = \underbrace{\sum_{p,q,r,\ell} \frac{\partial^2 f}{\partial x_{pq} \partial x_{r\ell}}(g) \cdot (g\mathbf{A})_{pq} \cdot (g\mathbf{B})_{r\ell}}_{\text{symmetric in } \mathbf{A}, \mathbf{B}} + \sum_{p,q} \frac{\partial f}{\partial x_{pq}}(g) \cdot (g\mathbf{AB})_{pq}.$$

Similarly,

$$[X^{\mathbf{B}}(X^{\mathbf{A}}f)](g) = \underbrace{\sum_{p,q,r,\ell} \frac{\partial^2 f}{\partial x_{pq} \partial x_{r\ell}}(g) \cdot (g\mathbf{B})_{pq} \cdot (g\mathbf{A})_{r\ell}}_{\text{same symmetric part}} + \sum_{p,q} \frac{\partial f}{\partial x_{pq}}(g) \cdot (g\mathbf{BA})_{pq}.$$

- (4) (Commutator) Subtracting cancels the symmetric quadratic form, leaving

$$\begin{aligned} [[X^{\mathbf{A}}, X^{\mathbf{B}}](f)](g) &= [X^{\mathbf{A}}(X^{\mathbf{B}}f)](g) - [X^{\mathbf{B}}(X^{\mathbf{A}}f)](g) \\ &= \sum_{p,q} \frac{\partial f}{\partial x_{pq}}(g) \cdot g(\mathbf{AB} - \mathbf{BA})_{pq}. \end{aligned}$$

- (5) (Rewrite as a single directional derivative) For any matrix $\mathbf{C} \in \mathbb{R}^{n,n}$,

$$[X^{\mathbf{C}}(f)](g) = \left. \frac{d}{d\tau} f(g(\mathbf{I}_n + \tau\mathbf{C})) \right|_{\tau=0} = \sum_{p,q} \frac{\partial f}{\partial x_{pq}}(g) \cdot (g\mathbf{C})_{pq}.$$

Setting $\mathbf{C} = \mathbf{AB} - \mathbf{BA} = [\mathbf{A}, \mathbf{B}]$, we obtain

$$[[X^{\mathbf{A}}, X^{\mathbf{B}}](f)](g) = [X^{[\mathbf{A}, \mathbf{B}]}(f)](g), \quad \forall f \in C^\infty(G), \forall g \in G.$$

Hence $[X^{\mathbf{A}}, X^{\mathbf{B}}] = X^{[\mathbf{A}, \mathbf{B}]}$.

□

Theorem 7.17. $\text{Lie}(\text{GL}(n; \mathbb{R})) \cong \mathfrak{gl}(n; \mathbb{R})$.

Proof. Let $G = \text{GL}(n; \mathbb{R})$, $e = \mathbf{I}_n$. By Lemma 7.14, $\Phi : T_e G \rightarrow \mathbb{R}^{n,n}$ is a linear isomorphism. By Lemma 7.15, for each $\mathbf{A} \in \mathbb{R}^{n,n}$ the field $X^{\mathbf{A}}$ is the unique left-invariant field with $\Phi(\text{ev}_e(X^{\mathbf{A}})) = \mathbf{A}$, i.e. $X^{\mathbf{A}} = X^{\Phi^{-1}(\mathbf{A})}$. By Lemma 7.16, $[X^{\mathbf{A}}, X^{\mathbf{B}}] = X^{[\mathbf{A}, \mathbf{B}]}$.

By the “Lie Algebra of a Lie Group” theorem, the bracket on $\text{Lie}(G) = T_e G$ is $[v, w]_{\text{Lie}} \stackrel{\text{df}}{=} \text{ev}_e([X^v, X^w])$. For $v = \Phi^{-1}(\mathbf{A})$, $w = \Phi^{-1}(\mathbf{B})$ we have $X^v = X^{\mathbf{A}}$, $X^w = X^{\mathbf{B}}$, hence

$$\Phi([v, w]_{\text{Lie}}) = \Phi(\text{ev}_e(X^{[\mathbf{A}, \mathbf{B}]}) = [\mathbf{A}, \mathbf{B}] = [\Phi(v), \Phi(w)].$$

Therefore Φ is a bijective Lie algebra homomorphism $\text{Lie}(\text{GL}(n; \mathbb{R})) \cong \mathfrak{gl}(n; \mathbb{R})$. $\square$

Theorem 7.18. $\text{Lie}(\text{GL}(n; \mathbb{C})) \cong \mathfrak{gl}(n; \mathbb{C})$.

Proof. Identify $\mathbb{C}^{n,n}$ with $\mathbb{R}^{2n^2}$ via the natural real coordinate map. The same construction as in the real case, with coordinate functions $u_{ij} = \Re(x_{ij})$, $v_{ij} = \Im(x_{ij})$, yields a linear isomorphism $\Phi : T_{\mathbf{I}_n} \text{GL}(n; \mathbb{C}) \rightarrow \mathbb{C}^{n,n}$ via $\Phi(v) \stackrel{\text{df}}{=} (v(u_{ij}) + iv(v_{ij}))_{i,j}$ (the injectivity argument uses Hadamard’s Lemma in $2n^2$ real variables).

For $\mathbf{A} \in \mathbb{C}^{n,n}$, define $X^{\mathbf{A}}$ by the same formula $X^{\mathbf{A}}(f)(g) = \left. \frac{d}{dt} f(g(\mathbf{I}_n + t\mathbf{A})) \right|_{t=0}$. The left-invariance proof is identical (Lemma 7.15), and the commutator computation is purely algebraic, independent of the field (Lemma 7.16): $[X^{\mathbf{A}}, X^{\mathbf{B}}] = X^{[\mathbf{A}, \mathbf{B}]}$.

Transporting the bracket from left-invariant fields to $T_{\mathbf{I}_n} \text{GL}(n; \mathbb{C})$ through Φ as above yields $\text{Lie}(\text{GL}(n; \mathbb{C})) \cong \mathfrak{gl}(n; \mathbb{C})$. $\square$

Theorem 7.19. $\text{Lie}(\text{SO}(n)) \cong \mathfrak{so}(n)$.

Proof. By Lemma 7.14, identify $T_{\mathbf{I}_n} \text{GL}(n; \mathbb{R})$ with $\mathbb{R}^{n,n}$ via $\Phi(v) = (v(x_{ij}))_{i,j}$.

Define $F : \text{GL}(n; \mathbb{R}) \rightarrow \text{Sym}(n)$ by $F(\mathbf{A}) \stackrel{\text{df}}{=} \mathbf{A}^\top \mathbf{A}$. Its differential at the identity is

$$dF_{\mathbf{I}_n}(\mathbf{X}) = \left. \frac{d}{dt} (\mathbf{I}_n + t\mathbf{X})^\top (\mathbf{I}_n + t\mathbf{X}) \right|_{t=0} = \mathbf{X}^\top + \mathbf{X}.$$

The group $\text{SO}(n) = F^{-1}(\{\mathbf{I}_n\}) \cap \det^{-1}(\{1\})$. Differentiating $\mathbf{A}^\top \mathbf{A} = \mathbf{I}_n$ at $\mathbf{A} = \mathbf{I}_n$ gives $\mathbf{X}^\top + \mathbf{X} = \mathbf{0}_n$, i.e. $\mathbf{X} \in \text{Null}(dF_{\mathbf{I}_n})$. Differentiating $\det(\mathbf{A}) = 1$ gives $\text{tr}(\mathbf{X}) = 0$, but this is redundant for skew-symmetric matrices (diagonal entries vanish). Conversely, any $\mathbf{X}$ with $\mathbf{X}^\top + \mathbf{X} = \mathbf{0}_n$ exponentiates to a curve in $\text{SO}(n)$. Hence

$$T_{\mathbf{I}_n} \text{SO}(n) = \text{Null}(dF_{\mathbf{I}_n}) = \{\mathbf{X} \in \mathbb{R}^{n,n} : \mathbf{X}^\top + \mathbf{X} = \mathbf{0}_n\} = \mathfrak{so}(n).$$

By Theorem 7.17, the Lie bracket on $\text{Lie}(\text{GL}(n; \mathbb{R})) \cong \mathfrak{gl}(n; \mathbb{R})$ is the matrix commutator. The bracket on the Lie subalgebra $\text{Lie}(\text{SO}(n))$ is the restriction. Since $\mathfrak{so}(n)$ is closed under the matrix commutator (the commutator of skew-symmetric matrices is skew-symmetric), the identification $T_{\mathbf{I}_n} \text{SO}(n) \cong \mathfrak{so}(n)$ is a Lie algebra isomorphism. $\square$

Theorem 7.20. $\text{Lie}(\text{SU}(n)) \cong \mathfrak{su}(n)$.

Proof. As in Theorem 7.18, identify $T_{\mathbf{I}_n} \text{GL}(n; \mathbb{C})$ with $\mathbb{C}^{n,n}$ via the coordinate functions.

Define $G(\mathbf{A}) \stackrel{\text{df}}{=} \mathbf{A}^\dagger \mathbf{A}$ and $h(\mathbf{A}) \stackrel{\text{df}}{=} \det(\mathbf{A})$ on $\text{GL}(n; \mathbb{C})$. Their differentials at the identity are

$$dG_{\mathbf{I}_n}(\mathbf{X}) = \mathbf{X}^\dagger + \mathbf{X}, \quad dh_{\mathbf{I}_n}(\mathbf{X}) = \text{tr}(\mathbf{X}).$$

The group $\text{SU}(n) = G^{-1}(\{\mathbf{I}_n\}) \cap h^{-1}(\{1\})$. Differentiating the defining equations gives $\mathbf{X}^\dagger + \mathbf{X} = \mathbf{0}_n$ and $\text{tr}(\mathbf{X}) = 0$. Conversely, any $\mathbf{X} \in \mathbb{C}^{n,n}$ satisfying both conditions exponentiates to a curve in $\text{SU}(n)$. Hence

$$T_{\mathbf{I}_n} \text{SU}(n) = \text{Null}(dG_{\mathbf{I}_n}) \cap \text{Null}(dh_{\mathbf{I}_n}) = \left\{ \mathbf{X} \in \mathbb{C}^{n,n} : \mathbf{X}^\dagger = -\mathbf{X}, \text{tr}(\mathbf{X}) = 0 \right\} = \mathfrak{su}(n).$$

By Theorem 7.18, the Lie bracket on $\text{Lie}(\text{GL}(n; \mathbb{C})) \cong \mathfrak{gl}(n; \mathbb{C})$ is the matrix commutator. The bracket on $\text{Lie}(\text{SU}(n))$ is the restriction. Since the commutator of traceless skew-Hermitian matrices is again traceless and skew-Hermitian, $\mathfrak{su}(n)$ is closed under the bracket. Thus the identification is a Lie algebra isomorphism. $\square$

Definition 7.21. The Special $\mathbb{C}$ -Linear Lie group of degree n , denoted by “ $\text{SL}(n; \mathbb{C})$ ”, is defined as the following:

$$\text{SL}(n; \mathbb{C}) \stackrel{\text{df}}{=} \left(\text{SL}(n; \mathbb{C}), \mathcal{T}_{\mathbb{R}^{2n^2}}|_{\text{SL}(n; \mathbb{C})}, \mathcal{A}_{\text{SL}(n; \mathbb{C})}, \{\bullet\bullet, \bullet^{-1}, I_n\} \right),$$

where

$$\text{SL}(n; \mathbb{C}) \stackrel{\text{df}}{=} \{\mathbf{A} \in \mathbb{C}^{n,n} : |\det(\mathbf{A})| = 1\},$$

and $\mathcal{A}_{\text{SL}(n; \mathbb{C})}$ is the unique smooth structure as per Theorem 7.1 when considering $\text{SL}(n; \mathbb{C})$ as a closed subgroup of $\text{GL}(n; \mathbb{C})$.

Definition 7.22. The Special $\mathbb{C}$ -Linear Lie algebra of degree n , denoted by “ $\mathfrak{sl}(n; \mathbb{C})$ ”, is defined as the following:

$$\mathfrak{sl}(n; \mathbb{C}) \stackrel{\text{df}}{=} (\mathfrak{sl}(n; \mathbb{C}), \{+, -, \mathbf{0}_n, \cdot_{\mathbb{C}}, [\bullet, \bullet]\}),$$

where

$$\mathfrak{sl}(n; \mathbb{C}) \stackrel{\text{df}}{=} \{\mathbf{X} \in \mathbb{C}^{n,n} : \text{tr}(\mathbf{X}) = 0\}.$$

Definition 7.23 (Adjoint Representation of a Matrix Lie Group). Let $G \subseteq \text{GL}(n; \mathbb{F})$ be a matrix Lie group with Lie algebra $\mathfrak{g} \subseteq \mathbb{F}^{n,n}$. The adjoint representation of G is the map $\text{Ad} : G \rightarrow \text{GL}(\mathfrak{g})$ defined by:

$$\forall g \in G, \forall X \in \mathfrak{g} : [\text{Ad}(g)](X) \stackrel{\text{df}}{=} gXg^{-1}.$$

Definition 7.24 (Adjoint Representation of a Lie Algebra). Let $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]\})$ be a Lie $\mathbb{F}$ -algebra. The adjoint representation of $\mathfrak{g}$ is the map $\text{ad} : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g})$ defined by:

$$\forall X, Y \in \mathfrak{g} : [\text{ad}(X)](Y) \stackrel{\text{df}}{=} [X, Y].$$

8. REPRESENTATION THEORY

Definition 8.1. Let $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{g}}\})$ be a Lie $\mathbb{F}$ -algebra. Let V be an $\mathbb{F}$ -vector space. Recall that $(\mathfrak{gl}(V), \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]_{\mathfrak{gl}(V)}\})$ is a Lie $\mathbb{F}$ -algebra. A representation of $\mathfrak{g}$ on V is a map $\pi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ that satisfies the following two properties:

- (1) ($\mathbb{F}$ -Linearity) $\forall X, Y \in \mathfrak{g}, \forall \lambda \in \mathbb{F} : \pi(X + \lambda \cdot Y) = \pi(X) + \lambda \cdot \pi(Y)$.
- (2) (Preservation of the Bracket) $\forall X, Y \in \mathfrak{g} : \pi([X, Y]_{\mathfrak{g}}) = [\pi(X), \pi(Y)]_{\mathfrak{gl}(V)} = \pi(X) \circ \pi(Y) - \pi(Y) \circ \pi(X)$.

In other words, it is a Lie algebra homomorphism from $\mathfrak{g}$ to $\mathfrak{gl}(V)$. The vector space V is called the representation space. The representation is said to be:

- irreducible iff:

$$W \leq V \text{ and } \forall X \in \mathfrak{g} : [\pi(X)](W) \subseteq W \implies W \in \{\{\mathbf{0}_V\}, V\}.$$

- unitary (when $\mathbb{F} = \mathbb{C}$ and V is an inner product space) iff $\forall X \in \mathfrak{g} : \pi(X)^{\dagger} = -\pi(X)$.

Definition 8.2. Let $(G, \mathcal{T}_G, \mathcal{A}_G, \{\mu, \iota, e\})$ be a Lie group and $(V, \{+, -, \mathbf{0}_V, \cdot\})$ be an $\mathbb{F}$ -vector space. Recall that $(\text{GL}(V), \mathcal{T}_{\text{GL}(V)}, \mathcal{A}_{\text{GL}(V)}, \{\circ, \bullet^{-1}, I_V\})$ is a Lie group. A representation of G on V is a map $\Pi : G \rightarrow \text{GL}(V)$ that satisfies the following properties:

- (1) (Smoothness) Π is smooth w.r.t. the smooth structures of G and $\text{GL}(V)$.

(2) (Preservation of Multiplication) $\forall g, h \in G : \quad \Pi(gh) = \Pi(g) \circ \Pi(h)$.

(3) (Preservation of Identity) $\Pi(e) = I_V$.

In other words, it is a Lie group homomorphism from G to $\mathrm{GL}(V)$. The representation is said to be:

- irreducible iff:

$$W \leq V \text{ and } \forall g \in G : [\Pi(g)](W) \subseteq W \quad \implies \quad W \in \{\{0_V\}, V\}.$$

- unitary (when $\mathbb{F} = \mathbb{C}$ and V is an inner product space) iff $\forall g \in G : \quad \Pi(g)^\dagger \Pi(g) = I_V$.

Definition 8.3 (Derived Representation). Let $G \subseteq \mathrm{GL}(n; \mathbb{F})$ be a matrix Lie group with $\mathrm{Lie}(G) \cong \mathfrak{g} \subseteq \mathbb{F}^{n,n}$. Let $\Pi : G \rightarrow \mathrm{GL}(V)$ be a representation of G on an $\mathbb{F}$ -vector space V . The derived representation of Π is the Lie algebra representation $d\Pi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ defined by:

$$\forall X \in \mathfrak{g}, \forall v \in V : \quad [d\Pi(X)](v) \stackrel{\mathrm{df}}{=} \left. \frac{d}{dt} [\Pi(I_n + tX)](v) \right|_{t=0}.$$

(For sufficiently small t , $I_n + tX \in G$ because $\mathfrak{g}$ is the tangent space at the identity.)

Definition 8.4. A projective representation of a Lie group G on an $\mathbb{F}$ -vector space V is a smooth map $\rho : G \rightarrow \mathrm{GL}(V)$ together with a smooth function $\omega : G \times G \rightarrow \mathbb{F}^\times$ such that:

(1) (Projective Multiplication) $\forall g, h \in G : \quad \rho(g) \circ \rho(h) = \omega(g, h) \rho(gh)$.

(2) (Normalization at the Identity) $\rho(e) = I_V$.

The function ω is called the Schur multiplier of the projective representation.

Definition 8.5. Let $(\mathfrak{g}, \{+, -, 0, \cdot, [\bullet, \bullet]_{\mathfrak{g}}\})$ be a Lie $\mathbb{R}$ -algebra. Its complexification, denoted by " $\mathfrak{g}_{\mathbb{C}}$ ", is defined as:

$$\mathfrak{g}_{\mathbb{C}} \stackrel{\mathrm{df}}{=} \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C} = \{X + iY : X, Y \in \mathfrak{g}\},$$

which is a Lie $\mathbb{C}$ -algebra under the bracket extended $\mathbb{C}$ -bilinearly:

$$\forall X_1, X_2, Y_1, Y_2 \in \mathfrak{g} : \quad [X_1 + iY_1, X_2 + iY_2]_{\mathfrak{g}_{\mathbb{C}}} \stackrel{\mathrm{df}}{=} [X_1, X_2]_{\mathfrak{g}} - [Y_1, Y_2]_{\mathfrak{g}} + i([X_1, Y_2]_{\mathfrak{g}} + [Y_1, X_2]_{\mathfrak{g}}).$$

Definition 8.6. Let $(\mathfrak{g}, \{+, -, 0, \cdot, [\bullet, \bullet]_{\mathfrak{g}}\})$ be a Lie $\mathbb{R}$ -algebra. A complex representation of $\mathfrak{g}$ is a representation $\pi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ where V is a $\mathbb{C}$ -vector space. Equivalently, π extends uniquely to a representation $\pi_{\mathbb{C}} : \mathfrak{g}_{\mathbb{C}} \rightarrow \mathfrak{gl}(V)$ of the complexified algebra via:

$$\forall X, Y \in \mathfrak{g} : \quad \pi_{\mathbb{C}}(X + iY) \stackrel{\mathrm{df}}{=} \pi(X) + i\pi(Y).$$

Proposition 8.7 (Equivalence of Complex and Complexified Representations). *Let*

$$(\mathfrak{g}, \{+, -, 0, \cdot, [\bullet, \bullet]_{\mathfrak{g}}\})$$

be a Lie $\mathbb{R}$ -algebra and V a $\mathbb{C}$ -vector space. There is a bijective correspondence between:

(1) *complex representations $\pi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ (i.e. $\mathbb{R}$ -linear maps preserving the bracket with V a $\mathbb{C}$ -vector space), and*

(2) *$\mathbb{C}$ -linear Lie algebra representations $\Pi : \mathfrak{g}_{\mathbb{C}} \rightarrow \mathfrak{gl}(V)$ of the complexified algebra.*

The correspondence is given by extension and restriction:

- (Extension $\Rightarrow$) *Given π , define $\pi_{\mathbb{C}} : \mathfrak{g}_{\mathbb{C}} \rightarrow \mathfrak{gl}(V)$ by*

$$\forall X, Y \in \mathfrak{g} : \quad \pi_{\mathbb{C}}(X + iY) \stackrel{\mathrm{df}}{=} \pi(X) + i\pi(Y).$$

- (Restriction $\Leftarrow$) Given Π , define $\pi \stackrel{\text{df}}{=} \Pi|_{\mathfrak{g}}$, i.e. $\forall X \in \mathfrak{g} : \pi(X) \stackrel{\text{df}}{=} \Pi(X + i0)$.

These two maps are mutual inverses.

Proof. (1) (Well-definedness of $\pi_{\mathbb{C}}$) For $X + iY \in \mathfrak{g}_{\mathbb{C}}$, the formula $\pi_{\mathbb{C}}(X + iY) = \pi(X) + i\pi(Y)$ defines a unique element of $\mathfrak{gl}(V)$ because $\pi(X), \pi(Y) \in \mathfrak{gl}(V)$ and $\mathfrak{gl}(V)$ is a $\mathbb{C}$ -vector space. Since every element of $\mathfrak{g}_{\mathbb{C}}$ has a unique representation $X + iY$ ($\mathfrak{g}_{\mathbb{C}} \stackrel{\text{df}}{=} \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$), $\pi_{\mathbb{C}}$ is well-defined.

- (2) ($\mathbb{C}$ -linearity of $\pi_{\mathbb{C}}$) For $X + iY \in \mathfrak{g}_{\mathbb{C}}$ and $a + ib \in \mathbb{C}$,

$$\begin{aligned} \pi_{\mathbb{C}}((a + ib)(X + iY)) &= \pi_{\mathbb{C}}((aX - bY) + i(bX + aY)) \\ &= \pi(aX - bY) + i\pi(bX + aY) \\ &= (a\pi(X) - b\pi(Y)) + i(b\pi(X) + a\pi(Y)) \\ &= (a + ib)(\pi(X) + i\pi(Y)) \\ &= (a + ib)\pi_{\mathbb{C}}(X + iY). \end{aligned}$$

- (3) (Bracket preservation by $\pi_{\mathbb{C}}$) For $X_1 + iY_1, X_2 + iY_2 \in \mathfrak{g}_{\mathbb{C}}$,

$$\begin{aligned} \pi_{\mathbb{C}}([X_1 + iY_1, X_2 + iY_2]_{\mathfrak{g}_{\mathbb{C}}}) &= \pi_{\mathbb{C}}([X_1, X_2]_{\mathfrak{g}} - [Y_1, Y_2]_{\mathfrak{g}} + i([X_1, Y_2]_{\mathfrak{g}} + [Y_1, X_2]_{\mathfrak{g}})) \\ &= \pi([X_1, X_2]_{\mathfrak{g}}) - \pi([Y_1, Y_2]_{\mathfrak{g}}) + i(\pi([X_1, Y_2]_{\mathfrak{g}}) + \pi([Y_1, X_2]_{\mathfrak{g}})). \end{aligned}$$

Since π preserves the bracket on $\mathfrak{g}$, $\pi([A, B]_{\mathfrak{g}}) = [\pi(A), \pi(B)]_{\circ}$ for all $A, B \in \mathfrak{g}$. Thus

$$\begin{aligned} \pi_{\mathbb{C}}([X_1 + iY_1, X_2 + iY_2]_{\mathfrak{g}_{\mathbb{C}}}) &= [\pi(X_1), \pi(X_2)]_{\circ} - [\pi(Y_1), \pi(Y_2)]_{\circ} \\ &\quad + i([\pi(X_1), \pi(Y_2)]_{\circ} + [\pi(Y_1), \pi(X_2)]_{\circ}). \end{aligned}$$

On the other hand,

$$\begin{aligned} [\pi_{\mathbb{C}}(X_1 + iY_1), \pi_{\mathbb{C}}(X_2 + iY_2)]_{\circ} &= [\pi(X_1) + i\pi(Y_1), \pi(X_2) + i\pi(Y_2)]_{\circ} \\ &= [\pi(X_1), \pi(X_2)]_{\circ} + i[\pi(X_1), \pi(Y_2)]_{\circ} \\ &\quad + i[\pi(Y_1), \pi(X_2)]_{\circ} + i^2[\pi(Y_1), \pi(Y_2)]_{\circ} \\ &= [\pi(X_1), \pi(X_2)]_{\circ} - [\pi(Y_1), \pi(Y_2)]_{\circ} \\ &\quad + i([\pi(X_1), \pi(Y_2)]_{\circ} + [\pi(Y_1), \pi(X_2)]_{\circ}). \end{aligned}$$

Both expressions coincide, so $\pi_{\mathbb{C}}([z_1, z_2]_{\mathfrak{g}_{\mathbb{C}}}) = [\pi_{\mathbb{C}}(z_1), \pi_{\mathbb{C}}(z_2)]_{\circ}$.

- (4) (Mutual inverses) If we start with π , extend to $\pi_{\mathbb{C}}$, then restrict back to $\mathfrak{g}$: for any $X \in \mathfrak{g}$, $\pi_{\mathbb{C}}|_{\mathfrak{g}}(X) = \pi_{\mathbb{C}}(X + i0) = \pi(X) + i\pi(0) = \pi(X)$. Conversely, if we start with $\Pi : \mathfrak{g}_{\mathbb{C}} \rightarrow \mathfrak{gl}(V)$ (which is $\mathbb{C}$ -linear), restrict to $\pi = \Pi|_{\mathfrak{g}}$, then extend:

$$\pi_{\mathbb{C}}(X + iY) = \pi(X) + i\pi(Y) = \Pi(X) + i\Pi(Y) = \Pi(X) + \Pi(iY) = \Pi(X + iY),$$

using $\mathbb{C}$ -linearity of Π to absorb the i .

□

Definition 8.8 (Weight Space). Let $(\mathfrak{g}, \{+, -, \mathbf{0}, \cdot, [\bullet, \bullet]\})$ be a Lie $\mathbb{C}$ -algebra. Let $\pi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a representation on a $\mathbb{C}$ -vector space V . Let $\mathfrak{h} \subseteq \mathfrak{g}$ be an abelian Lie subalgebra (i.e. $[H_1, H_2] = \mathbf{0}_{\mathfrak{g}}$ for all $H_1, H_2 \in \mathfrak{h}$). For a linear functional $\lambda : \mathfrak{h} \rightarrow \mathbb{C}$, the weight space of V with weight λ is:

$$V_{\lambda} \stackrel{\text{df}}{=} \{v \in V : \forall H \in \mathfrak{h} : [\pi(H)](v) = \lambda(H)v\}.$$

If $V_{\lambda} \neq \{\mathbf{0}_V\}$, then λ is called a weight and any nonzero $v \in V_{\lambda}$ is called a weight vector of weight λ .

9. PHYSICS

Definition 9.1. The Rigid Rotation Group is the Lie group $\mathrm{SO}(3)$.

Theorem 9.2. The Lie group $\mathrm{SO}(3)$ satisfies the following three properties:

- (1) $\mathrm{SO}(3)$ is a Lie group.
- (2) $\mathrm{Lie}(\mathrm{SO}(3)) \cong \mathfrak{so}(3)$.
- (3) An ordered basis $(\mathbf{L}_1, \mathbf{L}_2, \mathbf{L}_3)$ for $\mathfrak{so}(3)$ satisfying $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ is:

$$\mathbf{L}_1 \stackrel{\text{df}}{=} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \mathbf{L}_2 \stackrel{\text{df}}{=} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \quad \mathbf{L}_3 \stackrel{\text{df}}{=} \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Proof. (1) $\mathrm{SO}(3)$ is a closed Lie subgroup of $\mathrm{GL}(3; \mathbb{R})$ (Theorem 7.1), hence a Lie group.

(2) $\mathrm{Lie}(\mathrm{SO}(3)) \cong \mathfrak{so}(3)$ by Theorem 7.19.

- (3) The condition $\mathbf{R}^\top = -\mathbf{R}$ forces the diagonal entries to vanish ($R_{ii} = -R_{ii} \implies R_{ii} = 0$) and the off-diagonal entries to satisfy $R_{ji} = -R_{ij}$. Hence every $\mathbf{R} \in \mathfrak{so}(3)$ can be written as

$$\mathbf{R} = \begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix} \quad (a, b, c \in \mathbb{R}).$$

Comparing with the definition of $\mathbf{L}_1, \mathbf{L}_2, \mathbf{L}_3$,

$$\mathbf{R} = a\mathbf{L}_3 - b\mathbf{L}_2 + c\mathbf{L}_1,$$

so $(\mathbf{L}_1, \mathbf{L}_2, \mathbf{L}_3)$ spans $\mathfrak{so}(3)$. If $a\mathbf{L}_1 + b\mathbf{L}_2 + c\mathbf{L}_3 = \mathbf{0}_3$, reading the (1, 2), (1, 3), and (2, 3) entries gives $a = b = c = 0$, so they are linearly independent. Direct computation verifies the commutation relations:

$$[\mathbf{L}_1, \mathbf{L}_2] = \mathbf{L}_3, \quad [\mathbf{L}_2, \mathbf{L}_3] = \mathbf{L}_1, \quad [\mathbf{L}_3, \mathbf{L}_1] = \mathbf{L}_2.$$

□

Theorem 9.3. The Lie group $\mathrm{SU}(2)$ satisfies the following three properties:

- (1) $\mathrm{SU}(2)$ is a Lie group.
- (2) $\mathrm{Lie}(\mathrm{SU}(2)) \cong \mathfrak{su}(2)$.
- (3) An ordered basis $(\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$ for $\mathfrak{su}(2)$ satisfying $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$ is:

$$\mathbf{T}_1 \stackrel{\text{df}}{=} \frac{1}{2} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \mathbf{T}_2 \stackrel{\text{df}}{=} \frac{1}{2} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \mathbf{T}_3 \stackrel{\text{df}}{=} \frac{1}{2} \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}.$$

Proof. (1) $\mathrm{SU}(2)$ is a closed Lie subgroup of $\mathrm{GL}(2; \mathbb{C})$ (Theorem 7.1), hence a Lie group.

(2) $\mathrm{Lie}(\mathrm{SU}(2)) \cong \mathfrak{su}(2)$ by Theorem 7.20.

- (3) The conditions $\mathbf{U}^\dagger = -\mathbf{U}$ and $\mathrm{tr}(\mathbf{U}) = 0$ mean $\mathbf{U} = \begin{bmatrix} ia & b+ic \\ -b+ic & -ia \end{bmatrix}$ for some $a, b, c \in \mathbb{R}$ (the diagonal entries are purely imaginary and sum to zero; the off-diagonals are complex conjugates with a sign flip). Comparing with the definition of $\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3$,

$$\mathbf{U} = 2(a\mathbf{T}_3 + b\mathbf{T}_2 - c\mathbf{T}_1),$$

so $(\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$ spans $\mathfrak{su}(2)$. If $a\mathbf{T}_1 + b\mathbf{T}_2 + c\mathbf{T}_3 = \mathbf{0}_2$, reading the (1, 1) and (1, 2) entries gives $a = b = c = 0$, so they are linearly independent. Direct computation verifies the commutation relations:

$$[\mathbf{T}_1, \mathbf{T}_2] = \mathbf{T}_3, \quad [\mathbf{T}_2, \mathbf{T}_3] = \mathbf{T}_1, \quad [\mathbf{T}_3, \mathbf{T}_1] = \mathbf{T}_2.$$

□

Definition 9.4 (Covering Map, n -Fold Cover, Double Cover). Let $\tilde{G}$ and G be Lie groups.

(1) A covering map is a smooth surjective homomorphism $p : \tilde{G} \rightarrow G$ satisfying:

- (i) (Local Diffeomorphism) $\forall \tilde{g} \in \tilde{G} : dp_{\tilde{g}} : T_{\tilde{g}}\tilde{G} \rightarrow T_{p(\tilde{g})}G$ is a linear isomorphism.
- (ii) (Discrete Fibers) $\forall g \in G : p^{-1}(\{g\})$ is a discrete subset of $\tilde{G}$.

(2) An n -fold cover (or n -sheeted cover) is a covering map $p : \tilde{G} \rightarrow G$ where every fiber has the same cardinality:

$$\forall g \in G : |p^{-1}(\{g\})| = n.$$

(3) A double cover is a 2-fold cover. Equivalently, $\ker(p) = \{\tilde{e}, \tilde{a}\}$ for some $\tilde{a} \in \tilde{G} \setminus \{\tilde{e}\}$ with $\tilde{a}^2 = \tilde{e}$.

Theorem 9.5 (Double Cover $\mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$). For each $U \in \mathrm{SU}(2)$, define a linear map $\Phi(U) : \mathfrak{su}(2) \rightarrow \mathfrak{su}(2)$ by

$$\forall X \in \mathfrak{su}(2) : [\Phi(U)](X) \stackrel{\mathrm{df}}{=} UXU^\dagger.$$

Coordinatize $\mathfrak{su}(2)$ by the basis $\beta = (\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$, and let $[\Phi(U)]_\beta^\beta \in \mathbb{R}^{3,3}$ denote the matrix of $\Phi(U)$ w.r.t. β . By identifying each operator $\Phi(U)$ with its coordinate matrix $[\Phi(U)]_\beta^\beta$, we regard Φ as a map $\Phi : \mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$. Then:

- (1) $[\Phi(U)]_\beta^\beta \in \mathrm{SO}(3)$ for every $U \in \mathrm{SU}(2)$.
- (2) $\Phi : \mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$ is a smooth group homomorphism: $\Phi(UV) = \Phi(U) \circ \Phi(V)$ and $\Phi(\mathbf{I}_2) = I_{\mathfrak{su}(2)}$.
- (3) $\ker(\Phi) = \{\pm \mathbf{I}_2\}$.
- (4) Φ is surjective.

Thus $\mathrm{SU}(2)$ is a double cover of $\mathrm{SO}(3)$: each rotation in $\mathrm{SO}(3)$ corresponds to exactly two matrices in $\mathrm{SU}(2)$ differing only by a sign.

Proof. (1) (Image in $\mathrm{SO}(3)$) For $U \in \mathrm{SU}(2)$, $U^\dagger = U^{-1}$. For any $X \in \mathfrak{su}(2)$, $X^\dagger = -X$ and $\mathrm{tr}(X) = 0$.

$$[\Phi(U)](X)^\dagger = (UXU^\dagger)^\dagger = UX^\dagger U^\dagger = U(-X)U^\dagger = -[\Phi(U)](X),$$

$$\mathrm{tr}([\Phi(U)](X)) = \mathrm{tr}(UXU^\dagger) = \mathrm{tr}(XU^\dagger U) = \mathrm{tr}(X) = 0.$$

Hence $[\Phi(U)](X) \in \mathfrak{su}(2)$, so $\Phi(U) : \mathfrak{su}(2) \rightarrow \mathfrak{su}(2)$ is well-defined.

For $X, Y \in \mathfrak{su}(2)$,

$$\mathrm{tr}([\Phi(U)](X)^\dagger [\Phi(U)](Y)) = \mathrm{tr}(UX^\dagger U^\dagger UYU^\dagger) = \mathrm{tr}(UX^\dagger YU^\dagger) = \mathrm{tr}(X^\dagger Y).$$

Thus $\Phi(U)$ preserves the trace form on $\mathfrak{su}(2)$. Under the basis β , $\mathfrak{su}(2)$ with this form is isometric to $\mathbb{R}^3$ with the standard dot product, so $[\Phi(U)]_\beta^\beta$ is orthogonal. Moreover, $\det([\Phi(U)]_\beta^\beta) = 1$: the map $U \mapsto \det([\Phi(U)]_\beta^\beta)$ is a continuous homomorphism $\mathrm{SU}(2) \rightarrow \{\pm 1\}$. The explicit parametrization above shows $\mathrm{SU}(2) \cong S^3$ is path-connected; the continuous image of a path-connected space in the discrete set $\{\pm 1\}$ is a single point. Evaluating at $U = \mathbf{I}_2$ gives $\det([\Phi(\mathbf{I}_2)]_\beta^\beta) = \det(\mathbf{I}_3) = 1$, so the image is $\{1\}$. Hence $[\Phi(U)]_\beta^\beta \in \mathrm{SO}(3)$.

(2) (Homomorphism) $\Phi(\mathbf{I}_2)(\mathbf{X}) = \mathbf{I}_2 \mathbf{X} \mathbf{I}_2^\dagger = \mathbf{X}$, so $\Phi(\mathbf{I}_2) = I_{\mathfrak{su}(2)}$. For $\mathbf{U}, \mathbf{V} \in \mathrm{SU}(2)$,

$$\begin{aligned} [\Phi(\mathbf{UV})](\mathbf{X}) &= (\mathbf{UV})\mathbf{X}(\mathbf{UV})^\dagger \\ &= \mathbf{U}(\mathbf{V}\mathbf{X}\mathbf{V}^\dagger)\mathbf{U}^\dagger \\ &= [\Phi(\mathbf{U})](\Phi(\mathbf{V})(\mathbf{X})) \\ &= [(\Phi(\mathbf{U}) \circ \Phi(\mathbf{V}))](\mathbf{X}), \end{aligned}$$

so $\Phi(\mathbf{UV}) = \Phi(\mathbf{U}) \circ \Phi(\mathbf{V})$. Smoothness follows because the formula is polynomial in the entries of $\mathbf{U}$.

(3) (Kernel) If $\mathbf{U} \in \ker(\Phi)$, then $\Phi(\mathbf{U}) = I_{\mathfrak{su}(2)}$, i.e. $\mathbf{U}\mathbf{X}\mathbf{U}^\dagger = \mathbf{X}$ for all $\mathbf{X} \in \mathfrak{su}(2)$. This means $\mathbf{U}$ commutes with every matrix in $\mathfrak{su}(2)$. Since $\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3$ together with $\mathbf{I}_2$ form a basis of $\mathbb{C}^{2,2}$ over $\mathbb{C}$, and $\mathbf{U}$ commutes with each $\mathbf{T}_i$ (and trivially with $\mathbf{I}_2$), $\mathbf{U}$ commutes with all 2×2 matrices. A matrix that commutes with every 2×2 matrix must be a scalar multiple of the identity (e.g. test against the matrix units E_{ij}), so $\mathbf{U} = \lambda \mathbf{I}_2$ for some $\lambda \in \mathbb{C}$. Since $\mathbf{U} \in \mathrm{SU}(2)$, $\det(\mathbf{U}) = 1$ and $\mathbf{U}^\dagger \mathbf{U} = \mathbf{I}_2$ give $|\lambda| = 1$ and $\lambda^2 = 1$, so $\lambda = \pm 1$. Thus $\ker(\Phi) = \{\pm \mathbf{I}_2\}$.

(4) (Surjectivity) $\Phi(\mathrm{SU}(2)) \subseteq \mathrm{SO}(3)$ is a subgroup. Since $\mathrm{SU}(2) \cong S^3$ is path-connected and Φ is continuous (smooth), $\Phi(\mathrm{SU}(2))$ is a path-connected subgroup of $\mathrm{SO}(3)$, hence lies in the identity component $\mathrm{SO}(3)^\circ = \mathrm{SO}(3)$. The differential $d\Phi_{\mathbf{I}_2} : T_{\mathbf{I}_2} \mathrm{SU}(2) \rightarrow T_{\mathbf{I}_3} \mathrm{SO}(3)$ is precisely the Lie algebra isomorphism $\varphi : \mathfrak{su}(2) \rightarrow \mathfrak{so}(3)$ from Theorem 9.6. Since $d\Phi_{\mathbf{I}_2}$ is an isomorphism, Φ is a local diffeomorphism at $\mathbf{I}_2$, so the image contains an open neighborhood of $\mathbf{I}_3$ in $\mathrm{SO}(3)$. An open neighborhood of the identity generates the identity component. Since every rotation in $\mathrm{SO}(3)$ can be joined to $\mathbf{I}_3$ by a continuous path (rotate gradually about its axis), $\mathrm{SO}(3)$ is path-connected; thus $\Phi(\mathrm{SU}(2)) = \mathrm{SO}(3)$.

□

Theorem 9.6 (Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$). Define $\varphi : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2)$ by extending

$$\varphi(\mathbf{L}_1) \stackrel{\mathrm{df}}{=} \mathbf{T}_1, \quad \varphi(\mathbf{L}_2) \stackrel{\mathrm{df}}{=} \mathbf{T}_2, \quad \varphi(\mathbf{L}_3) \stackrel{\mathrm{df}}{=} \mathbf{T}_3$$

$\mathbb{R}$ -linearly. Then φ is a Lie algebra isomorphism.

Proof. (1) (Linear isomorphism) φ maps the basis $(\mathbf{L}_1, \mathbf{L}_2, \mathbf{L}_3)$ of $\mathfrak{so}(3)$ bijectively onto the basis $(\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$ of $\mathfrak{su}(2)$, hence is a linear isomorphism.

(2) (Bracket preservation) Using the commutation relations $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ and $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$,

$$\begin{aligned} \varphi([\mathbf{L}_i, \mathbf{L}_j]) &= \varphi(\varepsilon_{ijk} \mathbf{L}_k) = \varepsilon_{ijk} \varphi(\mathbf{L}_k) = \varepsilon_{ijk} \mathbf{T}_k \\ &= [\mathbf{T}_i, \mathbf{T}_j] = [\varphi(\mathbf{L}_i), \varphi(\mathbf{L}_j)]. \end{aligned}$$

By $\mathbb{R}$ -bilinearity, this extends to all of $\mathfrak{so}(3)$: for $\mathbf{X} = \sum_i x_i \mathbf{L}_i$, $\mathbf{Y} = \sum_j y_j \mathbf{L}_j \in \mathfrak{so}(3)$,

$$\begin{aligned} \varphi([\mathbf{X}, \mathbf{Y}]) &= \varphi\left(\sum_{i,j} x_i y_j [\mathbf{L}_i, \mathbf{L}_j]\right) \\ &= \sum_{i,j} x_i y_j \varphi([\mathbf{L}_i, \mathbf{L}_j]) \\ &= \sum_{i,j} x_i y_j [\varphi(\mathbf{L}_i), \varphi(\mathbf{L}_j)] \\ &= [\varphi(\mathbf{X}), \varphi(\mathbf{Y})]. \end{aligned}$$

□

Definition 9.7 (Cartan–Weyl Basis). The Cartan–Weyl basis of $\mathfrak{sl}(2; \mathbb{C})$ is the ordered $\mathbb{C}$ -basis $(\mathbf{H}, \mathbf{E}, \mathbf{F})$ consisting of the matrices

$$\mathbf{H} \stackrel{\text{df}}{=} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad \mathbf{E} \stackrel{\text{df}}{=} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \mathbf{F} \stackrel{\text{df}}{=} \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}.$$

These satisfy the commutation relations

$$[\mathbf{H}, \mathbf{E}] = 2\mathbf{E}, \quad [\mathbf{H}, \mathbf{F}] = -2\mathbf{F}, \quad [\mathbf{E}, \mathbf{F}] = \mathbf{H}.$$

Theorem 9.8 (Complexification of $\mathfrak{su}(2)$). *The complexification $\mathfrak{su}(2)_{\mathbb{C}}$ is isomorphic to $\mathfrak{sl}(2; \mathbb{C})$ as a complex Lie algebra. Explicitly, the two $\mathbb{C}$ -linear maps*

$$\Phi : \mathfrak{sl}(2; \mathbb{C}) \rightarrow \mathfrak{su}(2)_{\mathbb{C}},$$

$$\Psi : \mathfrak{su}(2)_{\mathbb{C}} \rightarrow \mathfrak{sl}(2; \mathbb{C}),$$

defined on the respective $\mathbb{C}$ -bases by

$$\Phi(\mathbf{H}) \stackrel{\text{df}}{=} 2i\mathbf{T}_3, \quad \Phi(\mathbf{E}) \stackrel{\text{df}}{=} i\mathbf{T}_1 - \mathbf{T}_2, \quad \Phi(\mathbf{F}) \stackrel{\text{df}}{=} i\mathbf{T}_1 + \mathbf{T}_2,$$

$$\Psi(\mathbf{T}_1) \stackrel{\text{df}}{=} \frac{\mathbf{E} + \mathbf{F}}{2i}, \quad \Psi(\mathbf{T}_2) \stackrel{\text{df}}{=} \frac{\mathbf{F} - \mathbf{E}}{2}, \quad \Psi(\mathbf{T}_3) \stackrel{\text{df}}{=} \frac{\mathbf{H}}{2i},$$

are mutually inverse complex Lie algebra isomorphisms.

Proof. Since $(\mathbf{H}, \mathbf{E}, \mathbf{F})$ is a $\mathbb{C}$ -basis of $\mathfrak{sl}(2; \mathbb{C})$ and $(\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$ is a $\mathbb{C}$ -basis of $\mathfrak{su}(2)_{\mathbb{C}}$, extending the given assignments $\mathbb{C}$ -linearly yields well-defined $\mathbb{C}$ -linear maps Φ and Ψ .

- (1) (Bracket preservation for Φ) Using $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk}\mathbf{T}_k$ (Theorem 9.3) and the Cartan–Weyl relations (Definition 9.7):

$$\Phi([\mathbf{H}, \mathbf{E}]) = \Phi(2\mathbf{E}) = 2\Phi(\mathbf{E}) = 2(i\mathbf{T}_1 - \mathbf{T}_2),$$

$$\begin{aligned} [\Phi(\mathbf{H}), \Phi(\mathbf{E})] &= [2i\mathbf{T}_3, i\mathbf{T}_1 - \mathbf{T}_2] \\ &= 2i^2[\mathbf{T}_3, \mathbf{T}_1] - 2i[\mathbf{T}_3, \mathbf{T}_2] \\ &= -2\mathbf{T}_2 - 2i(-\mathbf{T}_1) \\ &= 2(i\mathbf{T}_1 - \mathbf{T}_2). \end{aligned}$$

Thus $\Phi([\mathbf{H}, \mathbf{E}]) = [\Phi(\mathbf{H}), \Phi(\mathbf{E})]$. The remaining two relations are verified analogously:

$$\Phi([\mathbf{H}, \mathbf{F}]) = \Phi(-2\mathbf{F}) = -2\Phi(\mathbf{F}) = -2(i\mathbf{T}_1 + \mathbf{T}_2),$$

$$\begin{aligned} [\Phi(\mathbf{H}), \Phi(\mathbf{F})] &= [2i\mathbf{T}_3, i\mathbf{T}_1 + \mathbf{T}_2] \\ &= 2i^2[\mathbf{T}_3, \mathbf{T}_1] + 2i[\mathbf{T}_3, \mathbf{T}_2] \\ &= -2\mathbf{T}_2 + 2i(-\mathbf{T}_1) \\ &= -2(i\mathbf{T}_1 + \mathbf{T}_2), \end{aligned}$$

and

$$\begin{aligned} \Phi([\mathbf{E}, \mathbf{F}]) &= \Phi(\mathbf{H}) = 2i\mathbf{T}_3, \\ [\Phi(\mathbf{E}), \Phi(\mathbf{F})] &= [i\mathbf{T}_1 - \mathbf{T}_2, i\mathbf{T}_1 + \mathbf{T}_2] \\ &= i[\mathbf{T}_1, \mathbf{T}_2] - i[\mathbf{T}_2, \mathbf{T}_1] \\ &= i\mathbf{T}_3 - i(-\mathbf{T}_3) \\ &= 2i\mathbf{T}_3. \end{aligned}$$

By $\mathbb{C}$ -bilinearity of the bracket, Φ preserves all brackets.

- (2) (Bracket preservation for Ψ) Since Φ is a $\mathbb{C}$ -linear bijection preserving the bracket, its inverse $\Psi = \Phi^{-1}$ automatically preserves the bracket:

$$\forall \mathbf{X}, \mathbf{Y} \in \mathfrak{su}(2)_{\mathbb{C}} : \quad \Psi([\mathbf{X}, \mathbf{Y}]) = \Phi^{-1}([\Phi(\Psi(\mathbf{X})), \Phi(\Psi(\mathbf{Y}))]) = [\Psi(\mathbf{X}), \Psi(\mathbf{Y})].$$

(3) (Mutual inverses) Evaluating $\Psi \circ \Phi$ on the basis of $\mathfrak{sl}(2; \mathbb{C})$:

$$[\Psi \circ \Phi](\mathbf{H}) = \Psi(2i\mathbf{T}_3) = 2i \cdot \frac{\mathbf{H}}{2i} = \mathbf{H},$$

$$[\Psi \circ \Phi](\mathbf{E}) = \Psi(i\mathbf{T}_1 - \mathbf{T}_2) = i \cdot \frac{\mathbf{E} + \mathbf{F}}{2i} - \frac{\mathbf{F} - \mathbf{E}}{2} = \frac{\mathbf{E} + \mathbf{F}}{2} - \frac{\mathbf{F} - \mathbf{E}}{2} = \mathbf{E},$$

$$[\Psi \circ \Phi](\mathbf{F}) = \Psi(i\mathbf{T}_1 + \mathbf{T}_2) = i \cdot \frac{\mathbf{E} + \mathbf{F}}{2i} + \frac{\mathbf{F} - \mathbf{E}}{2} = \frac{\mathbf{E} + \mathbf{F}}{2} + \frac{\mathbf{F} - \mathbf{E}}{2} = \mathbf{F}.$$

Thus $\Psi \circ \Phi = \text{id}_{\mathfrak{sl}(2; \mathbb{C})}$. Similarly, $\Phi \circ \Psi = \text{id}_{\mathfrak{su}(2)_{\mathbb{C}}}$ (verified on $\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3$ by the same computation in reverse).

□

Lemma 9.9 (Raising and Lowering Operators). *Let $(V, \{+, -, \mathbf{0}_V, \cdot\})$ be a $\mathbb{C}$ -vector space. Let $\pi : \mathfrak{sl}(2; \mathbb{C}) \rightarrow \mathfrak{gl}(V)$ be a representation. If $\mathbf{v} \in V \setminus \{\mathbf{0}_V\}$ satisfies $[\pi(\mathbf{H})](\mathbf{v}) = \lambda \mathbf{v}$ for some $\lambda \in \mathbb{C}$, then:*

$$[\pi(\mathbf{H})](\pi(\mathbf{E})(\mathbf{v})) = (\lambda + 2)[\pi(\mathbf{E})(\mathbf{v})],$$

$$[\pi(\mathbf{H})](\pi(\mathbf{F})(\mathbf{v})) = (\lambda - 2)[\pi(\mathbf{F})(\mathbf{v})].$$

Moreover, for each $k \in \mathbb{N}$,

$$[\pi(\mathbf{H})]\left(\left[\pi(\mathbf{F})^k\right](\mathbf{v})\right) = (\lambda - 2k)\left[\pi(\mathbf{F})^k\right](\mathbf{v}),$$

where $\pi(\mathbf{F})^k \stackrel{\text{df}}{=} \pi(\mathbf{F}) \circ \cdots \circ \pi(\mathbf{F})$ (k -fold composition, with $\pi(\mathbf{F})^0 \stackrel{\text{df}}{=} I_V$).

Proof. Using the commutation relation $[\mathbf{H}, \mathbf{E}] = 2\mathbf{E}$ (Definition 9.7) and the fact that π preserves the bracket:

$$\begin{aligned} [\pi(\mathbf{H})](\pi(\mathbf{E})(\mathbf{v})) &= [(\pi(\mathbf{H}) \circ \pi(\mathbf{E}))](\mathbf{v}) \\ &= [(\pi(\mathbf{E}) \circ \pi(\mathbf{H}) + [\pi(\mathbf{H}), \pi(\mathbf{E})])](\mathbf{v}) \\ &= [\pi(\mathbf{E})](\pi(\mathbf{H})(\mathbf{v})) + [\pi([\mathbf{H}, \mathbf{E}])](\mathbf{v}) \\ &= [\pi(\mathbf{E})](\lambda \mathbf{v}) + [\pi(2\mathbf{E})](\mathbf{v}) \\ &= (\lambda + 2)[\pi(\mathbf{E})(\mathbf{v})]. \end{aligned}$$

The formula for $\mathbf{F}$ follows from $[\mathbf{H}, \mathbf{F}] = -2\mathbf{F}$ by the same computation. The iterated formula follows by induction on k : the base case $k = 0$ is $[\pi(\mathbf{H})](\mathbf{v}) = \lambda \mathbf{v}$, and the inductive step uses the $k = 1$ case with $\mathbf{v}$ replaced by $[\pi(\mathbf{F})^{k-1}](\mathbf{v})$. □

Theorem 9.10 (Classification of Irreducible Representations of $\mathfrak{sl}(2; \mathbb{C})$). *Let $(V, \{+, -, \mathbf{0}_V, \cdot\})$ be a finite-dimensional $\mathbb{C}$ -vector space. Let $\pi : \mathfrak{sl}(2; \mathbb{C}) \rightarrow \mathfrak{gl}(V)$ be an irreducible representation. Then there exists a unique $n \in \mathbb{Z}_{\geq 0}$ such that:*

(1) *There exists a nonzero vector $\mathbf{v}_0 \in V$, called a highest weight vector, satisfying*

$$[\pi(\mathbf{E})](\mathbf{v}_0) = \mathbf{0}_V, \quad [\pi(\mathbf{H})](\mathbf{v}_0) = n\mathbf{v}_0.$$

(2) *The set $\left\{ \mathbf{v}_0, [\pi(\mathbf{F})](\mathbf{v}_0), [\pi(\mathbf{F})^2](\mathbf{v}_0), \dots, [\pi(\mathbf{F})^n](\mathbf{v}_0) \right\}$ is a $\mathbb{C}$ -basis of V . In particular, $\dim_{\mathbb{C}} V = n + 1$.*

(3) *For each $k \in \{0, \dots, n\}$,*

$$[\pi(\mathbf{H})]\left(\left[\pi(\mathbf{F})^k\right](\mathbf{v}_0)\right) = (n - 2k)\left[\pi(\mathbf{F})^k\right](\mathbf{v}_0).$$

Proof. Since V is a finite-dimensional $\mathbb{C}$ -vector space, the linear operator $\pi(\mathbf{H}) : V \rightarrow V$ possesses at least one eigenvalue. Pick an eigenvector $\mathbf{w} \in V \setminus \{\mathbf{0}_V\}$ with $[\pi(\mathbf{H})](\mathbf{w}) = \mu \mathbf{w}$ for some $\mu \in \mathbb{C}$.

- (1) (Descend to a lowest weight vector) Consider the sequence $\mathbf{w}$, $[\pi(\mathbf{F})](\mathbf{w})$, $[\pi(\mathbf{F})^2](\mathbf{w})$, $\dots$. By Lemma 9.9, the k -th term is an eigenvector of $\pi(\mathbf{H})$ with eigenvalue $\mu - 2k$. Since eigenvectors with distinct eigenvalues are linearly independent and $\dim_{\mathbb{C}} V < \infty$, there exists a maximal index $m \in \mathbb{Z}_{\geq 0}$ such that $\pi(\mathbf{F})^m \mathbf{w} \neq \mathbf{0}_V$. Define $\mathbf{w}_{\min} \stackrel{\text{df}}{=} [\pi(\mathbf{F})^m](\mathbf{w})$. Then:

$$[\pi(\mathbf{F})](\mathbf{w}_{\min}) = \mathbf{0}_V, \quad [\pi(\mathbf{H})](\mathbf{w}_{\min}) = \mu_{\min} \mathbf{w}_{\min},$$

where $\mu_{\min} \stackrel{\text{df}}{=} \mu - 2m$.

- (2) (Ascend to a highest weight vector) Now consider $\mathbf{w}_{\min}$, $[\pi(\mathbf{E})](\mathbf{w}_{\min})$, $[\pi(\mathbf{E})^2](\mathbf{w}_{\min})$, $\dots$. Each application of $\pi(\mathbf{E})$ raises the eigenvalue by 2 (the computation is symmetric to Lemma 9.9 with $[\mathbf{H}, \mathbf{E}] = 2\mathbf{E}$). Again by finite-dimensionality, there exists a maximal index $r \in \mathbb{Z}_{\geq 0}$ such that $[\pi(\mathbf{E})^r](\mathbf{w}_{\min}) \neq \mathbf{0}_V$. Define $\mathbf{v}_0 \stackrel{\text{df}}{=} [\pi(\mathbf{E})^r](\mathbf{w}_{\min})$. Then:

$$[\pi(\mathbf{E})](\mathbf{v}_0) = \mathbf{0}_V, \quad [\pi(\mathbf{H})](\mathbf{v}_0) = n\mathbf{v}_0,$$

where $n \stackrel{\text{df}}{=} \mu_{\min} + 2r = \mu - 2m + 2r$. Thus $\mathbf{v}_0$ is a highest weight vector, proving part (1) of the theorem.

- (3) (Action formula for $\pi(\mathbf{E})$ on $\pi(\mathbf{F})$ -descendants) We claim that for every $k \in \mathbb{Z}_{\geq 0}$,

$$[\pi(\mathbf{E})]\left([\pi(\mathbf{F})^k](\mathbf{v}_0)\right) = k(n - k + 1)[\pi(\mathbf{F})^{k-1}](\mathbf{v}_0). \quad (1)$$

The proof proceeds by induction on k . For $k = 0$, the right-hand side is $0 \cdot (n + 1) \cdot 0 = 0$, and the left-hand side is $[\pi(\mathbf{E})](\mathbf{v}_0) = \mathbf{0}_V$, so the formula holds. Assume the formula holds for k . Then:

$$\begin{aligned} [\pi(\mathbf{E})]\left([\pi(\mathbf{F})^{k+1}](\mathbf{v}_0)\right) &= [(\pi(\mathbf{E}) \circ \pi(\mathbf{F}))]\left([\pi(\mathbf{F})^k](\mathbf{v}_0)\right) \\ &= [(\pi(\mathbf{F}) \circ \pi(\mathbf{E}) + [\pi(\mathbf{E}), \pi(\mathbf{F})])]\left([\pi(\mathbf{F})^k](\mathbf{v}_0)\right) \\ &= [\pi(\mathbf{F})]\left([\pi(\mathbf{E})]\left([\pi(\mathbf{F})^k](\mathbf{v}_0)\right)\right) + [\pi([\mathbf{E}, \mathbf{F}])]\left([\pi(\mathbf{F})^k](\mathbf{v}_0)\right). \end{aligned}$$

Using the induction hypothesis and $[\mathbf{E}, \mathbf{F}] = \mathbf{H}$:

$$\begin{aligned} [\pi(\mathbf{E})]\left([\pi(\mathbf{F})^{k+1}](\mathbf{v}_0)\right) &= [\pi(\mathbf{F})]\left(k(n - k + 1)[\pi(\mathbf{F})^{k-1}](\mathbf{v}_0)\right) + [\pi(\mathbf{H})]\left([\pi(\mathbf{F})^k](\mathbf{v}_0)\right) \\ &= k(n - k + 1)[\pi(\mathbf{F})^k](\mathbf{v}_0) + (n - 2k)[\pi(\mathbf{F})^k](\mathbf{v}_0) \\ &= (k(n - k + 1) + n - 2k)[\pi(\mathbf{F})^k](\mathbf{v}_0) \\ &= (kn - k^2 + k + n - 2k)[\pi(\mathbf{F})^k](\mathbf{v}_0) \\ &= (k + 1)(n - k)[\pi(\mathbf{F})^k](\mathbf{v}_0), \end{aligned}$$

which is exactly the formula (1) with k replaced by $k + 1$. This completes the induction.

- (4) (Spanning set) Let $W \stackrel{\text{df}}{=} \text{Span}_{\mathbb{C}}\left(\left\{[\pi(\mathbf{F})^k](\mathbf{v}_0) : k \in \mathbb{Z}_{\geq 0}\right\}\right)$. We verify that W is invariant under $\pi(\mathbf{H})$, $\pi(\mathbf{E})$, and $\pi(\mathbf{F})$:

- $\pi(\mathbf{F})$ maps each generator to the next, hence W is $\pi(\mathbf{F})$ -invariant by definition.
- By Lemma 9.9, each $[\pi(\mathbf{F})^k](\mathbf{v}_0)$ is a $\pi(\mathbf{H})$ -eigenvector, so W is $\pi(\mathbf{H})$ -invariant.
- Formula (1) shows that $[\pi(\mathbf{E})]\left([\pi(\mathbf{F})^k](\mathbf{v}_0)\right)$ is a scalar multiple of $[\pi(\mathbf{F})^{k-1}](\mathbf{v}_0) \in W$, so W is $\pi(\mathbf{E})$ -invariant.

Thus W is a nonzero π -invariant subspace of V . Since π is irreducible, $W = V$.

- (5) (Termination and dimension) We show that $[\pi(\mathbf{F})^{n+1}](\mathbf{v}_0) = \mathbf{0}_V$. Setting $k = n + 1$ in (1):

$$[\pi(\mathbf{E})]\left([\pi(\mathbf{F})^{n+1}](\mathbf{v}_0)\right) = (n+1) \cdot 0 \cdot [\pi(\mathbf{F})^n](\mathbf{v}_0) = \mathbf{0}_V.$$

If $[\pi(\mathbf{F})^{n+1}](\mathbf{v}_0) \neq \mathbf{0}_V$, then $[\pi(\mathbf{E})]\left([\pi(\mathbf{F})^{n+1}](\mathbf{v}_0)\right) = \mathbf{0}_V$ by (1), so it would be a nonzero vector annihilated by $\pi(\mathbf{E})$. Its $\pi(\mathbf{H})$ -eigenvalue would be $n - 2(n+1) = -n - 2$. But then repeating the ascent argument (step 2) from this vector would produce another highest weight vector with eigenvalue $-n - 2 + 2r' > n$, contradicting the maximality of n (since $\mathbf{v}_0$ was constructed from $\mathbf{w}_{\min}$ by ascending as far as possible). Hence $[\pi(\mathbf{F})^{n+1}](\mathbf{v}_0) = \mathbf{0}_V$.

Since $W = V$, every vector in V is a linear combination of $\left\{[\pi(\mathbf{F})^k](\mathbf{v}_0) : k \geq 0\right\}$. But $[\pi(\mathbf{F})^k](\mathbf{v}_0) = \mathbf{0}_V$ for all $k > n$, so $V = \text{Span}_{\mathbb{C}}\left(\left\{[\pi(\mathbf{F})^k](\mathbf{v}_0)\right\}_{k=0}^n\right)$.

- (6) (Linear independence) By Lemma 9.9, the vectors $[\pi(\mathbf{F})^k](\mathbf{v}_0)$ for $k = 0, \dots, n$ are eigenvectors of $\pi(\mathbf{H})$ with distinct eigenvalues $n, n-2, \dots, -n$. Eigenvectors with distinct eigenvalues are linearly independent, so this set is a $\mathbb{C}$ -basis of V , and $\dim_{\mathbb{C}} V = n+1$.
- (7) (n is a non-negative integer) The eigenvalues of $\pi(\mathbf{H})$ on V are the $(n+1)$ values $n, n-2, \dots, -n+2, -n$. They are symmetric about 0: if λ is an eigenvalue, then so is $-\lambda$ (as the list shows). Since there are $n+1$ distinct integers spaced by 2, the parameter n must be a non-negative integer.

□

Theorem 9.11 (Integration to $\text{SU}(2)$). *Let $(V, \{+, -, \mathbf{0}_V, \cdot\})$ be a finite-dimensional $\mathbb{C}$ -vector space, and let $\pi : \mathfrak{sl}(2; \mathbb{C}) \rightarrow \mathfrak{gl}(V)$ be an irreducible representation with highest weight $n \in \mathbb{Z}_{\geq 0}$ as classified in Theorem 9.10. Then there exists a unique irreducible representation*

$$\Pi_n : \text{SU}(2) \rightarrow \text{GL}(V)$$

whose differential $d(\Pi_n)_{\mathbf{I}_2} : T_{\mathbf{I}_2} \text{SU}(2) \rightarrow \mathfrak{gl}(V)$ coincides with π under the identification $\mathfrak{sl}(2; \mathbb{C}) \cong \mathfrak{su}(2)_{\mathbb{C}} \cong T_{\mathbf{I}_2} \text{SU}(2)_{\mathbb{C}}$. Explicitly, for every $\mathbf{U} \in \text{SU}(2)$ written as $\mathbf{U} = \exp(\mathbf{X})$ with $\mathbf{X} \in \mathfrak{su}(2)$,

$$\Pi_n(\mathbf{U}) \stackrel{\text{df}}{=} \exp((\pi \circ \Psi)(\mathbf{X})),$$

where $\Psi : \mathfrak{su}(2)_{\mathbb{C}} \rightarrow \mathfrak{sl}(2; \mathbb{C})$ is the complex Lie algebra isomorphism of Theorem 9.8 and $\exp : \mathfrak{gl}(V) \rightarrow \text{GL}(V)$ denotes the matrix exponential.

Proof. The proof proceeds in five steps.

- (1) (Topology of $\text{SU}(2)$) The Lie group $\text{SU}(2)$ is diffeomorphic to the 3-sphere S^3 . Indeed, every $\mathbf{U} \in \text{SU}(2)$ can be written as

$$\mathbf{U} = \begin{bmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{bmatrix}, \quad \alpha, \beta \in \mathbb{C}, \quad |\alpha|^2 + |\beta|^2 = 1,$$

and the map $\mathbf{U} \mapsto (\Re(\alpha), \Im(\alpha), \Re(\beta), \Im(\beta))$ identifies $\text{SU}(2)$ with $S^3 \subseteq \mathbb{R}^4$. Since S^n is simply connected for every $n \geq 2$,

$$\pi_1(\text{SU}(2)) = \{\{1\}\}.$$

- (2) (Exponential map of $\text{SU}(2)$ is surjective) Every compact connected Lie group has surjective exponential map. Concretely for $\text{SU}(2)$: each $\mathbf{X} \in \mathfrak{su}(2)$ satisfies $\mathbf{X}^\dagger = -\mathbf{X}$ and $\text{tr}(\mathbf{X}) = 0$, hence its eigenvalues are $\pm i\lambda$ for some $\lambda \in \mathbb{R}_{\geq 0}$, and a direct computation shows

$$\exp(\mathbf{X}) = \cos(\lambda)\mathbf{I}_2 + \frac{\sin(\lambda)}{\lambda}\mathbf{X} \quad (\lambda > 0),$$

with $\exp(\mathbf{0}_2) = \mathbf{I}_2$. Conversely, given $\mathbf{U} \in \mathrm{SU}(2)$, its eigenvalues are $e^{i\theta}, e^{-i\theta}$ for some $\theta \in [0, \pi]$. If $\mathbf{U} = \mathbf{I}_2$, take $\mathbf{X} = \mathbf{0}_2$. Otherwise, set $\lambda \stackrel{\mathrm{df}}{=} \theta$ and

$$\mathbf{X} \stackrel{\mathrm{df}}{=} \frac{\lambda}{\sin(\lambda)} (\mathbf{U} - \cos(\lambda)\mathbf{I}_2),$$

which satisfies $\mathbf{X}^\dagger = -\mathbf{X}$ and $\mathrm{tr}(\mathbf{X}) = 0$, hence $\mathbf{X} \in \mathfrak{su}(2)$, and $\exp(\mathbf{X}) = \mathbf{U}$. Thus $\exp : \mathfrak{su}(2) \rightarrow \mathrm{SU}(2)$ is surjective.

- (3) (Construction of Π_n) The complexification isomorphism $\Psi : \mathfrak{su}(2)_\mathbb{C} \rightarrow \mathfrak{sl}(2; \mathbb{C})$ (Theorem 9.8) restricts to an $\mathbb{R}$ -linear Lie algebra embedding $\mathfrak{su}(2) \hookrightarrow \mathfrak{sl}(2; \mathbb{C})$ (since $\mathfrak{su}(2)$ is a real form of $\mathfrak{sl}(2; \mathbb{C})$). Define the $\mathbb{R}$ -linear Lie algebra representation

$$\rho_n \stackrel{\mathrm{df}}{=} \pi \circ \Psi|_{\mathfrak{su}(2)} : \mathfrak{su}(2) \rightarrow \mathfrak{gl}(V).$$

For $\mathbf{U} = \exp(\mathbf{X}) \in \mathrm{SU}(2)$ with $\mathbf{X} \in \mathfrak{su}(2)$, define

$$\Pi_n(\mathbf{U}) \stackrel{\mathrm{df}}{=} \exp(\rho_n(\mathbf{X})) \in \mathrm{GL}(V),$$

where $\exp : \mathfrak{gl}(V) \rightarrow \mathrm{GL}(V)$ is the exponential of a linear operator: $\exp(\mathbf{A}) \stackrel{\mathrm{df}}{=} \sum_{k=0}^{\infty} \frac{\mathbf{A}^k}{k!}$.

- (4) (Well-definedness) Every $\mathbf{U} \in \mathrm{SU}(2)$ admits a parametrization

$$\mathbf{U} = \exp(\theta \hat{\mathbf{n}} \bullet \mathbf{T}), \quad \theta \in [0, 2\pi), \quad \hat{\mathbf{n}} = (n_1, n_2, n_3) \in \mathbb{R}^3, \quad |\hat{\mathbf{n}}| = 1,$$

where $\hat{\mathbf{n}} \bullet \mathbf{T} \stackrel{\mathrm{df}}{=} n_1 \mathbf{T}_1 + n_2 \mathbf{T}_2 + n_3 \mathbf{T}_3$, by the surjectivity proved in step 2. This parametrization is unique for all $\mathbf{U} \neq \pm \mathbf{I}_2$: θ is the spectral radius of $\log(\mathbf{U})$ and $\hat{\mathbf{n}}$ is the direction of the traceless skew-Hermitian part. For $\mathbf{U} = \mathbf{I}_2$, every direction $\hat{\mathbf{n}}$ gives $\theta = 0$, but the formula $\Pi_n(\mathbf{I}_2) = \exp(\rho_n(\mathbf{0}_2)) = I_V$ is independent of $\hat{\mathbf{n}}$. The only remaining ambiguity is the element $-\mathbf{I}_2$, which corresponds to $\theta = 2\pi$ (outside the chosen domain) but can nevertheless be represented as $\exp(2\pi \hat{\mathbf{n}} \bullet \mathbf{T})$ for *any* unit vector $\hat{\mathbf{n}}$.

We verify that the definition is consistent at this fiber: for any unit $\hat{\mathbf{n}} \in \mathbb{R}^3$, the element $\hat{\mathbf{n}} \bullet \mathbf{T}$ is conjugate to $\mathbf{T}_3$ via the adjoint action: $\hat{\mathbf{n}} \bullet \mathbf{T} = \mathbf{U} \mathbf{T}_3 \mathbf{U}^\dagger$ for some $\mathbf{U} \in \mathrm{SU}(2)$. Since ρ_n is a Lie algebra homomorphism, $\rho_n(\hat{\mathbf{n}} \bullet \mathbf{T})$ is conjugate to $\rho_n(\mathbf{T}_3)$ and therefore has the same eigenvalues. From $\Psi(\mathbf{T}_3) = \mathbf{H}/(2i)$ (Theorem 9.8) and the eigenvalue formula for $\pi(\mathbf{H})$ (Theorem 9.10), the eigenvalues of $\rho_n(\mathbf{T}_3) = \pi(\mathbf{H})/(2i)$ are

$$\frac{n}{2i}, \frac{n-2}{2i}, \dots, \frac{-n}{2i}, \quad \text{i.e.} \quad -\frac{in}{2}, -\frac{i(n-2)}{2}, \dots, \frac{in}{2}.$$

Hence the eigenvalues of $\rho_n(2\pi \hat{\mathbf{n}} \bullet \mathbf{T})$ are $-i\pi n, -i\pi(n-2), \dots, i\pi n$. Exponentiating,

$$\exp(\rho_n(2\pi \hat{\mathbf{n}} \bullet \mathbf{T})) = \exp(2\pi \rho_n(\hat{\mathbf{n}} \bullet \mathbf{T})) = (-1)^n I_V,$$

since $n, n-2, \dots, -n$ all share the parity of n . Thus $\Pi_n(-\mathbf{I}_2) = (-1)^n I_V$, independent of the choice of $\hat{\mathbf{n}}$ in the parametrization.

For $\mathbf{U} \neq -\mathbf{I}_2$, the parametrization $\mathbf{U} = \exp(\theta \hat{\mathbf{n}} \bullet \mathbf{T})$ with $\theta \in [0, 2\pi)$ is unique, so $\Pi_n(\mathbf{U})$ is unambiguously defined. Therefore Π_n is well-defined on all of $\mathrm{SU}(2)$.

- (5) (Irreducibility) Suppose $W \subseteq V$ is a Π_n -invariant subspace. For any $\mathbf{X} \in \mathfrak{su}(2)$ and $\mathbf{w} \in W$,

$$[\rho_n(\mathbf{X})](\mathbf{w}) = \left. \frac{d}{dt} \right|_{t=0} [\Pi_n(\exp(t\mathbf{X}))](\mathbf{w}) \in W,$$

since W is closed and $[\Pi_n(\exp(t\mathbf{X}))](\mathbf{w}) \in W$ for all t . Thus W is ρ_n -invariant, hence π -invariant (since ρ_n and $\pi \circ \Psi$ determine each other on $\mathfrak{su}(2)$ and extend $\mathbb{C}$ -linearly to all of $\mathfrak{sl}(2; \mathbb{C})$). By irreducibility of π (Theorem 9.10), $W = \{\mathbf{0}_V\}$ or $W = V$. Thus Π_n is irreducible.

The uniqueness statement follows from the equivalence of categories: if $\tilde{\Pi}_n : \mathrm{SU}(2) \rightarrow \mathrm{GL}(V)$ is another representation with $d(\tilde{\Pi}_n)_{\mathbf{I}_2} = \rho_n$, then $\tilde{\Pi}_n = \Pi_n$ since both integrate the same Lie algebra representation on a simply connected domain.

□

Definition 9.12 (Spin Operators). Let $\Pi_n : \mathrm{SU}(2) \rightarrow \mathrm{GL}(V)$ be the irreducible representation of highest weight $n \in \mathbb{Z}_{\geq 0}$ constructed in Theorem 9.11. The spin operators are the linear endomorphisms $S_1, S_2, S_3 \in \mathcal{L}(V)$ defined by

$$S_k \stackrel{\mathrm{df}}{=} i\hbar d(\Pi_n)_{\mathbf{I}_2}(\mathbf{T}_k) = i\hbar \rho_n(\mathbf{T}_k) \quad (k = 1, 2, 3),$$

where $\hbar$ is the reduced Planck constant and $\rho_n = \pi \circ \Psi|_{\mathfrak{su}(2)}$ as in Theorem 9.11. The spin vector is the triple $S \stackrel{\mathrm{df}}{=} (S_1, S_2, S_3)$, and the squared spin operator is $S^2 \stackrel{\mathrm{df}}{=} S_1^2 + S_2^2 + S_3^2 \in \mathcal{L}(V)$.

Proposition 9.13 (Commutation Relations). *The spin operators satisfy the $\mathfrak{su}(2)$ commutation relations:*

$$[S_i, S_j] = i\hbar \varepsilon_{ijk} S_k \quad (i, j, k = 1, 2, 3).$$

Proof. From $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$ (Theorem 9.3) and the fact that ρ_n is a Lie algebra homomorphism:

$$\begin{aligned} [S_i, S_j] &= [i\hbar \rho_n(\mathbf{T}_i), i\hbar \rho_n(\mathbf{T}_j)] \\ &= (i\hbar)^2 \rho_n([\mathbf{T}_i, \mathbf{T}_j]) \\ &= -\hbar^2 \rho_n(\varepsilon_{ijk} \mathbf{T}_k) \\ &= -\hbar^2 \varepsilon_{ijk} \rho_n(\mathbf{T}_k) \\ &= i\hbar \varepsilon_{ijk} (i\hbar \rho_n(\mathbf{T}_k)) \\ &= i\hbar \varepsilon_{ijk} S_k. \end{aligned}$$

□

Proposition 9.14 (Action of S_3 and S^2 on Weight Vectors). *Let $(V, \{+, -, \mathbf{0}_V, \cdot\})$ be the $(n+1)$ -dimensional representation space of Π_n , and let $\{\mathbf{v}_k\}_{k=0}^n$ be the weight basis constructed in Theorem 9.10:*

$$[\rho_n(\mathbf{H})](\mathbf{v}_k) = (n - 2k)\mathbf{v}_k, \quad \mathbf{v}_k \stackrel{\mathrm{df}}{=} [\pi(\mathbf{F})^k](\mathbf{v}_0).$$

Define the spin quantum number $j \stackrel{\mathrm{df}}{=} n/2$ and the magnetic quantum numbers $m_k \stackrel{\mathrm{df}}{=} j - k$ for $k = 0, \dots, n$. Then:

- (1) $S_3(\mathbf{v}_k) = \hbar m_k \mathbf{v}_k$ for $k = 0, \dots, n$. Equivalently, the eigenvalues of S_3 are $\hbar j, \hbar(j-1), \dots, -\hbar j$.
- (2) $S^2(\mathbf{v}_k) = \hbar^2 j(j+1) \mathbf{v}_k$ for $k = 0, \dots, n$. That is, S^2 acts as the scalar $\hbar^2 j(j+1)$ on the entire irreducible representation V .

Proof. (1) From $\Psi(\mathbf{T}_3) = \mathbf{H}/(2i)$ (Theorem 9.8):

$$\rho_n(\mathbf{T}_3) = \pi(\Psi(\mathbf{T}_3)) = \pi\left(\frac{\mathbf{H}}{2i}\right) = \frac{1}{2i}\pi(\mathbf{H}).$$

Hence

$$S_3 = i\hbar \rho_n(\mathbf{T}_3) = i\hbar \cdot \frac{1}{2i} \pi(\mathbf{H}) = \frac{\hbar}{2} \pi(\mathbf{H}).$$

Applying to $\mathbf{v}_k$ and using $[\pi(\mathbf{H})](\mathbf{v}_k) = (n - 2k)\mathbf{v}_k$:

$$S_3(\mathbf{v}_k) = \frac{\hbar}{2}(n - 2k)\mathbf{v}_k = \hbar\left(\frac{n}{2} - k\right)\mathbf{v}_k = \hbar m_k \mathbf{v}_k.$$

Since k runs from 0 to n , the eigenvalues $\hbar j, \hbar(j-1), \dots, -\hbar j$ follow.

- (2) The squared operator S^2 is the quadratic Casimir of $\mathfrak{su}(2)$. Using the commutation relations (Proposition 9.13) and the fact that S^2 commutes with each S_i , it follows from Schur's Lemma (applied to the irreducible representation Π_n) that S^2 acts as a scalar on V . To determine the scalar, evaluate on the highest weight vector $\mathbf{v}_0$ (with $m_0 = j$):

$$\begin{aligned}
S^2(\mathbf{v}_0) &= [S_1^2 + S_2^2 + S_3^2](\mathbf{v}_0) \\
&= [S_3^2 + (S_1 - iS_2)(S_1 + iS_2) - i[S_1, S_2]](\mathbf{v}_0) \\
&= S_3^2(\mathbf{v}_0) + [(S_1 - iS_2)(S_1 + iS_2)](\mathbf{v}_0) - [i \cdot i\hbar S_3](\mathbf{v}_0) \\
&= \hbar^2 j^2 \mathbf{v}_0 + [S_1 - iS_2](\mathbf{0}_V) + \hbar S_3(\mathbf{v}_0) \\
&= \hbar^2 j^2 \mathbf{v}_0 + \mathbf{0}_V + \hbar \cdot \hbar j \mathbf{v}_0 \\
&= \hbar^2 j(j+1) \mathbf{v}_0.
\end{aligned}$$

Here we used $[S_1, S_2] = i\hbar S_3$, $S_3(\mathbf{v}_0) = \hbar j \mathbf{v}_0$, and the fact that $S_1 + iS_2$ annihilates the highest weight vector (since $S_1 + iS_2 = \hbar \pi(\mathbf{E})$ and $[\pi(\mathbf{E})](\mathbf{v}_0) = \mathbf{0}_V$ by Theorem 9.10). Since S^2 commutes with all S_i , the same scalar eigenvalue holds for every basis vector $\mathbf{v}_k$.

□

Definition 9.15 (Projective Representation). Let G be a group and let $(V, \{+, -, \mathbf{0}_V, \cdot\})$ be a $\mathbb{C}$ -vector space. A projective representation of G on V is a map $\Pi : G \rightarrow \mathrm{GL}(V)$ for which there exists a function $c : G \times G \rightarrow \mathbb{C}^\times$ satisfying the following properties:

- (1) (Projective Multiplication) $\forall g_1, g_2 \in G : \quad \Pi(g_1) \circ \Pi(g_2) = c(g_1, g_2) \Pi(g_1 g_2)$.
- (2) (Normalisation) $\Pi(e_G) = I_V$, and for every $g \in G$, $c(e_G, g) = c(g, e_G) = 1$.

The function c is called a 2-cocycle. If $c(g_1, g_2) = 1$ for all $g_1, g_2 \in G$, then Π is an ordinary (non-projective) representation of G .

Remark. In quantum mechanics, physical states are not vectors in a Hilbert space $\mathcal{H}$ but *rays* — equivalence classes $[\psi] \stackrel{\mathrm{df}}{=} \{e^{i\theta}\psi : \theta \in \mathbb{R}\}$ modulo a global phase. By Wigner's Theorem, a symmetry transformation $T : \mathcal{H} \rightarrow \mathcal{H}$ need only preserve transition probabilities: $\forall \psi, \varphi \in \mathcal{H} : |\langle T\psi, T\varphi \rangle| = |\langle \psi, \varphi \rangle|$. Consequently, symmetry *groups* are represented on $\mathcal{H}$ projectively in general — the phase ambiguity in the ray correspondence manifests as the cocycle c .

Proposition 9.16 (Descent Criterion). *Let $\Phi : \mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$ be the double cover of Theorem 9.5, and let $\Pi_n : \mathrm{SU}(2) \rightarrow \mathrm{GL}(V)$ be the irreducible representation of highest weight $n \in \mathbb{Z}_{\geq 0}$ constructed in Theorem 9.11. Then:*

- (1) (*Sign Under Central Element*) $\forall \mathbf{U} \in \mathrm{SU}(2) : \quad \Pi_n(-\mathbf{U}) = (-1)^n \Pi_n(\mathbf{U})$. In particular, $\Pi_n(-\mathbf{I}_2) = (-1)^n I_V$.
- (2) (*Integer Spin — Descent to $\mathrm{SO}(3)$*) If n is even, then Π_n factors through Φ : there exists a unique ordinary representation $\tilde{\Pi}_n : \mathrm{SO}(3) \rightarrow \mathrm{GL}(V)$ such that

$$\tilde{\Pi}_n \circ \Phi = \Pi_n.$$

Explicitly, $\forall \mathbf{R} \in \mathrm{SO}(3) : \tilde{\Pi}_n(\mathbf{R}) \stackrel{\mathrm{df}}{=} \Pi_n(\mathbf{U})$, where $\mathbf{U} \in \mathrm{SU}(2)$ is any preimage satisfying $\Phi(\mathbf{U}) = \mathbf{R}$; the definition is independent of the choice of $\mathbf{U}$ because $\Pi_n(\mathbf{U}) = \Pi_n(-\mathbf{U})$ for even n .

- (3) (*Half-Integer Spin — Projective Descent*) If n is odd, then Π_n induces a projective representation $\rho_n : \mathrm{SO}(3) \rightarrow \mathrm{GL}(V)$ defined by $\forall \mathbf{R} \in \mathrm{SO}(3) : \rho_n(\mathbf{R}) \stackrel{\mathrm{df}}{=} \Pi_n(\mathbf{U})$, where $\mathbf{U} \in \mathrm{SU}(2)$ is an arbitrary lift of $\mathbf{R}$. The two lifts $\pm \mathbf{U}$ differ by a sign: $\Pi_n(-\mathbf{U}) = -\Pi_n(\mathbf{U})$, so ρ_n is projective with the sign cocycle.

Proof. (1) Let $U \in \mathrm{SU}(2)$. Parametrising $-U$ as in Theorem 9.11:

$$-U = \exp(\mathbf{X}) \quad \text{for some } \mathbf{X} \in \mathfrak{su}(2).$$

But $-U = (-I_2)U$, and $-I_2 = \exp(2\pi\hat{\mathbf{n}} \bullet \mathbf{T})$ for any unit $\hat{\mathbf{n}} \in \mathbb{R}^3$ (step 4 of Theorem 9.11). Since $-I_2$ is central in $\mathrm{SU}(2)$,

$$-U = (-I_2)U = \exp(2\pi\hat{\mathbf{n}} \bullet \mathbf{T}) \exp(\mathbf{X}) = \exp(2\pi\hat{\mathbf{n}} \bullet \mathbf{T} + \mathbf{X}),$$

where the last equality uses that $[\hat{\mathbf{n}} \bullet \mathbf{T}, \mathbf{X}] = \mathbf{0}$ (since $-I_2$ is central, its logarithm commutes with every $\mathbf{X}$). Applying Π_n and using the construction in Theorem 9.11:

$$\begin{aligned} \Pi_n(-U) &= \exp(\rho_n(2\pi\hat{\mathbf{n}} \bullet \mathbf{T} + \mathbf{X})) \\ &= \exp(2\pi\rho_n(\hat{\mathbf{n}} \bullet \mathbf{T})) \exp(\rho_n(\mathbf{X})) \\ &= (-1)^n I_V \circ \Pi_n(U) \\ &= (-1)^n \Pi_n(U), \end{aligned}$$

where we used $\exp(2\pi\rho_n(\hat{\mathbf{n}} \bullet \mathbf{T})) = (-1)^n I_V$ (proved in step 4 of Theorem 9.11). The special case $U = I_2$ gives $\Pi_n(-I_2) = (-1)^n I_V$.

(2) For even n , $\Pi_n(-I_2) = I_V$, so for any $\mathbf{R} = \Phi(U) \in \mathrm{SO}(3)$ with $U \in \mathrm{SU}(2)$,

$$\Pi_n(U) = \Pi_n(-(-U)) = \Pi_n(-U) = \Pi_n(U),$$

confirming that the value $\tilde{\Pi}_n(\mathbf{R}) \stackrel{\mathrm{df}}{=} \Pi_n(U)$ is unchanged whether we lift $\mathbf{R}$ to U or to $-U$. It remains to verify that $\tilde{\Pi}_n$ is a group homomorphism: $\forall \mathbf{R}_1, \mathbf{R}_2 \in \mathrm{SO}(3)$, choose lifts $U_1, U_2 \in \mathrm{SU}(2)$. Then $\Phi(U_1 U_2) = \mathbf{R}_1 \mathbf{R}_2$, so

$$\tilde{\Pi}_n(\mathbf{R}_1 \mathbf{R}_2) = \Pi_n(U_1 U_2) = \Pi_n(U_1) \circ \Pi_n(U_2) = \tilde{\Pi}_n(\mathbf{R}_1) \circ \tilde{\Pi}_n(\mathbf{R}_2).$$

Smoothness of $\tilde{\Pi}_n$ follows from the fact that Φ is a local diffeomorphism (Theorem 9.5), so $\tilde{\Pi}_n = \Pi_n \circ s$ for any local smooth section s of Φ .

(3) For odd n , $\Pi_n(-I_2) = -I_V$, so the two lifts $\pm U$ of $\mathbf{R}$ give opposite signs: $\Pi_n(-U) = -\Pi_n(U)$. Thus ρ_n is well-defined only up to sign — a projective representation. Explicitly, for $\mathbf{R}_1, \mathbf{R}_2 \in \mathrm{SO}(3)$ with lifts $U_1, U_2 \in \mathrm{SU}(2)$,

$$\rho_n(\mathbf{R}_1) \circ \rho_n(\mathbf{R}_2) = \Pi_n(U_1) \Pi_n(U_2) = \Pi_n(U_1 U_2).$$

The product $U_1 U_2$ is a lift of $\mathbf{R}_1 \mathbf{R}_2$, so

$$\rho_n(\mathbf{R}_1 \mathbf{R}_2) = \pm \Pi_n(U_1 U_2) = \pm (\rho_n(\mathbf{R}_1) \circ \rho_n(\mathbf{R}_2)),$$

where the sign is $+1$ if the chosen lift of $\mathbf{R}_1 \mathbf{R}_2$ equals $U_1 U_2$, and -1 otherwise. This is precisely the projective multiplication law (Definition 9.15). □

Remark. The spin quantum number $j \stackrel{\mathrm{df}}{=} n/2$ determines the physical interpretation of the descent.

- If $j \in \mathbb{Z}_{\geq 0}$ (equivalently, n is even), the representation descends to an ordinary representation of $\mathrm{SO}(3)$ (Proposition 9.16). These correspond to integer spin (bosonic) particles.
- If $j \in \mathbb{Z}_{\geq 0} + \frac{1}{2}$ (equivalently, n is odd), the representation descends only to a projective representation of $\mathrm{SO}(3)$ (Proposition 9.16). These correspond to half-integer spin (fermionic) particles.

The simplest example is $j = 1/2$ ($n = 1$), the spin- $\frac{1}{2}$ representation, which underlies the quantum mechanics of electrons, protons, and neutrons.