

Representations of Lie Groups

A Mathematical Dissection of Quantum Spin

Part I: Mathematical Foundations

Gordon Chan
Supervisor: Assoc. Prof. François Gay-Balmaz

28 July 2026

Outline

- 1 Topological Spaces
- 2 Topological Manifolds
- 3 Smooth Manifolds
- 4 Tangent Spaces
- 5 Algebraic Structures
- 6 Lie Groups
- 7 Matrix Lie Groups
- 8 The Rotation Group $SO(3)$
- 9 The Special Unitary Group $SU(2)$
- 10 The Double Cover $SU(2) \rightarrow SO(3)$
- 11 Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$
- 12 Summary and Next Steps

Definition (topological space)

A **topological space** is a pair $(X, \mathcal{T})$, where X is a set and $\mathcal{T} \subseteq \mathcal{P}(X)$ satisfies:

- 1 (Trivial bounds) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.
- 2 (Arbitrary unions) $\forall \{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T} : \bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}$.
- 3 (Finite intersections) $\forall U, V \in \mathcal{T} : U \cap V \in \mathcal{T}$.

Topological Spaces

Definition (topological space)

A **topological space** is a pair $(X, \mathcal{T})$, where X is a set and $\mathcal{T} \subseteq \mathcal{P}(X)$ satisfies:

- ① (Trivial bounds) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.
- ② (Arbitrary unions) $\forall \{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T} : \bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}$.
- ③ (Finite intersections) $\forall U, V \in \mathcal{T} : U \cap V \in \mathcal{T}$.

Why this matters. A topology tells us what “closeness” means without needing a distance function. Open sets are the primitive notion of “nearby” — they generalise open intervals in $\mathbb{R}$.

Topological Spaces

Definition (topological space)

A **topological space** is a pair $(X, \mathcal{T})$, where X is a set and $\mathcal{T} \subseteq \mathcal{P}(X)$ satisfies:

- ① (Trivial bounds) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.
- ② (Arbitrary unions) $\forall \{U_\alpha\}_{\alpha \in \mathcal{I}} \subseteq \mathcal{T} : \bigcup_{\alpha \in \mathcal{I}} U_\alpha \in \mathcal{T}$.
- ③ (Finite intersections) $\forall U, V \in \mathcal{T} : U \cap V \in \mathcal{T}$.

Why this matters. A topology tells us what “closeness” means without needing a distance function. Open sets are the primitive notion of “nearby” — they generalise open intervals in $\mathbb{R}$.

Key constructions. Product topology, subspace topology, metric topologies (e.g. the Euclidean topology on $\mathbb{R}^n$).

Definition (topological manifold)

A **topological manifold** is a topological space $(M, \mathcal{T})$ satisfying:

- ① (Hausdorff) Points can be separated by disjoint open sets.
- ② (Second-countable) The topology has a countable basis.
- ③ (Locally Euclidean) Every point has a neighbourhood homeomorphic to $\mathbb{R}^n$ for some fixed $n = \dim(M)$.

Definition (topological manifold)

A **topological manifold** is a topological space $(M, \mathcal{T})$ satisfying:

- ① (Hausdorff) Points can be separated by disjoint open sets.
- ② (Second-countable) The topology has a countable basis.
- ③ (Locally Euclidean) Every point has a neighbourhood homeomorphic to $\mathbb{R}^n$ for some fixed $n = \dim(M)$.

Intuition. A manifold is a space that *locally looks like* $\mathbb{R}^n$, but may have a nontrivial global shape (e.g. a sphere S^2 or a torus T^2).

Topological Manifolds

Definition (topological manifold)

A **topological manifold** is a topological space $(M, \mathcal{T})$ satisfying:

- ① (Hausdorff) Points can be separated by disjoint open sets.
- ② (Second-countable) The topology has a countable basis.
- ③ (Locally Euclidean) Every point has a neighbourhood homeomorphic to $\mathbb{R}^n$ for some fixed $n = \dim(M)$.

Intuition. A manifold is a space that *locally looks like* $\mathbb{R}^n$, but may have a nontrivial global shape (e.g. a sphere S^2 or a torus T^2).

Why Hausdorff + second-countable? These are technical conditions that prevent pathological spaces. They guarantee that partitions of unity exist.

Definition (smooth manifold)

A **smooth manifold** is a triple $(M, \mathcal{T}, \mathcal{A})$ where:

- ① $(M, \mathcal{T})$ is a topological manifold.
- ② $\mathcal{A} = \{(U_\alpha, \varphi_\alpha)\}_{\alpha \in \mathcal{I}}$ is a **smooth atlas**: each $(U_\alpha, \varphi_\alpha)$ is a chart, the charts cover M , and transition maps $\varphi_\beta \circ \varphi_\alpha^{-1}$ are C^∞ .
- ③ $\mathcal{A}$ is **maximal** with respect to smooth compatibility.

Definition (smooth manifold)

A **smooth manifold** is a triple $(M, \mathcal{T}, \mathcal{A})$ where:

- 1 $(M, \mathcal{T})$ is a topological manifold.
- 2 $\mathcal{A} = \{(U_\alpha, \varphi_\alpha)\}_{\alpha \in \mathcal{I}}$ is a **smooth atlas**: each $(U_\alpha, \varphi_\alpha)$ is a chart, the charts cover M , and transition maps $\varphi_\beta \circ \varphi_\alpha^{-1}$ are C^∞ .
- 3 $\mathcal{A}$ is **maximal** with respect to smooth compatibility.

Key idea. Charts are local coordinate systems. Smoothness means that when you switch between coordinate systems, the change-of-coordinates map is infinitely differentiable.

Smooth Maps and Diffeomorphisms

Definition (smooth map)

A map $F : M \rightarrow N$ between smooth manifolds is **smooth** if its coordinate representation $\psi \circ F \circ \varphi^{-1}$ is C^∞ for all compatible charts.

Smooth Maps and Diffeomorphisms

Definition (smooth map)

A map $F : M \rightarrow N$ between smooth manifolds is **smooth** if its coordinate representation $\psi \circ F \circ \varphi^{-1}$ is C^∞ for all compatible charts.

Definition (diffeomorphism)

$F : M \rightarrow N$ is a **diffeomorphism** if it is smooth, bijective, and F^{-1} is also smooth. Diffeomorphic manifolds are “the same” from the smooth perspective.

Smooth Maps and Diffeomorphisms

Definition (smooth map)

A map $F : M \rightarrow N$ between smooth manifolds is **smooth** if its coordinate representation $\psi \circ F \circ \varphi^{-1}$ is C^∞ for all compatible charts.

Definition (diffeomorphism)

$F : M \rightarrow N$ is a **diffeomorphism** if it is smooth, bijective, and F^{-1} is also smooth. Diffeomorphic manifolds are “the same” from the smooth perspective.

Example. $\mathbb{R}^n$ is diffeomorphic to the open unit ball $B_1(0)$. S^n is *not* diffeomorphic to $\mathbb{R}^n$ (compact vs. non-compact).

Tangent Spaces via Derivations

Motivation

How do we define tangent vectors to a smooth manifold without embedding it in $\mathbb{R}^N$? Answer: **derivations** — linear maps that satisfy the Leibniz rule.

Tangent Spaces via Derivations

Motivation

How do we define tangent vectors to a smooth manifold without embedding it in $\mathbb{R}^N$? Answer: **derivations** — linear maps that satisfy the Leibniz rule.

Definition (tangent space)

Let M be a smooth manifold, $p \in M$. The **stalk** $C_p^\infty M$ is the set of germs of smooth functions at p . The **tangent space** at p is

$$T_p M \stackrel{\text{df}}{=} \{v : C_p^\infty M \rightarrow \mathbb{R} : v \text{ is a derivation at } p\}.$$

$T_p M$ is an $\mathbb{R}$ -vector space with $\dim(T_p M) = \dim(M)$.

Tangent Spaces via Derivations

Motivation

How do we define tangent vectors to a smooth manifold without embedding it in $\mathbb{R}^N$? Answer: **derivations** — linear maps that satisfy the Leibniz rule.

Definition (tangent space)

Let M be a smooth manifold, $p \in M$. The **stalk** $C_p^\infty M$ is the set of germs of smooth functions at p . The **tangent space** at p is

$$T_p M \stackrel{\text{df}}{=} \{v : C_p^\infty M \rightarrow \mathbb{R} \mid v \text{ is a derivation at } p\}.$$

$T_p M$ is an $\mathbb{R}$ -vector space with $\dim(T_p M) = \dim(M)$.

Why derivations? For $\mathbb{R}^n$, a tangent vector v acts on functions by the directional derivative: $v(f) = \sum_i v^i \frac{\partial f}{\partial x^i} \Big|_p$. This algebraic characterisation generalises to any manifold.

The Differential

Definition (differential)

Let $F : M \rightarrow N$ be smooth, $p \in M$. The **differential of F at p** is the linear map $dF_p : T_p M \rightarrow T_{F(p)} N$ defined by

$$\forall v \in T_p M, \forall [f]_{F(p)} \in C_{F(p)}^\infty N : \quad [(dF_p(v))]([f]_{F(p)}) \stackrel{\text{df}}{=} v([f \circ F]_p).$$

The Differential

Definition (differential)

Let $F : M \rightarrow N$ be smooth, $p \in M$. The **differential of F at p** is the linear map $dF_p : T_p M \rightarrow T_{F(p)} N$ defined by

$$\forall v \in T_p M, \forall [f]_{F(p)} \in C_{F(p)}^\infty N : \quad [(dF_p(v))]([f]_{F(p)}) \stackrel{\text{df}}{=} v([f \circ F]_p).$$

Chain rule

For smooth maps $F : M \rightarrow N$ and $G : N \rightarrow P$,

$$d(G \circ F)_p = dG_{F(p)} \circ dF_p.$$

The Differential

Definition (differential)

Let $F : M \rightarrow N$ be smooth, $p \in M$. The **differential of F at p** is the linear map $dF_p : T_p M \rightarrow T_{F(p)} N$ defined by

$$\forall v \in T_p M, \forall [f]_{F(p)} \in C_{F(p)}^\infty N : \quad [(dF_p(v))]([f]_{F(p)}) \stackrel{\text{df}}{=} v([f \circ F]_p).$$

Chain rule

For smooth maps $F : M \rightarrow N$ and $G : N \rightarrow P$,

$$d(G \circ F)_p = dG_{F(p)} \circ dF_p.$$

Interpretation. The differential is the “best linear approximation” to a smooth map, generalising the Jacobian matrix from multivariable calculus.

Groups, Fields, and Vector Spaces

Definition (group)

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Groups, Fields, and Vector Spaces

Definition (group)

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Definition (field)

A set $\mathbb{F}$ (e.g. $\mathbb{R}, \mathbb{C}$) with addition, multiplication, and all the usual arithmetic axioms — a domain where linear algebra works.

Groups, Fields, and Vector Spaces

Definition (group)

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Definition (field)

A set $\mathbb{F}$ (e.g. $\mathbb{R}, \mathbb{C}$) with addition, multiplication, and all the usual arithmetic axioms — a domain where linear algebra works.

Definition (vector space)

An abelian group $(V, +)$ with scalar multiplication $\mathbb{F} \times V \rightarrow V$ satisfying distributivity and compatibility axioms.

Groups, Fields, and Vector Spaces

Definition (group)

A set G with an associative binary operation μ , an identity e , and inverses ι .
A group is the algebraic abstraction of **symmetry**.

Definition (field)

A set $\mathbb{F}$ (e.g. $\mathbb{R}, \mathbb{C}$) with addition, multiplication, and all the usual arithmetic axioms — a domain where linear algebra works.

Definition (vector space)

An abelian group $(V, +)$ with scalar multiplication $\mathbb{F} \times V \rightarrow V$ satisfying distributivity and compatibility axioms.

Why these matter. Lie groups combine the smooth structure of manifolds with the algebraic structure of groups. Lie algebras are vector spaces with an extra bracket operation.

Definition ($\mathbb{F}$ -algebra)

A vector space $\mathcal{A}$ over $\mathbb{F}$ together with a bilinear multiplication $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$.
(Examples: $\mathbb{R}^{n,n}$ with matrix multiplication, $C^\infty(M)$ with pointwise multiplication.)

Algebras and Lie Algebras

Definition ($\mathbb{F}$ -algebra)

A vector space $\mathcal{A}$ over $\mathbb{F}$ together with a bilinear multiplication $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$. (Examples: $\mathbb{R}^{n,n}$ with matrix multiplication, $C^\infty(M)$ with pointwise multiplication.)

Definition (Lie algebra)

An $\mathbb{F}$ -vector space $\mathfrak{g}$ with a bilinear bracket $[\bullet, \bullet] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying:

- 1 (Skew-symmetry) $[Y, X] = -[X, Y]$
- 2 (Jacobi identity) $[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$

Algebras and Lie Algebras

Definition ($\mathbb{F}$ -algebra)

A vector space $\mathcal{A}$ over $\mathbb{F}$ together with a bilinear multiplication $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$. (Examples: $\mathbb{R}^{n,n}$ with matrix multiplication, $C^\infty(M)$ with pointwise multiplication.)

Definition (Lie algebra)

An $\mathbb{F}$ -vector space $\mathfrak{g}$ with a bilinear bracket $[\bullet, \bullet] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying:

- ① (Skew-symmetry) $[Y, X] = -[X, Y]$
- ② (Jacobi identity) $[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$

Key example. $(\mathbb{R}^3, \times)$ — the cross product makes $\mathbb{R}^3$ a Lie algebra. $[v, w] = v \times w$ satisfies skew-symmetry and Jacobi.

Lie Groups

Definition (Lie group)

A **Lie group** is a smooth manifold $(G, \mathcal{T}, \mathcal{A})$ that is also a group $(G, \{\mu, \iota, e\})$ such that:

- 1 $\mu : G \times G \rightarrow G$ (multiplication) is smooth.
- 2 $\iota : G \rightarrow G$ (inversion) is smooth.

Lie Groups

Definition (Lie group)

A **Lie group** is a smooth manifold $(G, \mathcal{T}, \mathcal{A})$ that is also a group $(G, \{\mu, \iota, e\})$ such that:

- 1 $\mu : G \times G \rightarrow G$ (multiplication) is smooth.
- 2 $\iota : G \rightarrow G$ (inversion) is smooth.

The synthesis. A Lie group unifies geometry (smooth manifold) with algebra (group). This allows us to study continuous symmetries using both differential calculus and group theory.

Lie Groups

Definition (Lie group)

A **Lie group** is a smooth manifold $(G, \mathcal{T}, \mathcal{A})$ that is also a group $(G, \{\mu, \iota, e\})$ such that:

- 1 $\mu : G \times G \rightarrow G$ (multiplication) is smooth.
- 2 $\iota : G \rightarrow G$ (inversion) is smooth.

The synthesis. A Lie group unifies geometry (smooth manifold) with algebra (group). This allows us to study continuous symmetries using both differential calculus and group theory.

Examples.

- $(\mathbb{R}^n, +)$ — the translation group.
- $\mathrm{GL}(n, \mathbb{R})$ — invertible $n \times n$ real matrices.
- $\mathrm{SO}(n)$ — rotations in $\mathbb{R}^n$.
- $\mathrm{SU}(n)$ — unitary transformations with determinant 1.

Left Translation and Left-Invariant Vector Fields

Definition (left translation)

For $g \in G$, $L_g : G \rightarrow G$ is defined by $L_g(h) \stackrel{\text{df}}{=} gh$. L_g is a diffeomorphism.

Left Translation and Left-Invariant Vector Fields

Definition (left translation)

For $g \in G$, $L_g : G \rightarrow G$ is defined by $L_g(h) \stackrel{\text{df}}{=} gh$. L_g is a diffeomorphism.

Definition (left-invariant vector field)

A smooth vector field X on G is **left-invariant** if

$$\forall g \in G, \forall f \in C^\infty(G) : \quad X(f \circ L_g) = X(f) \circ L_g.$$

Left Translation and Left-Invariant Vector Fields

Definition (left translation)

For $g \in G$, $L_g : G \rightarrow G$ is defined by $L_g(h) \stackrel{\text{df}}{=} gh$. L_g is a diffeomorphism.

Definition (left-invariant vector field)

A smooth vector field X on G is **left-invariant** if

$$\forall g \in G, \forall f \in C^\infty(G) : \quad X(f \circ L_g) = X(f) \circ L_g.$$

Geometric meaning. A left-invariant vector field is completely determined by its value at the identity e . Translating the vector at e by L_g gives the vector at any point g .

The Lie Algebra of a Lie Group

Theorem (tangent space at identity is a Lie algebra)

Let G be a Lie group. Define $\mathfrak{g} = T_e G$ (tangent space at identity). For each $v \in \mathfrak{g}$, there exists a **unique** left-invariant vector field X^v with $X^v|_e = v$. Define the bracket on $\mathfrak{g}$ by

$$[v, w]_{\mathfrak{g}} \stackrel{\text{df}}{=} [X^v, X^w]|_e.$$

Then $(\mathfrak{g}, [\bullet, \bullet]_{\mathfrak{g}})$ is a Lie $\mathbb{R}$ -algebra with $\dim(\mathfrak{g}) = \dim(G)$.

The Lie Algebra of a Lie Group

Theorem (tangent space at identity is a Lie algebra)

Let G be a Lie group. Define $\mathfrak{g} = T_e G$ (tangent space at identity). For each $v \in \mathfrak{g}$, there exists a **unique** left-invariant vector field X^v with $X^v|_e = v$. Define the bracket on $\mathfrak{g}$ by

$$[v, w]_{\mathfrak{g}} \stackrel{\text{df}}{=} [X^v, X^w]_o|_e.$$

Then $(\mathfrak{g}, [\bullet, \bullet]_{\mathfrak{g}})$ is a Lie $\mathbb{R}$ -algebra with $\dim(\mathfrak{g}) = \dim(G)$.

The bridge. Every Lie group gives rise to a Lie algebra (its tangent space at the identity). The Lie algebra captures the **infinitesimal** structure of the group.

The Lie Algebra of a Lie Group

Theorem (tangent space at identity is a Lie algebra)

Let G be a Lie group. Define $\mathfrak{g} = T_e G$ (tangent space at identity). For each $v \in \mathfrak{g}$, there exists a **unique** left-invariant vector field X^v with $X^v|_e = v$. Define the bracket on $\mathfrak{g}$ by

$$[v, w]_{\mathfrak{g}} \stackrel{\text{df}}{=} [X^v, X^w]_o|_e.$$

Then $(\mathfrak{g}, [\bullet, \bullet]_{\mathfrak{g}})$ is a Lie $\mathbb{R}$ -algebra with $\dim(\mathfrak{g}) = \dim(G)$.

The bridge. Every Lie group gives rise to a Lie algebra (its tangent space at the identity). The Lie algebra captures the **infinitesimal** structure of the group.

Remark. We denote the Lie algebra of a Lie group G by $\text{Lie}(G)$.

The Closed Subgroup Theorem (É. Cartan)

Theorem (Closed Subgroup Theorem)

Let G be a Lie group and $H \subseteq G$ a closed subgroup. Then H admits a unique smooth structure making it a Lie group such that the inclusion $H \hookrightarrow G$ is a smooth embedding.

The Closed Subgroup Theorem (É. Cartan)

Theorem (Closed Subgroup Theorem)

Let G be a Lie group and $H \subseteq G$ a closed subgroup. Then H admits a unique smooth structure making it a Lie group such that the inclusion $H \hookrightarrow G$ is a smooth embedding.

Powerful consequence. To show that a subset of matrices is a Lie group, you only need to check that it is a closed subgroup of some $GL(n, \mathbb{F})$. No need to construct charts explicitly.

The Closed Subgroup Theorem (É. Cartan)

Theorem (Closed Subgroup Theorem)

Let G be a Lie group and $H \subseteq G$ a closed subgroup. Then H admits a unique smooth structure making it a Lie group such that the inclusion $H \hookrightarrow G$ is a smooth embedding.

Powerful consequence. To show that a subset of matrices is a Lie group, you only need to check that it is a closed subgroup of some $\mathrm{GL}(n, \mathbb{F})$. No need to construct charts explicitly.

Example. $\mathrm{SO}(n)$, $\mathrm{SU}(n)$, $\mathrm{SL}(n, \mathbb{R})$, and $\mathrm{Sp}(2n)$ are all closed subgroups of $\mathrm{GL}(N, \mathbb{F})$ and therefore Lie groups.

The General Linear Group $GL(n, \mathbb{F})$

Example (the general linear group)

$$GL(n, \mathbb{F}) = \{\mathbf{A} \in \mathbb{F}^{n,n} : \det(\mathbf{A}) \neq 0\}.$$

The General Linear Group $GL(n, \mathbb{F})$

Example (the general linear group)

$$GL(n, \mathbb{F}) = \{\mathbf{A} \in \mathbb{F}^{n,n} : \det(\mathbf{A}) \neq 0\}.$$

- $GL(n, \mathbb{F})$ is an open subset of $\mathbb{F}^{n,n} \cong \mathbb{F}^{n^2}$, hence a smooth manifold of dimension n^2 .

The General Linear Group $GL(n, \mathbb{F})$

Example (the general linear group)

$$GL(n, \mathbb{F}) = \{\mathbf{A} \in \mathbb{F}^{n,n} : \det(\mathbf{A}) \neq 0\}.$$

- $GL(n, \mathbb{F})$ is an open subset of $\mathbb{F}^{n,n} \cong \mathbb{F}^{n^2}$, hence a smooth manifold of dimension n^2 .
- Its Lie algebra is $\mathfrak{gl}(n, \mathbb{F}) = \mathbb{F}^{n,n}$ (all $n \times n$ matrices) with the commutator bracket:

$$[\mathbf{A}, \mathbf{B}] = \mathbf{AB} - \mathbf{BA}.$$

The General Linear Group $GL(n, \mathbb{F})$

Example (the general linear group)

$$GL(n, \mathbb{F}) = \{\mathbf{A} \in \mathbb{F}^{n,n} : \det(\mathbf{A}) \neq 0\}.$$

- $GL(n, \mathbb{F})$ is an open subset of $\mathbb{F}^{n,n} \cong \mathbb{F}^{n^2}$, hence a smooth manifold of dimension n^2 .
- Its Lie algebra is $\mathfrak{gl}(n, \mathbb{F}) = \mathbb{F}^{n,n}$ (all $n \times n$ matrices) with the commutator bracket:

$$[\mathbf{A}, \mathbf{B}] = \mathbf{AB} - \mathbf{BA}.$$

- Most matrix Lie groups arise as closed subgroups of $GL(n, \mathbb{F})$. Their Lie algebras are subspaces of $\mathfrak{gl}(n, \mathbb{F})$ closed under the commutator.

The Special Orthogonal Group $SO(3)$

Example (the special orthogonal group — rigid rotations)

$$SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} : \mathbf{R}^T \mathbf{R} = \mathbf{I}_3, \det(\mathbf{R}) = 1 \};$$

$$\mathfrak{so}(3) = T_{\mathbf{I}_3} SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} : \mathbf{R}^T = -\mathbf{R} \}.$$

$\dim(\mathfrak{so}(3)) = \dim(SO(3)) = 3$, with basis $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$:

$$\mathbf{L}_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \mathbf{L}_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \quad \mathbf{L}_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The Special Orthogonal Group $SO(3)$

Example (the special orthogonal group — rigid rotations)

$$SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} : \mathbf{R}^T \mathbf{R} = \mathbf{I}_3, \det(\mathbf{R}) = 1 \};$$

$$\mathfrak{so}(3) = T_{\mathbf{I}_3} SO(3) = \{ \mathbf{R} \in \mathbb{R}^{3,3} : \mathbf{R}^T = -\mathbf{R} \}.$$

$\dim(\mathfrak{so}(3)) = \dim(SO(3)) = 3$, with basis $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$:

$$\mathbf{L}_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \mathbf{L}_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \quad \mathbf{L}_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Remark. $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ — the same structure as the cross product on $\mathbb{R}^3$.

The Special Unitary Group $SU(2)$

Example (the special unitary group)

$$SU(2) = \left\{ \mathbf{U} \in \mathbb{C}^{2,2} : \mathbf{U}^\dagger \mathbf{U} = \mathbf{I}_2, \det(\mathbf{U}) = 1 \right\};$$

$$\mathfrak{su}(2) = T_{\mathbf{I}_2} SU(2) = \left\{ \mathbf{U} \in \mathbb{C}^{2,2} : \mathbf{U}^\dagger = -\mathbf{U}, \operatorname{tr}(\mathbf{U}) = 0 \right\}.$$

$\dim(\mathfrak{su}(2)) = \dim(SU(2)) = 3$, with basis $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$:

$$\mathbf{T}_1 = \frac{1}{2} \begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix}, \quad \mathbf{T}_2 = \frac{1}{2} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \mathbf{T}_3 = \frac{1}{2} \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}.$$

The Special Unitary Group $SU(2)$

Example (the special unitary group)

$$SU(2) = \left\{ \mathbf{U} \in \mathbb{C}^{2,2} : \mathbf{U}^\dagger \mathbf{U} = \mathbf{I}_2, \det(\mathbf{U}) = 1 \right\};$$

$$\mathfrak{su}(2) = T_{\mathbf{I}_2} SU(2) = \left\{ \mathbf{U} \in \mathbb{C}^{2,2} : \mathbf{U}^\dagger = -\mathbf{U}, \operatorname{tr}(\mathbf{U}) = 0 \right\}.$$

$\dim(\mathfrak{su}(2)) = \dim(SU(2)) = 3$, with basis $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$:

$$\mathbf{T}_1 = \frac{1}{2} \begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix}, \quad \mathbf{T}_2 = \frac{1}{2} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \mathbf{T}_3 = \frac{1}{2} \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}.$$

Remark. $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$ — **identical** to $\mathfrak{so}(3)$. This is the first hint of the isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$.

The Double Cover $\Phi : \mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$

Construction

For $\mathbf{U} \in \mathrm{SU}(2)$, define $\Phi(\mathbf{U}) : \mathfrak{su}(2) \rightarrow \mathfrak{su}(2)$ by

$$[\Phi(\mathbf{U})](\mathbf{X}) \stackrel{\mathrm{df}}{=} \mathbf{U}\mathbf{X}\mathbf{U}^\dagger.$$

Using the basis $\beta = (\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$, identify $\mathfrak{su}(2) \cong \mathbb{R}^3$. Then the matrix $[\Phi(\mathbf{U})]_\beta^\beta$ lies in $\mathrm{SO}(3)$.

The Double Cover $\Phi : \text{SU}(2) \rightarrow \text{SO}(3)$

Construction

For $\mathbf{U} \in \text{SU}(2)$, define $\Phi(\mathbf{U}) : \mathfrak{su}(2) \rightarrow \mathfrak{su}(2)$ by

$$[\Phi(\mathbf{U})](\mathbf{X}) \stackrel{\text{df}}{=} \mathbf{U}\mathbf{X}\mathbf{U}^\dagger.$$

Using the basis $\beta = (\mathbf{T}_1, \mathbf{T}_2, \mathbf{T}_3)$, identify $\mathfrak{su}(2) \cong \mathbb{R}^3$. Then the matrix $[\Phi(\mathbf{U})]^\beta_\beta$ lies in $\text{SO}(3)$.

Theorem (double cover $\text{SU}(2) \rightarrow \text{SO}(3)$)



$\Phi : \text{SU}(2) \rightarrow \text{SO}(3)$ is:

- 1 A smooth group homomorphism: $\Phi(\mathbf{UV}) = \Phi(\mathbf{U}) \circ \Phi(\mathbf{V})$.
- 2 Surjective.
- 3 Has kernel $\ker(\Phi) = \{\pm \mathbf{I}_2\}$ (two elements).


Thus $\text{SU}(2)$ is a **double cover** of $\text{SO}(3)$.

The Double Cover: Geometric Picture


$$\mathrm{SU}(2) \cong S^3$$



$$\mathrm{SO}(3) \cong \mathbb{R}P^3$$



The Double Cover: Geometric Picture



Why two-to-one? Both $\mathbf{U}$ and $-\mathbf{U}$ give the same rotation. A rotation by angle θ about axis $\hat{\mathbf{n}}$ lifts to $\pm \cos(\theta/2)\mathbf{I}_2 \mp i \sin(\theta/2)(\hat{\mathbf{n}} \cdot \boldsymbol{\sigma})$. Increasing θ by 2π flips the sign.

Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$

Theorem (Lie algebra isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$)

Define $\varphi : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2)$ by $\varphi(\mathbf{L}_i) \stackrel{\text{df}}{=} \mathbf{T}_i$ (and extend $\mathbb{R}$ -linearly). Then φ is a Lie algebra isomorphism.

Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$

Theorem (Lie algebra isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$)

Define $\varphi : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2)$ by $\varphi(\mathbf{L}_i) \stackrel{\text{df}}{=} \mathbf{T}_i$ (and extend $\mathbb{R}$ -linearly). Then φ is a Lie algebra isomorphism.

Proof (essence)

- φ maps a basis to a basis $\implies$ linear isomorphism.
- Both satisfy $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ and $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$.
- Therefore $\varphi([\mathbf{L}_i, \mathbf{L}_j]) = \varphi(\varepsilon_{ijk} \mathbf{L}_k) = \varepsilon_{ijk} \mathbf{T}_k = [\mathbf{T}_i, \mathbf{T}_j] = [\varphi(\mathbf{L}_i), \varphi(\mathbf{L}_j)]$.

Lie Algebra Isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$

Theorem (Lie algebra isomorphism $\mathfrak{so}(3) \cong \mathfrak{su}(2)$)

Define $\varphi : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2)$ by $\varphi(\mathbf{L}_i) \stackrel{\text{df}}{=} \mathbf{T}_i$ (and extend $\mathbb{R}$ -linearly). Then φ is a Lie algebra isomorphism.

Proof (essence)

- φ maps a basis to a basis $\implies$ linear isomorphism.
- Both satisfy $[\mathbf{L}_i, \mathbf{L}_j] = \varepsilon_{ijk} \mathbf{L}_k$ and $[\mathbf{T}_i, \mathbf{T}_j] = \varepsilon_{ijk} \mathbf{T}_k$.
- Therefore $\varphi([\mathbf{L}_i, \mathbf{L}_j]) = \varphi(\varepsilon_{ijk} \mathbf{L}_k) = \varepsilon_{ijk} \mathbf{T}_k = [\mathbf{T}_i, \mathbf{T}_j] = [\varphi(\mathbf{L}_i), \varphi(\mathbf{L}_j)]$.

Significance. Although $SU(2)$ and $SO(3)$ are different as groups (double cover vs. quotient), their Lie algebras are **identical**. Infinitesimal rotations are the same in both pictures.

What We Have Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds**
— the geometric foundation.

What We Have Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds**
— the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.

What We Have Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- ③ **Lie groups** = smooth manifolds + groups — where geometry meets algebra.

What We Have Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- ③ **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- ④ Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).

What We Have Built So Far

- ① **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- ② **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- ③ **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- ④ Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).
- ⑤ The **Closed Subgroup Theorem** lets us recognise matrix Lie groups easily.

What We Have Built So Far

- 1 **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- 2 **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- 3 **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- 4 Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).
- 5 The **Closed Subgroup Theorem** lets us recognise matrix Lie groups easily.
- 6 $SO(3)$ (rotations) and $SU(2)$ (unitary 2×2 matrices) are the central examples.

What We Have Built So Far

- 1 **Topological spaces** $\rightarrow$ **topological manifolds** $\rightarrow$ **smooth manifolds** — the geometric foundation.
- 2 **Groups, fields, vector spaces, Lie algebras** — the algebraic foundation.
- 3 **Lie groups** = smooth manifolds + groups — where geometry meets algebra.
- 4 Every Lie group has a **Lie algebra** (tangent space at identity with commutator bracket).
- 5 The **Closed Subgroup Theorem** lets us recognise matrix Lie groups easily.
- 6 $SO(3)$ (rotations) and $SU(2)$ (unitary 2×2 matrices) are the central examples.
- 7 $SU(2)$ is a **double cover** of $SO(3)$. Their Lie algebras are **isomorphic**: $\mathfrak{so}(3) \cong \mathfrak{su}(2)$.

What Comes Next (Part II)

Representation theory (next meeting)

- Representations of Lie algebras and Lie groups.
- Complexification $\mathfrak{su}(2)_{\mathbb{C}} \cong \mathfrak{sl}(2, \mathbb{C})$ and the Cartan–Weyl basis (H, E, F) .
- Classification of irreducible representations of $\mathfrak{su}(2)$: for each $j \in \{0, \frac{1}{2}, 1, \frac{3}{2}, \dots\}$, a unique irrep of dimension $2j + 1$.
- Integration: every $\mathfrak{su}(2)$ -representation exponentiates to an $SU(2)$ -representation.
- Projective representations and the descent to $SO(3)$: integer $j \rightarrow$ ordinary representations, half-integer $j \rightarrow$ spinor (projective) representations.
- Physical spin operators and the 4π periodicity of spinors.

- Hall, B. C. (2015). *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction* (2nd ed.). Springer.
- Lee, J. M. (2012). *Introduction to Smooth Manifolds* (2nd ed.). Springer.
- Steinacker, H. (2024). *Lie Groups and Lie Algebras for Physicists*. Lecture notes, University of Vienna.
- Full paper with proofs: github.com/ganddoffcha/quantum-spin

Thank you.

Questions?