

Product Glossary



	Product	Active Ingredient	Indication
Sl. No.	Cardiology		
1	STATIX®	Atorvastatin 10/20/40/80 mg tablets	Controls elevated cholesterol levels
2	STATIX® F	Atorvastatin 10 mg + Micronised Fenofibrate 200 mg tablets	For Diabetic Dyslipidemia
3	STATIX® - EZ	Atorvastatin 10 mg + Ezetimibe 10 mg tablets	Controls extremely high levels of cholesterol
4	TELMISAT®	Telmisartan 20/40/80 mg tablets	Offers 24 hour blood pressure control
5	TELMISAT®-H	Telmisartan 40/80 mg + Hydrochlorothiazide 12.5 mg tablets	In uncontrolled Hypertension
6	TELMISAT® AM	Telmisartan 40 mg + Amlodipine 5 mg tablets	In Diabetic Hypertensives
7	ACTIBLOK - IPR	Metoprolol Immediate & Patterned Release 25/50/100 mg tablets	In patients of Hypertension, Angina, IHD and Heart Failure
8	ACTIBLOK AM®	Metoprolol Succinate IPR 25/50 mg +Amlodipine 5 mg tablets	In Uncontrolled Hypertension
9	BESTOR®	Rosuvastatin 5/10/20 mg tablets	For the management of Dyslipidemia and Atherosclerosis
10	BESTOR® FN	Rosuvastatin 10mg + Nanonized Fenofibrate 145 mg	In diabetes patients with combined hyperlipidmia
11	ZARGO®	Losartan Potassium 25/50 mg tablets	In hypertensive patients with CV co-morbidity
12	ZARGO® - H	Losartan Potassium 50 mg + Hydrochlorothiazide 12.5 mg tablets	In severe Hypertensive patients
13	MYOKINASE®	Recombinant Streptokinase for injection 1,500,000 IU	In patients of Acute Myocardial Infarction
14	DYNALIX®	Enoxaparin 40/60 mg Pre Filled Syringe	In patients of Acute Coronary Syndrome and Prophylaxis of Deep Vein Thrombosis
15	CLOTIDE®	Eptifibatide 10 ml bolus/100 ml for infusion	In ACS patients, undergoing Percutaneous Coronary Interventions
16	TIROZEST™	Tirozest HCL 5 mg/100 ml vial	In high risk ACS patients with NSTEMI

Sl. No.	Diabetology		
1	INSUPen®	INSUPen is a reusable insulin pen that can deliver insulin doses from 1 unit (0.01mL) to 60 units (0.60 mL) in increments of 1 unit.	INSUPen® is to be used only with Biocon's Insugen-R Refill™, Insugen-N Refill™, Insugen-30/70 Refill™, Insugen 50/50 Refill™ and BASALOG Refill™, refilled 3mL insulin cartridges
2	Insugen-30/70 Refill™	Each ml contains Human Insulin (rDNA origin), 100 IU.	To reduce blood glucose levels in all types of diabetes.
3	Insugen-50/50 Refill™	Each ml contains Human Insulin (rDNA origin), 100 IU.	To reduce blood glucose levels in all types of diabetes.
4	Insugen-N Refill™	Each ml contains Human Insulin (rDNA origin), 100 IU.	To reduce blood glucose levels in all types of diabetes.
5	Insugen-R Refill™	Each ml contains Human Insulin (rDNA origin), 100 IU.	To reduce blood glucose levels in all types of diabetes.
6	BASALOG Refill™	Each ml contains Insulin Glargine (rDNA Origin), 100 IU.	To provide 24 hours basal insulin action in all types of diabetes
7	INSUGEN® 30/70 40 IU	Each ml contains Human Insulin (rDNA origin), IP 40 IU	To reduce blood glucose levels in all types of diabetes
8	INSUGEN® 50/50 40 IU	Each ml contains Human Insulin (rDNA origin), IP 40 IU	To reduce blood glucose levels in all types of diabetes
9	INSUGEN® N 40 IU	Each ml contains Human Insulin (rDNA origin), IP 40 IU	To reduce blood glucose levels in all types of diabetes
10	INSUGEN® R 40 IU	Each ml contains Human Insulin (rDNA origin), IP 40 IU	To reduce blood glucose levels in all types of diabetes
11	Insugen® 30/70 100 IU	Each ml contains Human Insulin (rDNA origin), IP 100 IU	To reduce blood glucose levels in all types of diabetes
12	Insugen® 50/50 100 IU	Each ml contains Human Insulin (rDNA origin), IP 100 IU	To reduce blood glucose levels in all types of diabetes
13	Insugen® N 100 IU	Each ml contains Human Insulin (rDNA origin), IP 100 IU	To reduce blood glucose levels in all types of diabetes
14	Insugen® R 100 IU	Each ml contains Human Insulin (rDNA origin), IP 100 IU	To reduce blood glucose levels in all types of diabetes
15	BASALOG®	Each ml contains Insulin Glargine (rDNA Origin) 100 IU	To provide 24 hours basal insulin action in all types of diabetes
16	BLISTO®	Glimepiride 1/2/4 mg	For monotherapy or combination therapy of type 2 diabetes
17	BLISTO MF®	Glimepiride 1 mg + Metformin 500 mg SR & Glimepiride 2/4 mg + Metformin 1000 mg SR	For type 2 diabetes inadequately controlled with a single drug
18	METADOZE-IPR®	Metformin 500/850 mg IPR	For monotherapy or combination therapy of type 2 diabetes
19	TriGPM®	Glimepiride 1/2 mg + Pioglitazone 15 mg + Metformin 500 mg ER	For type 2 diabetes inadequately controlled with a combination therapy of 2 drugs
20	ZUKER®-MF	Gliclazide 80 mg + Metformin 500 mg SR	For type 2 diabetes inadequately controlled with a single drug
21	PIODART®	Pioglitazone 15/30 mg	For monotherapy or combination therapy of type 2 diabetes
22	PIODART®-MF	Pioglitazone 15 mg + Metformin ER 500 mg	For type 2 diabetes inadequately controlled with a single drug
23	OLISAT®	Orlistat 60/120 mg	For weight reduction in overweight or obese people
24	GABIL®	Gabapentin 300 mg + Methylcobalamin 500 mcg	To reduce symptoms of diabetic neuropathy
25	GMAB® Plus	GLA 100 mg + Methylcobalamin 1500 mcg + ALA 100 mg + Benfothiamine 100 mg + Elemental Zinc 15 mg	Nutritional supplementation in patients with diabetes
26	VOLICOSE®	Voglibose 0.2/0.3 mg	To reduce post-prandial plasma glucose levels in patients with diabetes
27	METADOSE® V	Voglibose 0.2/0.3 mg+ Metformin 500 mg For type 2 diabetes	For type 2 diabetes inadequately controlled with a single drug

Sl. No.	Nephrology		
1	ERYPRO safe™	Recombinant Human Erythropoietin Alpha injection in strengths of 2000/3000/4000/5000/6000/10000 IU	For the treatment of patients with anemia due to chronic renal failure, either on dialysis or not on dialysis
2	ERYPRO™ - Vial	Recombinant Human Erythropoietin Alpha 2000/4000/10000 IU	For the treatment of patients with anemia due to chronic renal failure, either on dialysis or not on dialysis
3	ERYPRO™ – PREFIL	Recombinant Human Erythropoietin Alpha 2000/4000 IU	For the treatment of patients with anemia due to chronic renal failure, either on dialysis or not on dialysis
4	TACROGRAF™	Tacrolimus 25/0.5/1/2 mg capsules	Prophylaxis of organ rejection in patients receiving allogenic liver, kidney or heart transplantation
5	RENODAPT®	Mycophenolate Mofetil 250 mg capsules and 500/750 mg tablets	Prophylaxis of organ rejection in patients receiving allogenic renal, heart or liver transplantation
6	RENODAPT®-S	Mycophenolic Acid 180/360 mg tablets	Prophylaxis of organ rejection in patients receiving allogenic renal transplantation
7	CYCLOPHIL ME® (ORAL SOLUTION)	Cyclosporine Oral Solution USP 100 mg/ml	Prophylaxis of organ rejection in kidney, liver and heart allogenic transplantation
8	CYCLOPHIL ME®	Cyclosporine USP 25/50/100 mg capsules	Prophylaxis of organ rejection in kidney, liver and heart allogenic transplantation
9	RAPACAN™	Sirolimus 1mg tablets	Prophylaxis of organ rejection in patients aged 13yrs or older receiving renal transplantation
10	Advacan™	Everolimus 0.25/0.5 mg tablets	Prophylaxis of organ rejection in adult patients at low to moderate immunologic risk receiving a kidney or cardiac transplantation
11	Narita™+	Whey protein supplement fortified with vitamins and minerals, 200 gm tin	Nutritional supplement in patients of chronic renal failure, including patients on dialysis
12	BIOSEV™	Sevelamer HCl 400/800 mg tablets	Control of serum phosphorous in patients with chronic kidney disease on dialysis
13	NEPORIN™	Heparin Sodium 5000/25000 IU	For adequate anticoagulation in haemodialysis

Sl. No.	Oncotherapeutics		
1	BIOMAb EGFR®	Nimotuzumab 200 mg humanized monoclonal antibody targeting epidermal growth factor receptor	For the treatment of locally advanced squamous cell carcinoma of head and neck, along with radiation and/or chemotherapy
2	Abraxane®	Paclitaxel protein-bound particles for injectable suspension (albumin-bound)	For the treatment of metastatic breast cancer
3	ERYPRO safe™	Recombinant Human Erythropoietin Alpha 40000 IU	
4	NUFIL safe™	Filgrastim (Recombinant Human Granulocyte Colony Stimulating Factor) 300 µg	For the treatment of chemotherapy induced neutropenia
5	NUFIL™	Filgrastim (Recombinant Human Granulocyte Colony Stimulating Factor) 300 µg	For the treatment of chemotherapy induced neutropenia
6	NUFIL Sf™	Filgrastim (Recombinant Human Granulocyte Colony Stimulating Factor) 300 µg	For the treatment of chemotherapy induced neutropenia
7	Genxtor®	Temsirolimus 25mg I.V	For the treatment of advanced renal cell carcinoma
8	Evertor™	Everolimus 5 mg/10 mg For the treatment of patients with advanced renal cell carcinoma	
9	XORTIB™	Capecitabine IP 150/500mg	<p>Colorectal Cancer</p> <ul style="list-style-type: none"> As a single agent for adjuvant treatment in patients with colorectal cancer Dukes' C Colon cancer who have undergone complete resection of the primary tumor For the treatment of metastatic <p>Breast Cancer</p> <ul style="list-style-type: none"> In combination with Docetaxel for the treatment of patients with locally advanced or metastatic breast cancer after failure of prior anthracycline containing chemotherapy <p>Gastric Cancer</p> <ul style="list-style-type: none"> For first-line treatment of advanced gastric cancer in combination with a platinum-based regimen
10	ATSURE®	Gemcitabine Hydrochloride - 200 mg/1 gm./ 1.4 gm.	<p>Breast Cancer</p> <ul style="list-style-type: none"> In combination with paclitaxel as first-line treatment of patients with metastatic breast cancer, after failure of prior anthracycline containing adjuvant chemotherapy, unless anthracyclines were clinically contraindicated. <p>Non-Small Cell Lung Cancer</p> <ul style="list-style-type: none"> In combination with Cisplatin for the first-line treatment of patients with inoperable locally advanced (Stage III A or IIIB) or metastatic (Stage IV) NSCLC. <p>Pancreatic Cancer</p> <ul style="list-style-type: none"> First-line treatment for patients with locally advanced (nonresectable Stage II or Stage III) or metastatic (Stage IV) adenocarcinoma of the pancreas. Also indicated for patients previously treated with 5-FU. <p>Ovarian Cancer</p> <ul style="list-style-type: none"> In combination with carboplatin for the treatment of patients with advanced ovarian cancer that has relapsed at least 6 months after completion of platinum-based therapy.
11	SYMXY™	Pemetrexed 100/500 mg	<p>Non-Small Cell Lung Cancer</p> <ul style="list-style-type: none"> In combination with Cisplatin for the initial treatment of patients with locally advanced or metastatic nonsquamous NSCLC. For the maintenance treatment of patients with locally advanced or metastatic nonsquamous NSCLC whose disease has not progressed after four cycles of platinum-based first-line chemotherapy. As a single-agent for the treatment of patients with locally advanced or metastatic nonsquamous NSCLC after prior chemotherapy. <p>Mesothelioma</p> <ul style="list-style-type: none"> In combination with Cisplatin for the treatment of patients with malignant pleural mesothelioma whose disease is unresectable or who are otherwise not candidates for curative surgery
12	HUMIX®	Lenalidomide – 5/10/25 mg	<p>Multiple Myeloma</p> <ul style="list-style-type: none"> In combination with dexamethasone for the treatment of patients with multiple myeloma who have received at least one prior therapy <p>Myelodysplastic Syndromes</p> <ul style="list-style-type: none"> For the treatment of patients with transfusion-dependent anaemia due to low or intermediate 1-risk Myelodysplastic syndromes associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities
13	ADNEXA®	Epirubicin Hydrochloride BP 10/50 mg	As a component of adjuvant therapy in patients with evidence of axillary node tumour involvement following resection of primary breast cancer.
14	XTIDE®	Oxaliplatin USP 50/100 mg	For the treatment of advanced colorectal cancer
15	TAZENTA™	Bortezomib 2mg	For the treatment of patients with multiple myeloma In mantle cell lymphoma who have received at least 1 prior therapy
16	XNTA™	Bendamustine Hydrochloride 100 mg	<p>Chronic Lymphocytic Leukaemia (CLL) For the treatment of patients with CLL</p> <p>Non-Hodgkin's Lymphoma For the treatment of patients with indolent B-cell non-Hodgkin's lymphoma that has progressed during or within six months of treatment with rituximab or a rituximab containing regimen</p>

Sl. No.	Immunotherapy		
1	ALZUMAb™	Each vial contains 25 mg of itolizumab. Itolizumab is a humanized recombinant anti-CD-6 monoclonal antibody. Each mL contains 5 mg of itolizumab for intravenous use.	A novel biologic for the treatment of moderate-to-severe psoriasis. ALZUMAb™ is indicated for the treatment of patients with active moderate to severe chronic plaque psoriasis who are candidates for systemic therapy.
2	PICON®	Pimecrolimus cream 1% w/w	Mild to Moderate atopic Dermatitis
3	TBIS®	Tacrolimus ointment 0.03%/0.1% w/w	Moderate to Severe Dermatitis
4	CALPSOR™	Calcipotriol ointment 0.005%w/w	Mild to Moderate Plaque Psoriasis
5	CALPSOR™C	Calcipotriol ointment 0.005%w/w + Clobetasol Propionate 0.05% w/w	Moderate to Severe Plaque Psoriasis
6	PSORID™	Cyclosporin capsules 25/50/100 mg Cyclosporin oral solution 100 mg/ml	Severe Psoriasis

Sl. No.	Comprehensive Care		
1	CASOGYN®	Casogyn 50 & 70	<ul style="list-style-type: none"> • Treatment of invasive candidiasis. • Treatment of invasive aspergillosis who are refractory to other antifungals • Empirical therapy for fungal infections (such as Candida or Aspergillus) in febrile neutropenia patients
2	NUCYTIN NDD	<ul style="list-style-type: none"> • 1g of Meropenem • 100ml of 0.9% NaCl Sodium Chloride for Intravenous Infusion • IV Set and Connector. 	<ul style="list-style-type: none"> • Skin and Skin Structure Infections • Intra-abdominal Infections • Bacterial Meningitis
3	MICELUM NDD	<ul style="list-style-type: none"> • 1 vial of Micelum NDD containing 500mg of Imipenem and 500mg of Cilastatin • 100ml of 0.9% NaCl Sodium Chloride Intravenous Infusion • IV Set and Connector. 	<ul style="list-style-type: none"> • Lower respiratory tract infections. • Urinary tract infections • Intra-abdominal infections. • Gynaecologic infections. • Bacterial septicemia. • Bone and joint infections. • Skin and skin structure infections. • Endocarditis.
4	PROLOP	Each 100mL contains Paracetamol IP-----1.0% w/v Mannitol IP-----5.0%w/v Water for injections IP---q.s.	<ul style="list-style-type: none"> • Management of mild to moderate pain • Management of moderate to severe pain with adjunctive opioid analgesics • Reduction of fever
5	FERVAY	FERVAY-250™ & 500™	<ul style="list-style-type: none"> • Pharyngitis/tonsillitis, • Acute bacterial otitis media, • Acute bacterial maxillary sinusitis, • Acute bacterial exacerbations of chronic bronchitis and • Secondary bacterial infections of acute bronchitis, • Uncomplicated skin and skin-structure infections, • Uncomplicated urinary tract infections, • Uncomplicated gonorrhea, • Early Lyme disease
6	FERVAY	FERVAY 1.5 g & FERVAY 750 mg	<ul style="list-style-type: none"> • Lower respiratory tract infections, • Urinary tract infections, • Skin and soft tissue infections, • Bone and joint infections, • Septicaemia, • Meningitis, • Gonorrhea • Prophylaxis against post-operative infection
7	KOOLISTIN	KOOLISTIN®1MIU & 2MIU	<ul style="list-style-type: none"> • Serious infections caused by Gram-negative bacteria, including those of the lower respiratory tract and urinary tract, where commonly used antibacterial agent may be ineffective because of bacterial resistance.
8	OGMIOS	OGMIOS – 1000MG+200MG INJ	<ul style="list-style-type: none"> • Upper Respiratory Tract Infections (including ENT): tonsillitis, sinusitis, otitis media • Lower Respiratory Tract Infections: acute exacerbations of chronic bronchitis, lobar- and broncho-pneumonia • Genito-urinary Tract Infections: cystitis, urethritis, pyelonephritis, female genital infections • Skin and Soft Tissue Infections • Bone and Joint Infections: osteomyelitis • Other Infections: septic abortion, puerperal sepsis, intra-abdominal sepsis, septicaemia, peritonitis, post-surgical infections • Prophylaxis against infection, which may be associated with major surgical procedures such as gastrointestinal, pelvic, head and neck, cardiac, renal, joint replacement and biliary tract surgery
9	OGMIOS	OGMIOS – 500MG+125MG TABS	<ul style="list-style-type: none"> • Acute bacterial sinusitis (adequately diagnosed), • Acute otitis media, • Acute exacerbations of chronic bronchitis (adequately diagnosed), • Community acquired pneumonia, • Cystitis, • Pyelonephritis, • Skin and soft tissue infections in particular cellulitis, animal bites, severe dental abscess with spreading cellulitis, • Bone and joint infections in particular osteomyelitis
10	DAPTOBIO™	DAPTOMYCIN 350MG Each vial contains: Daptomycin 350 mg Sodium Hydroxide (for pH adjustment) IP q.s	<ul style="list-style-type: none"> • Complicated skin and skin structure infections and • Staphylococcus aureus bloodstream infections (bacteremia), including those with right-sided infective endocarditis
11	VONFUNG NDD	<ul style="list-style-type: none"> • 1 vial of Vonfung NDD containing 200mg of voriconazole • 100ml of 0.9% NaCl Sodium Chloride Intravenous Infusion • IV Set and Connector 	<ul style="list-style-type: none"> • Invasive aspergillosis • Candidemia (nonneutropenics) and disseminated candidiasis in skin, abdomen, kidney, bladder wall, and wounds • Esophageal candidiasis
12	DORIPLUG	Doripenem monohydrate equivalent to 500 mg doripenem	<ul style="list-style-type: none"> • Nosocomial pneumonia, including ventilator-associated pneumonia (VAP) • Complicated intra-abdominal infections • Complicated urinary tract infections (UTI), including pyelonephritis and concurrent bacteremia
13	TIGEPLUG	Tigecycline 50mg Excipients q.s.	<ul style="list-style-type: none"> • Complicated skin and soft tissue infections • Complicated intra-abdominal infections
14	CYTO SORB	Cytosorb is an extracorporeal cytokine filter, which helps in removal of excess cytokines causing Multi Organ Failure in Sepsis patient. This device is developed by CYTOSORBENTS CORPORATION –USA, has received CE approval for usage in European countries for sepsis management. Studies done in European countries (European sepsis study) have established the safety of this device in sepsis patient and also showed significant reduction in Cytokines	In the management of Sepsis
15	THINWES	Each 10 ml contains: Terlipressin (as acetate) 1mg	<ul style="list-style-type: none"> • Bleeding oesophageal Varices. • Bleeding in gastrointestinal and urogenital systems such as gastric and duodenal ulcerations, functional and similar types of metrorrhagia, etc. • Bleeding as a result of surgery particularly of the abdominal and pelvic area. • Local application in gynaecological operations, such as cervix porta uteri.
16	HEPDOZE	Tenofovir disoproxil fumarate IP 300mg	<ul style="list-style-type: none"> • Chronic hepatitis B in adults and pediatric patients 12 years and older. • In combination with other antiretroviral agents, for the treatment of HIV-1 infection in adults and pediatric patients 2 years and older.
17	VARIOCT	<ul style="list-style-type: none"> • VARIOCT™ 50: Each mL contains Octreotide Acetate equivalent to Octreotide 50 mcg. • VARIOCT™ 100: Each mL contains Octreotide Acetate equivalent to Octreotide 100 mcg. 	<ol style="list-style-type: none"> 1. Acromegaly 2. Carcinoid tumors 3. Vasoactive intestinal peptide tumors
18	STERONEX	STERONEX 125/500/1000 mg INJECTIONS	<ol style="list-style-type: none"> 1. Neurological conditions 2. Collagen diseases 3. Dermatologic diseases 4. Severe, intractable allergic states 5. Rheumatic disorders 6. Inflammatory bowel diseases 7. Respiratory disorders 8. Hematologic disorders 9. Neoplastic disorders 10. Renal disease 11. Other conditions
19	CELRIM®	Cefepime 1g	<ul style="list-style-type: none"> • Pneumonia • Febrile Neutropenia • Urinary Tract Infections • Uncomplicated Skin and Skin Structure Infections • Complicated Intra-abdominal Infections
20	CELRIM TZ®	Cefepime 1g & Tazo bactam 0.125 g	<ul style="list-style-type: none"> • Urinary tract infections • Skin and skin structure infections and • Complicated intra-abdominal infections
21	Biopiper TZ™	Piperacillin 4 g & Tazobactam 0.5 g	<ul style="list-style-type: none"> • Intra-abdominal Infections • Skin and skin structure infections • Postpartum endometritis or pelvic inflammatory disease • Community-acquired pneumonia
22	Biopiper TZ™ 2.25	Piperacillin 2g and tazobactam 250mg	<ul style="list-style-type: none"> • Intra- abdominal Infections • Skin and skin structure infections • Postpartum endometritis or pelvic inflammatory disease • Community-acquired pneumonia

23	PENMER® – 500	Meropenem 500 mg	<ul style="list-style-type: none"> • Skin and Skin Structure Infections • Intra-abdominal Infections • Bacterial Meningitis
24	PENMER®	Meropenem 1g	<ul style="list-style-type: none"> • Skin and Skin Structure Infections • Intra-abdominal Infections • Bacterial Meningitis
25	IMICELUM™	Imipenem 500 mg & Cilastatin 500 mg	<ul style="list-style-type: none"> • Lower respiratory tract infections • Urinary tract infections • Intra-abdominal infections • Gynaecologic infections • Bacterial septicemia • Bone and joint infections • Skin and skin structure infections • Endocarditis
26	ENTAVAR™	Linezolid I.V 200 mg/300 ml	<ul style="list-style-type: none"> In Gram-Positive infections: • VRE infections • Nosocomial pneumonia • Skin and skin structure infections • Diabetic foot infections
27	ENTAVAR™ – 600	Linezolid Tablet 600 mg	<ul style="list-style-type: none"> In Gram-Positive infections: • VRE infections • Nosocomial pneumonia • Skin and skin structure infections • Diabetic foot infections
28	SUPRAVA TZ™	Ceftriaxone 1 g + Tazobactam 125mg	<ul style="list-style-type: none"> • Lower respiratory tract infections • community-acquired pneumonia • Acute bacterial otitis media • Skin and skin structure infections • Urinary tract infections • Uncomplicated gonorrhoea
29	SUPRAVA™	Ceftriaxone 500mg / 1g	<ul style="list-style-type: none"> • Lower respiratory tract infections • Acute bacterial otitis media • Skin and skin structure infections • Urinary tract infections • Bone and joint infections • Intra-abdominal infections • Meningitis • Surgical prophylaxis
30	CEGAVA™	1. Cefoperazone 1g and Sulbactam 500mg 2. Cefoperazone 500mg and Sulbactam 500mg 3. Cefoperazone 500mg and Sulbactam 500mg	<ul style="list-style-type: none"> • Respiratory Tract Infections • Urinary Tract Infections • Intra-abdominal infections • Skin and Soft Tissue Infections • Bone and Joint Infections • Pelvic Inflammatory Disease

31	CEGAVA TZ™	Cefoperazone 1g and Tazobactam 125mg	<ul style="list-style-type: none"> • Respiratory Tract Infections • Urinary Tract Infections • Intra-abdominal infections • Intra-abdominal infections • Skin and Soft Tissue Infections • Bone and Joint Infections
32	TECONIN™	Teicoplanin 200/400mg	Gram positive infections in the following: <ul style="list-style-type: none"> • skin and soft tissue infections • urinary tract infections • lower respiratory tract infections • joint and bone infections • septicaemia • endocarditis • peritonitis
33	THINLA™	Heparin 5000 IU / 25000IU	<ul style="list-style-type: none"> • Dialysis and continuous venous hemofiltration • As prophylaxis in ACS, PCI, DVT
34	MATABAC™	Sulbactam 1g / 2g	Acinetobacter induced: <ul style="list-style-type: none"> • Nosocomial Pneumonia • Bacteremia • UTI • Surgical Site wound infections • Post neurological meningitis

Sl. No.	Bioproducts		
1	IVNEX™ Safe	Human Normal Immunoglobulin for Intravenous Administration Ph. Eur. Sg	<ul style="list-style-type: none"> • Primary Immunodeficiency Diseases • Guillain Barre Syndrome • Idiopathic Thrombocytopenic Purpura • Kawasaki Disease
2	ZOMATOR™	Somatostatin 250 mcg and 3 mg	<ul style="list-style-type: none"> • Bleeding oesophageal Varices • Non variceal upper GI bleeding • Treatment of postoperative complications and prophylaxis following pancreatic surgery • Treatment of intestinal, biliary and pancreatic fistulae