

The synaptic dynamics of learning and memory

Subhaneil Lahiri


Stanford University, Applied Physics

May 13, 2019


“A memory frontier for complex synapses”, S Lahiri and S Ganguli.
Adv. Neural Inf. Process. Syst. 26, pp. 1034–1042., (2013).

What is a synapse?

Comp-neuro/deep-learning



Cellular biology




[Cajal (1899)]

What is a synapse?

Comp-neuro/deep-learning

Cellular biology

$$W_{ij}$$



[Klann (2002)]

Storage capacity of synaptic memory


Hopfield model, perceptron: capacity $\propto N$, (# synapses).

Assumes unbounded analogue synapses

With discrete, finite synapses:

\implies memory capacity $\sim \mathcal{O}(\log N)$.


[Amit and Fusi (1992), Amit and Fusi (1994)]




[Petersen et al. (1998), O'Connor et al. (2005)]

New memories overwrite old \implies stability-plasticity dilemma.

Synapses are complex




[Coba et al. (2009)]




[Montgomery and Madison (2002)]

Synapses are complex




[Coba et al. (2009)]



[Montgomery and Madison (2002)]


Cascade model




Capacity $\propto N^{2/3}$.

[Fusi et al. (2005)]

Synapses are complex




[Coba et al. (2009)]



[Montgomery and Madison (2002)]

Cascade model



Capacity $\propto N^{2/3}$. [Fusi et al. (2005)]

Capacity $\propto N$. [Benna and Fusi (2016)]

My approach

We want to study the structure-function relationship of biological processes.

Not trying to build a *single* model.

Instead, we build a broad framework of models to find:

- underlying mechanisms and principles.
- trade-offs between aspects of performance (e.g. learning vs. memory).
- properties of models that best manage these trade-offs.


Outline

1 Timescales of memory


- Quantifying memory quality
- Frontiers of memory
- Implications of memory limits

2 Designing experiments

Synaptic learning and memory



Synaptic learning and memory




Models of complex synaptic dynamics



Models of complex synaptic dynamics


- Internal functional state of synapse \rightarrow synaptic weight. ● weak
- Candidate plasticity events \rightarrow transitions between states ● strong



[Fusi et al. (2005), Fusi and Abbott (2007), Barrett and van Rossum (2008)]
[Smith et al. (2006), Lahiri and Ganguli (2013)]


Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Models of complex synaptic dynamics


- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong


Potential event



Depression event


Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Models of complex synaptic dynamics


- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics


- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong




Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong

Potential event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong

Potential event



Depression event


Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight. ● weak
- Candidate plasticity events \rightarrow transitions between states ● strong


- weak

- strong




Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).


Synaptic memory curves




Synaptic memory curves




Synaptic memory curves




Synaptic memory curves




Synaptic memory curves



Synaptic memory curves



Synaptic memory curves



Recognition memory: has this pattern been seen before?

Hypothesis test statistic: $\vec{w}(t) \cdot \vec{w}_{\text{test}}$.

[Sommer and Dayan (1998)]

Quantifying memory quality

Test if $\vec{w}(t) \cdot \vec{w}_{\text{test}} \geq \text{threshold}$?

[Sommer and Dayan (1998)]



Compare with null-distribution: $P[\vec{w}_{\text{null}} \cdot \vec{w}_{\text{test}}]$

$$\text{SNR}(t) = \frac{\langle \vec{w}(t) \cdot \vec{w}_{\text{test}} \rangle - \langle \vec{w}_{\text{null}} \cdot \vec{w}_{\text{test}} \rangle}{\sqrt{\text{Var}(\vec{w}_{\text{null}} \cdot \vec{w}_{\text{test}})}},$$

\Rightarrow discriminability: KL divergence, Chernoff distance, ROC curve, ...


Specific models of complex synaptic dynamics

Two example models of complex synapses.



[Fusi et al. (2005), Leibold and Kempster (2008), Ben-Dayan Rubin and Fusi (2007)]

These have different memory storage properties



Parameters for synaptic dynamics

$f^{\text{pot/dep}}$ = fraction of events that are pot/dep,

pot. event: $\mathbf{M}_{ij}^{\text{pot}}$ = transition prob. $i \rightarrow j$,

dep. event: $\mathbf{M}_{ij}^{\text{dep}}$ = transition prob. $i \rightarrow j$.

Constraints:

$$f^{\text{pot/dep}}, \mathbf{M}_{ij}^{\text{pot/dep}} \in [0, 1], \quad f^{\text{pot}} + f^{\text{dep}} = \sum_j \mathbf{M}_{ij}^{\text{pot/dep}} = 1.$$

Eigenmodes are more convenient parameters

Eigenmodes:

$$\left(f^{\text{pot}}\mathbf{M}^{\text{pot}} + f^{\text{dep}}\mathbf{M}^{\text{dep}}\right)\mathbf{u}_a = \lambda_a\mathbf{u}_a.$$

Contribution to SNR:


$$\text{SNR}(t) = \sum_a \mathcal{I}_a e^{-t/\tau_a}.$$

What are the constraints?

Upper bounds on measures of memory


Initial SNR:

$$\mathcal{I} = \text{SNR}(0) = \sum_a \mathcal{I}_a \leq \sqrt{N}.$$



Area under curve:


$$\mathcal{A} = \int_0^\infty \text{SNR}(t) dt = \sum_a \mathcal{I}_a \tau_a \leq \sqrt{N}(M-1)/r.$$



[Lahiri and Ganguli (2013)]


Proven envelope: memory frontier

Upper bound on memory curve at *any* timescale.





Proven envelope: memory frontier



Upper bound on memory curve at *any* timescale.





Models that maximize memory for one timescale



Models that maximize memory for one timescale




Models that maximize memory for one timescale



Synaptic diversity and timescales of memory

Different synapses have different molecular structures.




Emes RD, Grant SGN, 2012.
Annu. Rev. Neurosci. 35:111-31

[Emes and Grant (2012)]

Synaptic diversity and timescales of memory

Different synapses have different molecular structures.




Emes RD, Grant SGN. 2012.
Annu. Rev. Neurosci. 35:111–31

[Emes and Grant (2012)]

Memories stored in different places for different timescales

[Squire and Alvarez (1995)]

[McClelland et al. (1995)]



[Born and Wilhelm (2012)]

Also: Cerebellar cortex → nuclei.

[Attwell et al. (2002)]



[Cooke et al. (2004)]

Synaptic structure and function: general principles

Real synapses limited by molecular building blocks.
Evolution had larger set of priorities.

What can we conclude?

Short timescale \longrightarrow Moderate timescale \longrightarrow Long timescale





Synaptic structure and function: general principles

Real synapses limited by molecular building blocks.
Evolution had larger set of priorities.

What can we conclude?

Short timescale \longrightarrow Moderate timescale \longrightarrow Long timescale





short topology \longrightarrow long topology

Synaptic structure and function: general principles

Real synapses limited by molecular building blocks.
Evolution had larger set of priorities.

What can we conclude?

Short timescale \longrightarrow Moderate timescale \longrightarrow Long timescale



short topology \longrightarrow long topology

deterministic synapse \longrightarrow stochastic synapse

Experimental tests?

Traditional experiments:



Experimental tests?

Traditional experiments:



To fit a model: long sequence of small plasticity events.
Observe the changes in synaptic efficacy.



Experimental tests?

Traditional experiments:



To fit a model: long sequence of small plasticity events.
Observe the changes in synaptic efficacy.




EM algorithms:

Sequence of hidden states \rightarrow estimate transition probabilities


Transition probabilities \rightarrow estimate sequence of hidden states

[Baum et al. (1970), Rabiner and Juang (1993), Dempster et al. (2007)]


Simulated experiment




Estimated model



pot



Transition prob.



Transition prob.



dep

Problem: need *long* sequences.

Whole cell patch of postsynaptic neuron → Ca washout.

Conclusions

- We have formulated a general theory of learning and memory with complex synapses.
- We find a memory envelope: a single curve that cannot be exceeded by the memory curve of *any* synaptic model.
- We understood which types of synaptic structure are useful for storing memories for different timescales.
- We studied more than a single model. We studied *all possible models*, to extract general principles relating synaptic structure to function

Acknowledgements

Surya Ganguli

Jascha Sohl-Dickstein

Friedemann Zenke

Sam Ocko

Stephane Deny

Jonathan Kadmon

Madhu Advani

Peiran Gao

Niru Maheswaranathan

Ben Poole

Kiah Hardcastle

Lane McIntosh

Alex Williams

Christopher Stock

Sarah Harvey

Aran Nayebi

Jennifer Raymond

Barbara Nguyen-Vu

Grace Zhao

Aparna Suvrathan

Rhea Kimpo

Carla Shatz

Hanmi Lee


David Sussillo

Stefano Fusi


Marcus Benna

Funding: Swartz Foundation, Stanford Bio-X Genentech fellowship.

Model of circuit




Baseline



Evidence: level of depression

Basal level of GluR2 phosphorylation at serine 880 in AMPA receptor.






Biochemical signature of PF-Pk LTD.

Shows that # depressed synapses in flocculus is larger in KO than WT.


Evidence: saturation by CF stimulation

Use Channelrhodopsin to stimulate CF \rightarrow increase LTD in PF-Pk synapses
 \rightarrow simulate saturation in WT.






Other models that fail

Multistate synapse




Pooled resource model





[Amit and Fusi (1994)]

Other models that work

Non-uniform multistate model




Cascade model



[Fusi et al. (2005)]

Mathematical explanation



Serial synapse: $\pi_i \sim \mathcal{N} \left(\frac{q^{\text{pot}}}{q^{\text{dep}}} \right)^i$.

Learning rate $\sim \pi_{M/2} \left(\frac{q^{\text{dep}}}{q^{\text{pot}}} \right) = \mathcal{N} \left(\frac{q^{\text{pot}}}{q^{\text{dep}}} \right)^{\frac{M}{2}-1}$.

For $M > 2$: larger $q^{\text{dep}} \implies$ slower learning.


For $M = 2$: larger $q^{\text{dep}} \implies$ larger $\mathcal{N} \implies$ faster learning.

Initial SNR as flux

Initial SNR is closely related to flux between strong & weak states


$$\text{SNR}(0) \leq \frac{4\sqrt{N}}{r} \Phi_{-+}.$$

Max when potentiation guarantees $\mathbf{w} \rightarrow +1$,
depression guarantees $\mathbf{w} \rightarrow -1$.



Two-state model

Two-state model equivalent to previous slide:



$$\Rightarrow \text{SNR}(t) = \sqrt{N} (4f^{\text{pot}} f^{\text{dep}}) e^{-rt}.$$

Maximal initial SNR:


$$\text{SNR}(0) \leq \sqrt{N}.$$

Area under memory curve

$$\mathcal{A} = \int_0^\infty dt \text{ SNR}(t), \quad \overline{\text{SNR}}(\tau) \rightarrow \frac{\mathcal{A}}{\tau} \quad \text{as } \tau \rightarrow \infty.$$

Area bounds memory lifetime:

$$\begin{aligned} \text{SNR}(\text{lifetime}) &= 1 \\ \Rightarrow \quad \text{lifetime} &< \mathcal{A}. \end{aligned}$$



This area has an upper bound:


$$\mathcal{A} \leq \sqrt{N}(M-1)/r.$$

Saturated by a model with linear chain topology.

Proof of area bound

For any model, we can construct perturbations that

- preserve equilibrium distribution,
- increase area.



e.g. decrease “shortcut” transitions, increase bypassed “direct” ones.

Endpoint: linear chain

The area of this model is


$$A = \frac{2\sqrt{N}}{r} \sum_k \pi_k |k - \langle k \rangle|.$$

Max: equilibrium probability distribution concentrated at both ends.

[Barrett and van Rossum (2008)]

Saturating model

Make end states “sticky”



Has long decay time, but terrible initial SNR.

$$\lim_{\epsilon \rightarrow 0} A = \sqrt{N}(M-1)/r.$$

Technical detail: ordering states

Let \mathbf{T}_{ij} = mean first passage time from state i to state j . Then:

$$\eta = \sum_j \mathbf{T}_{ij} \pi_j,$$

is independent of the initial state i (Kemeney's constant).


[Kemeny and Snell (1960)]

We define:

$$\eta_i^+ = \sum_{j \in \text{strong}} \mathbf{T}_{ij} \pi_j, \quad \eta_i^- = \sum_{j \in \text{weak}} \mathbf{T}_{ij} \pi_j.$$

They can be used to arrange the states in an order (increasing η^- or decreasing η^+). [back](#)

Intuition for using topology



$$\begin{aligned}\mathcal{I} &\propto q, & \max_a \tau_a &\propto \frac{1}{q}, \\ \mathcal{I} &\propto \frac{1}{M}, & \max_a \tau_a &\propto M^2,\end{aligned}$$

\Rightarrow

$$\begin{aligned}\text{Stochasticity: } \mathcal{I} &\propto \frac{1}{\tau_{\max}}, \\ \text{Topology: } \mathcal{I} &\propto \frac{1}{\sqrt{\tau_{\max}}}.\end{aligned}$$

References VII



Christian Leibold and Richard Kempter.

“Sparseness Constrains the Prolongation of Memory Lifetime via Synaptic Metaplasticity”.

Cereb. Cortex, 18(1):67–77, (2008) .



Daniel D Ben-Dayan Rubin and Stefano Fusi.

“Long memory lifetimes require complex synapses and limited sparseness”.

Front. Comput. Neurosci., 1:1–14, (November, 2007) .



Richard D. Emes and Seth G.N. Grant.

“Evolution of Synapse Complexity and Diversity”.

Annual Review of Neuroscience, 35(1):111–131, (2012) .

47

References IX



Phillip J.E. Attwell, Samuel F. Cooke, and Christopher H. Yeo.

“Cerebellar Function in Consolidation of a Motor Memory”.

Neuron, 34(6):1011–1020, (jun, 2002) .



Samuel F Cooke, Phillip J E Attwell, and Christopher H Yeo.

“Temporal properties of cerebellar-dependent memory consolidation.”

J. Neurosci., 24(12):2934–41, (mar, 2004) .

46



LE Baum, T Petrie, George Soules, and Norman Weiss.

“A maximization technique occurring in the statistical analysis of probabilistic functions of Markov chains”.

The annals of mathematical statistics, 41(1):164–171, (1970) .

51

52

