

Learning and memory with complex synaptic plasticity

Subhaneil Lahiri

Stanford University, Applied Physics

March 29, 2019

What is a synapse?


Experimentalists

Theorists

What is a synapse?

Experimentalists

Theorists




[Klann (2002)]

What is a synapse?

Experimentalists

Theorists



[Klann (2002)]

W_{ij}

Storage capacity of synaptic memory


Hopfield, perceptron have capacity $\propto N$, (# synapses).

Assumes unbounded analogue synapses

With discrete, finite synapses:


\implies memory capacity $\sim \mathcal{O}(\log N)$.

[Amit and Fusi (1992), Amit and Fusi (1994)]




[Petersen et al. (1998), O'Connor et al. (2005)]

Synapses are complex



[Coba et al. (2009)]



[Montgomery and Madison (2002)]

Outline

1 Learning with enhanced plasticity

- Effects of enhanced plasticity on cerebellar learning
- Synaptic models of cerebellar learning
- Learning outcomes of mice and models

2 Memory over different timescales

- Quantifying memory quality
- Frontiers of memory
- Implications of memory limits

3 Designing experiments

Section 1

Learning with enhanced plasticity

Benefits of enhanced plasticity?

Learning requires synaptic plasticity.




Can we enhance learning by enhancing plasticity?





Enhanced plasticity *can* enhance learning


Overexpress NR2B



Inhibit CN



Knockout Hdac2



Fear conditioning

[Tang et al. (1999)]

Novel object recog.


[Malleret et al. (2001)]

Fear conditioning

[Guan et al. (2009)]

Enhanced plasticity can *impair* learning


Mutate PSD-95



Water maze

[Migaud et al. (1998)]


Knockout PTP δ



Water maze

[Uetani et al. (2000)]


Delete Tmod2



Fear cond.

[Cox et al. (2003)]

Knockout FMR1



Eyeblink

[Koekkoek et al. (2005)]

also: [Hayashi et al. (2004), Rutten et al. (2008)]

Overview

Sometimes enhanced plasticity → enhanced learning.
Sometimes enhanced plasticity → impaired learning.

Why? How? When?



Overview

Sometimes enhanced plasticity → enhanced learning.
Sometimes enhanced plasticity → impaired learning.




Why? How? When?

Mice with enhanced cerebellar plasticity can show **both** impaired and enhanced learning.

Simple synapses **cannot** explain behaviour. **Complex synapses** are required.
→ predictions for synaptic physiology.

Vestibulo-Occular Reflex



Eye movements compensate for head movements
⇒ stabilise image on retina.

Requires control of VOR gain = $\frac{\text{eye velocity}}{\text{head velocity}}$.


Needs to be adjusted as eye muscles age, etc.

Vestibulo-Occular Reflex training

VOR Increase
Training



VOR Decrease
Training




VOR increase: LTD in PF-Pk synapses.
VOR decrease: different mechanism,
also reverses LTD in PF-Pk.

[Marr (1969), Albus (1971), Ito (1972)]

Enhanced plasticity impairs learning

Expectation: enhanced LTD \rightarrow enhanced learning.

VOR Increase
Training




Experiment: enhanced plasticity \rightarrow impaired learning.

Knockout of MHC-I K^bD^b molecules in PF-Pk synapses
 \rightarrow lower threshold for LTD

[McConnell et al. (2009)]


Depletion hypothesis

Learning rate \sim intrinsic plasticity rate \times # synapses available for LTD.




Depletion hypothesis

Learning rate \sim intrinsic plasticity rate \times # synapses available for LTD.



Depletion hypothesis

Learning rate \sim intrinsic plasticity rate \times # synapses available for LTD.




Question 1: depletion effect competes with enhanced intrinsic plasticity.
When is depletion effect stronger?


Replenishment by reverse-training




Replenishment by reverse-training




Replenishment by reverse-training




Replenishment by reverse-training




Replenishment by reverse-training



Enh. Plast.



WT



Question 2: How can replenishment ever impair learning?


Models of complex synaptic dynamics



Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
- Candidate plasticity events → transitions between states

● weak
● strong



States: NMDAR subunit composition, CaMK II autophosphorylation, activating PKC, p38 MAPK,...


[Fusi et al. (2005), Fusi and Abbott (2007), Barrett and van Rossum (2008)]

[Smith et al. (2006); Lahiri and Ganguly (2013)]

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong

Potentiation event




Depression event

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Models of complex synaptic dynamics

- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong


Potentiation




Depression

Metaplasticity: change propensity for plasticity
(independent of change in synaptic weight).

Modelling VOR experiments




Modelling VOR experiments



Modelling VOR experiments


PF-Pk LTD → VOR increase



Training: different CF activity \implies
change frequency of pot/dep events.

Modelling VOR experiments


PF-Pk LTD → VOR increase



Training: different CF activity \implies
change frequency of pot/dep events.

Modelling VOR experiments


PF-Pk LTD → VOR increase



Training: different CF activity \Rightarrow
change frequency of pot/dep events.

Modelling VOR experiments


PF-Pk LTD → VOR increase



Training: different CF activity \Rightarrow
change frequency of pot/dep events.

Modelling VOR experiments


PF-Pk LTD → VOR increase



Training: different CF activity \implies
change frequency of pot/dep events.

Modelling VOR experiments

PF-Pk LTD → VOR increase




Training: different CF activity \implies
change frequency of pot/dep events.

Learning: decrease in average synaptic weight.

Modelling VOR experiments

PF-Pk LTD → VOR increase




Training: different CF activity \Rightarrow
change frequency of pot/dep events.

Learning: decrease in average synaptic weight.

Mutation: lower threshold for LTD \Rightarrow
increase transition probability for depression events.

Modelling VOR experiments

PF-Pk LTD → VOR increase



Training: different CF activity \Rightarrow
change frequency of pot/dep events.

Learning: decrease in average synaptic weight.

Mutation: lower threshold for LTD \Rightarrow
increase transition probability for depression events.

Questions

Depletion effect competes with enhanced intrinsic plasticity.

Question 1: When is the depletion effect stronger?

Reverse training impairs learning in wild-type.


Question 2: How can replenishment ever impair learning?

Enhanced plasticity → enhanced/impaired learning


Big question: Why? When?

Simple synapses cannot explain the data

Multistate synapse




VOR Increase
Training




Simple synapses cannot explain the data

Two-state model




VOR Increase
Training




Simple synapses cannot explain the data

Two-state model



VOR Increase
Training




Simple synapses cannot explain the data

Two-state model



Initial distribution




depletion effect

<
enhanced plasticity


⇒ enhanced learning

Simple synapses cannot explain the data

Two-state model



VOR Increase Training



Simple synapses cannot explain the data



Two-state model



Initial distribution

WT
w/ rev-training

synapses



reverse training





replenishment




enhanced learning

Complex metaplastic synapses can explain the data



Serial model




VOR Increase Training



Complex metaplastic synapses can explain the data





VOR Increase
Training





Complex metaplastic synapses can explain the data

Serial model





amplified depletion
>
enhanced plasticity
⇒ impaired learning

Complex metaplastic synapses can explain the data



Complex metaplastic synapses can explain the data




Serial model



reverse training
+
“stubborn” metaplasticity
⇒ impaired learning


Complex metaplastic synapses can explain the data

Serial model



Complex metaplastic synapses can explain the data

Serial model



starting point:
labile states




enhanced plasticity
⇒ impaired learning

starting point:
stubborn states



enhanced plasticity
⇒ enhanced learning


Essential features





The success of the serial model relies on two features:

- Complexity - needed for depletion to dominate enhanced plasticity,
- Stubbornness - repeated potentiation impairs subsequent depression.

Fail:



Succeed:



Conclusions

- Diverse behavioural patterns:
Enhanced plasticity → enhance/impair learning (prior experience).
Reverse-training → enhance/impair learning (plasticity rates).
- **enhanced LTD vs. depletion** → learning outcome.
- Predictions for synaptic physiology:
Complexity: necessary to amplify depletion.
Stubbornness: repeated potentiation impairs subsequent depression.
- We used behaviour to constrain the dynamics of synaptic plasticity.



Section 2

Memory over different timescales

Storage capacity of synaptic memory

Hopfield, perceptron have capacity $\propto N$, (# synapses).

Assumes unbounded analog synapses

With discrete, finite synapses:

\Rightarrow memory capacity $\sim \mathcal{O}(\log N)$.


[Amit and Fusi (1992), Amit and Fusi (1994)]

New memories overwrite old \Rightarrow stability-plasticity dilemma.

Models of complex synaptic dynamics

There are N identical synapses with M internal functional states.





- Internal functional state of synapse → synaptic weight.
 - Candidate plasticity events → transitions between states
- weak
● strong




States: #AMPAR, #NMDAR, NMDAR subunit composition,
CaMK II autophosphorylation, activating PKC, p38 MAPK,...

[Fusi et al. (2005), Fusi and Abbott (2007), Barrett and van Rossum (2008)]

Specific models of complex synaptic dynamics




Synaptic memory curves




Synapses store a sequence of memories.

Recognition memory performance described by SNR.




General principles relating structure and function?

Synaptic structure



Synaptic function



- What are the fundamental limits of memory?
- Which models achieve these limits?
- What are the theoretical principles behind the optimal models?

Parameters for synaptic dynamics

$f^{\text{pot/dep}}$ = fraction of events that are pot/dep,

pot. event: M_{ij}^{pot} = transition prob. $i \rightarrow j$,

$$\mathbf{W}^{\text{pot}} = f^{\text{pot}}(\mathbf{M}^{\text{pot}} - I),$$

dep. event: M_{ij}^{dep} = transition prob. $i \rightarrow j$,

$$\mathbf{W}^{\text{dep}} = f^{\text{dep}}(\mathbf{M}^{\text{dep}} - I).$$

Constraints:

$$f^{\text{pot/dep}}, M_{ij}^{\text{pot/dep}} \in [0, 1], \quad f^{\text{pot}} + f^{\text{dep}} = \sum_j M_{ij}^{\text{pot/dep}} = 1.$$


Memory curve given by

$$\begin{aligned}\overline{\text{SNR}}(\tau) &= \sqrt{N} \pi (\mathbf{W}^{\text{pot}} - \mathbf{W}^{\text{dep}}) \left[\mathbf{I} - r\tau (\mathbf{W}^{\text{pot}} + \mathbf{W}^{\text{dep}}) \right]^{-1} \mathbf{w}. \\ &= \sqrt{N} \sum_a \frac{\mathcal{I}_a}{1 + r\tau/\tau_a}.\end{aligned}$$

Upper bounds on measures of memory


Initial SNR:

$$\mathcal{I} = \text{SNR}(0) = \sum_a \mathcal{I}_a \leq \sqrt{N}.$$



Area under curve:


$$\mathcal{A} = \int_0^\infty \text{SNR}(t) dt = \sum_a \mathcal{I}_a \tau_a \leq \sqrt{N}(M-1)/r.$$



[Lahiri and Ganguli (2013)]


Proven envelope: memory frontier

Upper bound on memory curve at *any* timescale.





Proven envelope: memory frontier



Upper bound on memory curve at *any* timescale.





Models that maximize memory for one timescale



Models that maximize memory for one timescale




Models that maximize memory for one timescale



Synaptic diversity and timescales of memory

Different synapses have different molecular structures.




[A] Emes RD, Grant SGN. 2012.
Annu. Rev. Neurosci. 35:111–31

[Emes and Grant (2012)]

Synaptic diversity and timescales of memory

Different synapses have different molecular structures.




[Emes RD, Grant SGN. 2012.
Annu. Rev. Neurosci. 35:111–31]

[Emes and Grant (2012)]

Memories stored in different places for different timescales

[Squire and Alvarez (1995)]

[McClelland et al. (1995)]



[Born and Wilhelm (2012)]

Also: Cerebellar cortex → nuclei.


[Attwell et al. (2002)]

[Cooke et al. (2004)]

Synaptic structure and function: general principles

Real synapses limited by molecular building blocks.
Evolution had larger set of priorities.


What can we conclude?



Synaptic structure and function: general principles

Real synapses limited by molecular building blocks.
Evolution had larger set of priorities.


What can we conclude?



Synaptic structure and function: general principles

Real synapses limited by molecular building blocks.
Evolution had larger set of priorities.

What can we conclude?



Section 3

Designing experiments

Experimental tests?

Traditional experiments:



Experimental tests?

Traditional experiments:




To fit a model: long sequence of small plasticity events.

Observe the changes in synaptic efficacy.



Simulated experiment




EM algorithms:

Sequence of hidden states → estimate transition probabilities
Transition probabilities → estimate sequence of hidden states

[Baum et al. (1970), Rabiner and Juang (1993), Dempster et al. (2007)]

Simulated experiment




EM algorithms:

Sequence of hidden states → estimate transition probabilities
Transition probabilities → estimate sequence of hidden states

[Baum et al. (1970), Rabiner and Juang (1993), Dempster et al. (2007)]

Simulated experiment




EM algorithms:

Sequence of hidden states → estimate transition probabilities
Transition probabilities → estimate sequence of hidden states

[Baum et al. (1970), Rabiner and Juang (1993), Dempster et al. (2007)]

Simulated experiment




EM algorithms:

Sequence of hidden states → estimate transition probabilities
Transition probabilities → estimate sequence of hidden states

[Baum et al. (1970), Rabiner and Juang (1993), Dempster et al. (2007)]

Simulated experiment



EM algorithms:

Sequence of hidden states → estimate transition probabilities
Transition probabilities → estimate sequence of hidden states

[Baum et al. (1970), Rabiner and Juang (1993), Dempster et al. (2007)]

Experimental problems

- Need single synapses.
- Need long sequences of plasticity events.
- Need to control types of candidate plasticity events.
- Need to measure synaptic efficacies.

When we patch the postsynaptic neuron → Ca washout.

Summary

- We have formulated a general theory of learning and memory with complex synapses.
- We find a memory envelope: a single curve that cannot be exceeded by the memory curve of *any* synaptic model.
- We understood which types of synaptic structure are useful for storing memories for different timescales.
- We studied more than a single model. We studied *all possible models*, to extract general principles relating synaptic structure to function

Acknowledgements

Surya Ganguli

Jascha Sohl-Dickstein
Friedemann Zenke
Sam Ocko
Stephane Deny
Jonathan Kadmon
Madhu Advani
Peiran Gao

David Sussillo

Niru Maheswaranathan

Ben Poole
Kiah Hardcastle
Lane McIntosh
Alex Williams
Christopher Stock
Sarah Harvey
Aran Nayebi

Stefano Fusi

Jennifer Raymond

Barbara Nguyen-Vu
Grace Zhao

Aparna Suvrathan
Rhea Kimpo


Carla Shatz

Hanmi Lee


Marcus Benna

Funding: Swartz Foundation, Stanford Bio-X Genentech fellowship.

Model of circuit




Baseline



Evidence: level of depression

Basal level of GluR2 phosphorylation at serine 880 in AMPA receptor.






Biochemical signature of PF-Pk LTD.

Shows that # depressed synapses in flocculus is larger in KO than WT.


Evidence: saturation by CF stimulation

Use Channelrhodopsin to stimulate CF → increase LTD in PF-Pk synapses
→ simulate saturation in WT.






Other models that fail

Multistate synapse




Pooled resource model





[Amit and Fusi (1994)]

Other models that work

Non-uniform multistate model




Cascade model



[Fusi et al. (2005)]

Mathematical explanation



Serial synapse: $\pi_i \sim \mathcal{N} \left(\frac{q^{\text{pot}}}{q^{\text{dep}}} \right)^i.$


Learning rate $\sim \pi_{M/2} \left(\frac{q^{\text{dep}}}{q^{\text{pot}}} \right) = \mathcal{N} \left(\frac{q^{\text{pot}}}{q^{\text{dep}}} \right)^{\frac{M}{2}-1}.$

For $M > 2$: larger q^{dep} \implies slower learning.

For $M = 2$: larger q^{dep} \implies larger \mathcal{N} \implies faster learning.

Quantifying memory quality

Test if $\vec{w}_{\text{ideal}} \cdot \vec{w}(t) \geq \theta$?



$$\text{SNR}(t) = \frac{\langle \vec{w}_{\text{ideal}} \cdot \vec{w}(t) \rangle - \langle \vec{w}_{\text{ideal}} \cdot \vec{w}(\infty) \rangle}{\sqrt{\text{Var}(\vec{w}_{\text{ideal}} \cdot \vec{w}(\infty))}},$$

$$\overline{\text{SNR}}(\tau) = \int d\tau \frac{e^{-t/\tau}}{\tau} \text{SNR}(t).$$

$$\text{NNR}(t) = \sqrt{\frac{\text{Var}(\vec{w}_{\text{ideal}} \cdot \vec{w}(t))}{\text{Var}(\vec{w}_{\text{ideal}} \cdot \vec{w}(\infty))}}.$$


Also: KL divergence, Chernoff distance, . . .

Initial SNR as flux

Initial SNR is closely related to flux between strong & weak states


$$\text{SNR}(0) \leq \frac{4\sqrt{N}}{r} \Phi_{-+}.$$

Max when potentiation guarantees $\mathbf{w} \rightarrow +1$,
depression guarantees $\mathbf{w} \rightarrow -1$.



Two-state model

Two-state model equivalent to previous slide:



Maximal initial SNR:


$$\text{SNR}(0) \leq \sqrt{N}.$$

Area under memory curve

$$\mathcal{A} = \int_0^\infty dt \text{ SNR}(t), \quad \overline{\text{SNR}}(\tau) \rightarrow \frac{\mathcal{A}}{\tau} \quad \text{as} \quad \tau \rightarrow \infty.$$

Area bounds memory lifetime:

$$\begin{aligned} \text{SNR(lifetime)} &= 1 \\ \implies \text{lifetime} &< \mathcal{A}. \end{aligned}$$



This area has an upper bound:

$$\mathcal{A} \leq \sqrt{N(M-1)}/r.$$


Saturated by a model with linear chain topology.

Proof of area bound

For any model, we can construct perturbations that

- preserve equilibrium distribution,
- increase area.

details



e.g. decrease “shortcut” transitions, increase bypassed “direct” ones.
Endpoint: linear chain

The area of this model is


$$A = \frac{2\sqrt{N}}{r} \sum_k \pi_k |k - \langle k \rangle|.$$

Max: equilibrium probability distribution concentrated at both ends.

[Barrett and van Rossum (2008)]

Saturating model

Make end states “sticky”



Has long decay time, but terrible initial SNR.

$$\lim_{\varepsilon \rightarrow 0} A = \sqrt{N}(M - 1)/r.$$

Technical detail: ordering states

Let \mathbf{T}_{ij} = mean first passage time from state i to state j . Then:

$$\eta = \sum_j \mathbf{T}_{ij} \pi_j,$$

is independent of the initial state i (Kemeney's constant).

[Kemeny and Snell (1960)]


We define:

$$\eta_i^+ = \sum_{j \in \text{strong}} \mathbf{T}_{ij} \pi_j, \quad \eta_i^- = \sum_{j \in \text{weak}} \mathbf{T}_{ij} \pi_j.$$

They can be used to arrange the states in an order (increasing η^- or decreasing η^+). [back](#)

Technical detail: upper/lower triangular


With states in order:



Endpoint: potentiation goes right, depression goes left.

[back](#)

Intuition for using topology



$$\begin{array}{lll} \mathcal{I} \propto q, & \max_a \tau_a \propto \frac{1}{q}, & \xrightarrow{\quad} \text{Stochasticity: } \mathcal{I} \propto \frac{1}{\tau_{\max}}, \\ \mathcal{I} \propto \frac{1}{M}, & \max_a \tau_a \propto M^2, & \text{Topology: } \mathcal{I} \propto \frac{1}{\sqrt{\tau_{\max}}}. \end{array}$$

References I



Eric Klann.

“Metaplastic Protein Phosphatases”.

Learning and Memory, 9(4):153–155, (2002) ,

<http://learnmem.cshlp.org/content/9/4/153.full.pdf+html>.

2 3 4



D. J. Amit and S. Fusi.

“Constraints on learning in dynamic synapses”.

Network: Comp. Neural, 3(4):443–464, (1992) .

5 68



D. J. Amit and S. Fusi.

“Learning in neural networks with material synapses”.

Neural Comput., 6(5):957–982, (1994) .

5 68 100

References II



Carl C. H. Petersen, Robert C. Malenka, Roger A. Nicoll, and John J. Hopfield.

“All-or-none potentiation at CA3-CA1 synapses”.

Proc. Natl. Acad. Sci. U.S.A., 95(8):4732–4737, (1998) .

5



Daniel H. O'Connor, Gayle M. Wittenberg, and Samuel S.-H. Wang.

“Graded bidirectional synaptic plasticity is composed of switch-like unitary events”.

Proc. Natl. Acad. Sci. U.S.A., 102(27):9679–9684, (2005) .

5

References III



M. P. Coba, A. J. Pocklington, M. O. Collins, M. V. Kopanitsa, R. T. Uren, S. Swamy, M. D. Croning, J. S. Choudhary, and S. G. Grant.

“Neurotransmitters drive combinatorial multistate postsynaptic density networks”.

Sci Signal, 2(68):ra19, (2009) .

6



Johanna M. Montgomery and Daniel V. Madison.

“State-Dependent Heterogeneity in Synaptic Depression between Pyramidal Cell Pairs”.

Neuron, 33(5):765 – 777, (2002) .

6

References IV



Y. P. Tang, E. Shimizu, G. R. Dube, C. Rampon, G. A. Kerchner, M. Zhuo, G. Liu, and J. Z. Tsien.

“Genetic enhancement of learning and memory in mice”.

Nature, 401(6748):63–69, (Sep, 1999) .

10



Gaël Malleret, Ursula Haditsch, David Genoux, Matthew W. Jones, Tim V.P. Bliss, Amanda M. Vanhoose, Carl Weitlauf, Eric R. Kandel, Danny G. Winder, and Isabelle M. Mansuy.

“Inducible and Reversible Enhancement of Learning, Memory, and Long-Term Potentiation by Genetic Inhibition of Calcineurin”.

Cell, 104(5):675 – 686, (2001) .

10

References V



J. S. Guan, S. J. Haggarty, E. Giacometti, J. H. Dannenberg, N. Joseph, J. Gao, T. J. Nieland, Y. Zhou, X. Wang, R. Mazitschek, J. E. Bradner, R. A. DePinho, R. Jaenisch, and L. H. Tsai.

"HDAC2 negatively regulates memory formation and synaptic plasticity".

Nature, 459(7243):55–60, (May, 2009) .

10



M. Migaud, P. Charlesworth, M. Dempster, L. C. Webster, A. M. Watabe, M. Makhinson, Y. He, M. F. Ramsay, R. G. Morris, J. H. Morrison, T. J. O'Dell, and S. G. Grant.

"Enhanced long-term potentiation and impaired learning in mice with mutant postsynaptic density-95 protein".

Nature, 396(6710):433–439, (December, 1998) .

11

References VI



N. Uetani, K. Kato, H. Ogura, K. Mizuno, K. Kawano, K. Mikoshiba, H. Yakura, M. Asano, and Y. Iwakura.

“Impaired learning with enhanced hippocampal long-term potentiation in PTPdelta-deficient mice”.

EMBO J., 19(12):2775–2785, (June, 2000) .

11



Patrick R Cox, Velia Fowler, Bisong Xu, J.David Sweatt, Richard Paylor, and Huda Y Zoghbi.

“Mice lacking tropomodulin-2 show enhanced long-term potentiation, hyperactivity, and deficits in learning and memory”.

Molecular and Cellular Neuroscience, 23(1):1 – 12, (2003) .

11

References VII



S.K.E. Koekkoek, K. Yamaguchi, B.A. Milojkovic, B.R. Dortland, T.J.H. Ruigrok, R. Maex, W. De Graaf, A.E. Smit, F. VanderWerf, C.E. Bakker, R. Willemse, T. Ikeda, S. Kakizawa, K. Onodera, D.L. Nelson, E. Mientjes, M. Joosten, E. De Schutter, B.A. Oostra, M. Ito, and C.I. De Zeeuw.

“Deletion of *FMR1* in Purkinje Cells Enhances Parallel Fiber LTD, Enlarges Spines, and Attenuates Cerebellar Eyelid Conditioning in Fragile X Syndrome”.

Neuron, 47(3):339 – 352, (2005) .

11



Mansuo L Hayashi, Se-Young Choi, B.S. Shankaranarayana Rao, Hae-Yoon Jung, Hey-Kyoung Lee, Dawei Zhang, Sumantra Chattarji, Alfredo Kirkwood, and Susumu Tonegawa.

“Altered Cortical Synaptic Morphology and Impaired Memory Consolidation in Forebrain- Specific Dominant-Negative {PAK} Transgenic Mice”.

Neuron, 42(5):773 – 787, (2004) .

11

References VIII



Kris Rutten, Dinah L. Misner, Melissa Works, Arjan Blokland, Thomas J. Novak, Luca Santarelli, and Tanya L. Wallace.

“Enhanced long-term potentiation and impaired learning in phosphodiesterase 4D-knockout (PDE4D-/-) mice”.

European Journal of Neuroscience, 28(3):625–632, (2008) .

11



David Marr.

“A theory of cerebellar cortex”.

The Journal of Physiology, 202(2):437–470, (1969) .

15



James S. Albus.

“A theory of cerebellar function”.

Mathematical Biosciences, 10(1):25 – 61, (1971) .

15

References IX



Masao Ito.

“Neural design of the cerebellar motor control system”.

Brain research, 40(1):81–84, (1972) .

15



Michael J. McConnell, Yanhua H. Huang, Akash Datwani, and Carla J. Shatz.

“H2-Kb and H2-Db regulate cerebellar long-term depression and limit motor learning”.

Proc. Natl. Acad. Sci. U.S.A., 106(16):6784–6789, (2009) .

16



S. Fusi, P. J. Drew, and L. F. Abbott.

“Cascade models of synaptically stored memories”.

Neuron, 45(4):599–611, (February, 2005) .

26

69

70

101

References X



S. Fusi and L. F. Abbott.

“Limits on the memory storage capacity of bounded synapses”.

Nat. Neurosci., 10(4):485–493, (Apr, 2007) .

26

69



A. B. Barrett and M. C. van Rossum.

“Optimal learning rules for discrete synapses”.

PLoS Comput. Biol., 4(11):e1000230, (November, 2008) .

26

69

107



Maurice A Smith, Ali Ghazizadeh, and Reza Shadmehr.

“Interacting Adaptive Processes with Different Timescales Underlie Short-Term Motor Learning”.

PLoS Biol., 4(6):e179, (May, 2006) .

26

References XI



Subhaneil Lahiri and Surya Ganguli.

“A memory frontier for complex synapses”.

In C.J.C. Burges, L. Bottou, M. Welling, Z. Ghahramani, and K.Q. Weinberger, editors, *Adv. Neural Inf. Process. Syst. 26*, pages 1034–1042. NIPS, 2013.

URL <https://papers.nips.cc/paper/4872-a-memory-frontier-for-complex-synapses>.

26

74



Daniel D Ben-Dayan Rubin and Stefano Fusi.

“Long memory lifetimes require complex synapses and limited sparseness”.

Front. Comput. Neurosci., 1:1–14, (November, 2007) .

70

References XII



Christian Leibold and Richard Kempter.

“Sparseness Constrains the Prolongation of Memory Lifetime via Synaptic Metaplasticity”.

Cereb. Cortex, 18(1):67–77, (2008) .

70



Marcus K. Benna and Stefano Fusi.

“Computational principles of synaptic memory consolidation”.

Nature Neuroscience, 19(12):1697–1706, (July, 2016) , arXiv:1507.07580 [q-bio.NC].

70

References XIII



Terry Elliott.

"The mean time to express synaptic plasticity in integrate-and-express, stochastic models of synaptic plasticity induction", jan 2011.

ISSN 08997667.

URL

http://www.mitpressjournals.org/doi/10.1162/NECO_a_00061.

70



Richard D. Emes and Seth G.N. Grant.

"Evolution of Synapse Complexity and Diversity".

Annual Review of Neuroscience, 35(1):111–131, (2012) .

80

81

References XIV



Larry R Squire and Pablo Alvarez.

“Retrograde amnesia and memory consolidation: a neurobiological perspective”.

Current Opinion in Neurobiology, 5(2):169–177, (April, 1995) .

80

81



James L McClelland, Bruce L McNaughton, and Randall C O'Reilly.

“Why there are complementary learning systems in the hippocampus and neocortex: Insights from the successes and failures of connectionist models of learning and memory.”, 1995.

80

81



Jan Born and Ines Wilhelm.

“System consolidation of memory during sleep.”.

Psychological research, 76(2):192–203, (mar, 2012) .

80

81

References XV



Phillip J.E. Attwell, Samuel F. Cooke, and Christopher H. Yeo.
“Cerebellar Function in Consolidation of a Motor Memory”.
Neuron, 34(6):1011–1020, (jun, 2002) .

80 81



Samuel F Cooke, Phillip J E Attwell, and Christopher H Yeo.
“Temporal properties of cerebellar-dependent memory consolidation.”.
J. Neurosci., 24(12):2934–41, (mar, 2004) .

80 81



LE Baum, T Petrie, George Soules, and Norman Weiss.
“A maximization technique occurring in the statistical analysis of probabilistic functions of Markov chains”.
The annals of mathematical statistics, 41(1):164–171, (1970) .

88 89 90 91 92

References XVI



Lawrence R Rabiner and Biing-Hwang Juang.

Fundamentals of speech recognition, volume 14 of *Signal Processing*.

Prentice Hall, Inc., Upper Saddle River, NJ, USA, 1993.

ISBN 0-13-015157-2.

88

89

90

91

92



A. P. Dempster, N. M. Laird, and D. B. Rubin.

“Maximum Likelihood from Incomplete Data via the EM Algorithm”.

Journal of the Royal Statistical Society. Series B (Methodological), (October, 2007) .

88

89

90

91

92

References XVII



Daniel Hsu, Sham M. Kakade, and Tong Zhang.

“A spectral algorithm for learning Hidden Markov Models”.

Journal of Computer and System Sciences, 78(5):1460 – 1480, (2012) ,
arXiv:0811.4413 [cs.LG].

{JCSS} Special Issue: Cloud Computing 2011.

88

89

90

91

92



J.G. Kemeny and J.L. Snell.

Finite markov chains.

Springer, 1960.

109