

# 4th year Review - Statistical Physics Perspectives on Learning in High Dimensions

Advisor: Surya Ganguli

Madhu Advani

Stanford University

April 6, 2014

## Current Research

Optimal Tractable High Dimensional M-estimation

## Future Directions

- ① LN and GLM extensions of M-estimation also Optimal Signal Processing (Structured Coefficients)
- ② Random Dimensionality Reduction
- ③ Phase transitions in clustering Behavior

## Previous Research and Future Plan

- ① Review Paper
- ② Timeline

# Problem Setup

- Consider Statistical inference with  $N$  data points and  $P$  unknowns (predictors).
- (Easy) Classical Regime:  $\kappa = P/N \rightarrow 0$
- (Hard) High Dimensional Regime:  $\kappa = P/N \neq 0$

Outputs generated by:  $y_i = \mathbf{X}_i \cdot \mathbf{w}^0 + \epsilon_i \quad i \in [1, \dots, N]$

- We want to find  $\mathbf{w}^0 \in \mathcal{R}^P$
- Noise  $\epsilon$  not necessarily gaussian

$$\hat{\mathbf{w}} = \arg \min_{\mathbf{w}} \left[ \sum_i \rho(y_i - \mathbf{X}_i \cdot \mathbf{w}) \right]$$

E.g.  $\rho(x) = x^2, |x|, -\ln f(x)$

# Maximum Likelihood

# Classical vs High Dimensional Optimal M-estimation

## Classical

- $N \rightarrow \infty, P/N = \kappa \rightarrow 0$
- $\rho_{\text{opt}} = -\log f$
- $\langle\langle (\hat{w}_i - w_i^0)^2 \rangle\rangle \geq \frac{\kappa}{\int \frac{\kappa}{f'^2}}$

## High Dimensional

- $N, P \rightarrow \infty, P/N = \kappa \in [0, 1]$
- $\rho_{\text{opt}}(x) = -\inf_y \left[ \ln(\zeta_{\hat{q}_0}(y)) + \frac{(x-y)^2}{2\hat{q}_0} \right] \quad \zeta = f * \phi_{\hat{q}_0}$
- $\langle\langle (\hat{w}_i - w_i^0)^2 \rangle\rangle \geq \frac{\kappa}{\int \frac{\kappa}{\zeta'^2}}$

# Adding a Regularizer

$$P(\mathbf{w}^0 | \mathbf{X}, \mathbf{y}) = \frac{P(\mathbf{X}, \mathbf{y} | \mathbf{w}^0) P(\mathbf{w}^0)}{P(\mathbf{X}, \mathbf{y})}$$

## Maximum a Priori

$$\hat{\mathbf{w}}_{\text{MAP}} = \arg \min_{\mathbf{w}} \left[ \sum_i -\log f(y_i - \mathbf{X}_i \cdot \mathbf{w}) + \sum_j -\log g(w_j) \right]$$

## Regularized M-estimation

$$\hat{\mathbf{w}} = \arg \min_{\mathbf{w}} \left[ \sum_i \rho(y_i - \mathbf{X}_i \cdot \mathbf{w}) + \sum_j \sigma(w_j) \right]$$

Note separability. Solvable for convex strategy  $\sigma, \rho$

$$\hat{\mathbf{w}} = \arg \min_{\mathbf{w}} \sum_a \rho(y_a - \mathbf{X}_a \cdot \mathbf{w}) + \sum_i \sigma(w_i)$$

## Applications

- Maximum Likelihood (ML) and MAP commonly applied to High Dimensional Bio-informatics problems, where we should expect poor performance
- Deriving/Understanding the Optimal M-estimator has potential applications for statistical inference and signal processing.
- Tractability of this form of optimization popular: Compressed Sensing, LASSO, Elastic Net

# Statistical Physics Formulation

Define a spin glass system to solve M-estimator inference

## Spin Glass System

Define continuous spins  $\mathbf{u} = \mathbf{w}^0 - \mathbf{w}$ . Let the Energy of the system be a function of these spins

$$E_{\Lambda}(\mathbf{u}) = \sum_i \rho(\mathbf{X}_i \cdot \mathbf{u} + \epsilon_i) + \sum_a \sigma(w_a^0 - u_a)$$

This in turn induces an equilibrium probability distribution on the state

$$P_G(\mathbf{u}) = \frac{e^{-\beta E_{\Lambda}(\mathbf{u})}}{Z_{\Lambda}} \quad Z_{\Lambda} = \int e^{-\beta E_{\Lambda}(\mathbf{u})} d\mathbf{u}$$

$$\lim_{\beta \rightarrow \infty} P_G(\mathbf{u}) = \delta(\mathbf{u} - \mathbf{w}^0 + \hat{\mathbf{w}})$$



## Coupled Equations Relating Order Parameters

$$\left\langle \left\langle (\text{prox}_{c\rho}(\sqrt{q}z + \epsilon) - \sqrt{q}z - \epsilon)^2 \right\rangle \right\rangle_{z,\epsilon} = \kappa q$$

$$\left\langle \left\langle \text{prox}'_{c\rho}(\sqrt{q}z + \epsilon) \right\rangle \right\rangle_{z,\epsilon} = 1 - \kappa$$

## Order Parameters

$$q = c =$$


## Definition

$$\begin{aligned} \text{prox}_f(x) &= \arg \min_y \left[ \frac{(x - y)^2}{2} + f(y) \right] \\ &= [I + \partial f]^{-1}(x) \end{aligned}$$

- Note that both  $f$  and  $\text{prox}_f$  are functions
- Known for many functions: e.g. soft thresholding operator for L1 norm
- Used in proximal algorithms: minima of  $f \Leftrightarrow$  fixed points  $\text{prox}_f$

## Order Parameters

nothing here yet



# Unregularized M-estimators

- $X_{ij} \in \mathcal{N}(0, 1/P)$
- $\rho$  convex
- $\epsilon$  iid

## Optimal M-estimator

$$\rho_{\text{opt}}(x) = -\inf_y \left[ \ln(\zeta_{\hat{q}}(y)) + \frac{(x-y)^2}{2\hat{q}} \right] \quad \zeta_{\hat{q}} = f * \phi_{\hat{q}}$$

$$\hat{q} = \min q \quad \text{s.t.} \quad q l_q = \kappa \quad l_q = \int_{-\infty}^{\infty} \frac{\zeta_q'^2(y)}{\zeta_q(y)} dy$$

- Best possible asymptotic Mean squared error for any convex M-estimator is  $\hat{q}$