

# The Influence of Temperature on Ozone Production under varying NO<sub>x</sub> Conditions – a modelling study

J. Coates<sup>1</sup>, K. A. Mar<sup>1</sup>, N. Ojha<sup>2</sup>, and T. M. Butler<sup>1</sup>

<sup>1</sup>Institute for Advanced Sustainability Studies, Potsdam, Germany

<sup>2</sup>Atmospheric Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany

*Correspondence to:* J. Coates (jane.coates@iass-potsdam.de)

**Abstract.** Surface ozone is a secondary air pollutant produced during the atmospheric photochemical degradation of emitted volatile organic compounds (VOCs) in the presence of sunlight and nitrogen oxides (NO<sub>x</sub>). Temperature directly influences ozone production through speeding up the rates of the chemical reactions and increasing the emissions of VOCs, such as isoprene, from vegetation. In this study, we used a box model to examine the non-linear relationship between ozone, NO<sub>x</sub> and temperature, and compared this to previous observational studies. Under high-NO<sub>x</sub> conditions, an increase in ozone from 20 °C to 40 °C of up to 20 ppbv was due to faster reaction rates while increased isoprene emissions added up to a further 11 ppbv of ozone. The increased oxidation rate of emitted VOC with temperature controlled the rate of O<sub>x</sub> production, the net influence of peroxy nitrates increased net O<sub>x</sub> production per molecule of emitted VOC oxidised. The rate of increase in ozone mixing ratios with temperature from our box model simulations was about half the rate of increase in ozone with temperature observed over central Europe or simulated by a regional chemistry transport model. Modifying the box model setup to approximate stagnant meteorological conditions increased the rate of increase of ozone with temperature as the accumulation of oxidants enhanced ozone production through the increased production of peroxy radicals from the secondary degradation of emitted VOCs. The box model simulations approximating stagnant conditions and the maximal ozone production chemical regime reproduced the 2 ppbv increase in ozone per °C from the observational and regional model data over central Europe. The simulated ozone-temperature relationship was more sensitive to mixing than the choice of chemical mechanism. Our analysis suggests that reductions in NO<sub>x</sub> emissions would be required to offset the additional ozone production due to an increase in temperature in the future.

## 1 Introduction

Surface-level ozone (O<sub>3</sub>) is a secondary air pollutant formed during the photochemical degradation of volatile organic compounds (VOCs) in the presence of nitrogen oxides (NO<sub>x</sub> ≡ NO + NO<sub>2</sub>). Due to the photochemical nature of ozone production, it is strongly influenced by meteorological variables such as temperature (Jacob and Winner, 2009). In particular, heatwaves which are characterised by high temperatures and stagnant meteorological conditions tend to have high ozone levels as was the case during the European heatwave in 2003 (Solberg et al., 2008; Vautard et al., 2005). Furthermore, Otero et al. (2016) showed that temperature was a major meteorological driver of summertime ozone concentrations in many areas of central Europe.

Temperature primarily influences ozone production in two ways: speeding up the rates of many chemical reactions, and increasing emissions of VOCs from biogenic sources (BVOCs) (Sillman and Samson, 1995). While emissions of anthropogenic VOCs (AVOCs) are generally not dependent on temperature, evaporative emissions of some AVOCs do increase with temperature (Rubin et al., 2006). The review of Pusede et al. (2015) provides further details of the temperature-dependent processes  
5 impacting ozone production.

Regional modelling studies over the US (Sillman and Samson, 1995; Steiner et al., 2006; Dawson et al., 2007) examined the sensitivity of ozone production during a pollution episode to increased temperatures. These studies noted that increased temperatures (without changing VOC or NO<sub>x</sub>-conditions) led to higher ozone levels, often exceeding local air quality guidelines. Sillman and Samson (1995) and Dawson et al. (2007) varied the temperature dependence of the PAN (peroxy acetyl  
10 nitrate) decomposition rate during simulations of the eastern US determining the sensitivity of ozone production with temperature to the PAN decomposition rate. In addition to the influence of PAN decomposition on ozone production, Steiner et al. (2006) correlated the increase in ozone mixing ratios with temperature over California to increased mixing ratios of formaldehyde, a secondary degradation product of many VOCs and an important radical source. Steiner et al. (2006) also noted increased emissions of BVOCs at higher temperatures in urban areas with high NO<sub>x</sub> emissions also increased ozone levels  
15 with temperature.

Pusede et al. (2014) used an analytical model constrained by observations over the San Joaquin Valley, California to infer a non-linear relationship between ozone, temperature and NO<sub>x</sub>, similar to the well-known non-linear relationship of ozone production on NO<sub>x</sub> and VOC levels (Sillman, 1999). Moreover, Pusede et al. (2014) showed that temperature can be used as a surrogate for VOC levels when considering the relationship of ozone under different NO<sub>x</sub> conditions.

20 Environmental chamber studies have also been used to analyse the relationship of ozone with temperature using a fixed mixture of VOCs. The chamber experiments of Carter et al. (1979) and Hatakeyama et al. (1991) showed increases in ozone from a VOC mix with temperature. Both studies compared the concentration time series of ozone and nitrogen-containing compounds (NO<sub>x</sub>, PAN, HNO<sub>3</sub>) at various temperatures linking the maximum ozone concentration to the decrease in PAN concentrations at temperatures greater than 303 K.

25 Despite many studies considering the ozone-temperature relationship from an observational and chamber study perspective, modelling studies focusing on the detailed chemical processes of the ozone-temperature relationship under different NO<sub>x</sub> conditions have not been performed (to our knowledge). The regional modelling studies described previously concentrated on reproducing ozone levels (using a single chemical mechanism) over regions with known meteorology and NO<sub>x</sub> conditions then varying the temperature. These modelling studies did not consider the relationship between ozone, NO<sub>x</sub> and temperature. The  
30 review of Pusede et al. (2015) also highlights a general lack of modelling studies looking at the relationship of ozone with temperature under different NO<sub>x</sub> conditions.

Comparisons of different chemical mechanisms, such as Emmerson and Evans (2009) and Coates and Butler (2015), showed that different representations of tropospheric chemistry influenced ozone production. Neither of these studies examined the ozone-temperature relationship differences between chemical mechanisms. Furthermore, Rasmussen et al. (2013) acknowledged that the modelled ozone-temperature relationship may be sensitive to the choice of chemical mechanism and recom-  
35

mended investigating this sensitivity. Comparing the ozone-temperature relationship predicted by different chemical mechanisms is potentially important for modelling of future air quality due to the expected increase in heatwaves (Karl and Trenberth, 2003).

In this study, we use an idealised box model to determine how ozone levels vary with temperature under different  $\text{NO}_x$  conditions. We determine whether faster chemical reaction rates or increased BVOC emissions have a greater influence on instantaneous ozone production with higher temperature under different  $\text{NO}_x$  conditions. Furthermore, we compare the ozone-temperature relationship produced by different chemical mechanisms. We compare the rate of increase of ozone with temperature obtained from the box model to both observations and regional model output. Additionally, we look at the role of stagnation on the rate of increase of ozone with temperature.

## 2 Methodology

### 2.1 Model Setup

We used the MECCA box model (Sander et al., 2005) to determine the important gas-phase chemical processes for ozone production under different temperatures and  $\text{NO}_x$  conditions. The MECCA box model was set up as described in Coates and Butler (2015) and updated to include vertical mixing with the free troposphere using a diurnal cycle for the PBL height. The vertical mixing scheme was based on the approach of Lourens et al. (2016) with the model using the mean mixing layer height from the BAERLIN campaign over Berlin, Germany (Bonn et al., 2016).

Simulations were run for daylight hours in one full day using a latitude of  $51^\circ\text{N}$ , broadly representative of urban conditions in central Europe. Methane was fixed at 1.7 ppmv throughout the model run, carbon monoxide (CO) and ozone were initialised at 200 ppbv and 40 ppbv and then allowed to evolve freely throughout the simulation. All VOC emissions were held constant until noon simulating a plume of freshly-emitted VOC. the mixing ratios of  $\text{O}_3$ , CO and  $\text{CH}_4$  in the free troposphere were respectively set to 50 ppbv, 116 ppbv and 1.8 ppmv. These conditions were taken from the MATCH-MPIC chemical weather forecast model on the 21st March (the start date of the simulations). The model results (<http://cwfi.iass-potsdam.de/>) at the 700 hPa height were chosen and the daily average was used as input into the boxmodel.

Separate box model simulations were performed by systematically varying the temperature between 288 and 313 K (15–40 °C) in steps of 0.5 K. NO emissions were systematically varied between  $5.0 \times 10^9$  and  $1.5 \times 10^{12}$  molecules(NO)  $\text{cm}^{-2} \text{s}^{-1}$  in steps of  $1 \times 10^{10}$  molecules(NO)  $\text{cm}^{-2} \text{s}^{-1}$  at each temperature step. At 20 °C, these NO emissions corresponded to peak  $\text{NO}_x$  mixing ratios of 0.02 ppbv and 10 ppbv respectively, this range of  $\text{NO}_x$  mixing ratios covers the  $\text{NO}_x$  conditions found in pristine and urban conditions (von Schneidemesser et al., 2015).

All simulations were repeated using different chemical mechanisms to investigate whether the relationship between ozone, temperature and  $\text{NO}_x$  changes using different representations of ozone production chemistry. The reference chemical mechanism was the near-explicit Master Chemical Mechanism, MCMv3.2, (Jenkin et al., 1997, 2003; Saunders et al., 2003; Rickard et al., 2015). The reduced chemical mechanisms in our study were Common Representative Intermediates, CRIv2 (Jenkin et al., 2008), Model for OZone and Related Chemical Tracers, MOZART-4 (Emmons et al., 2010), Regional Acid Deposition Model,

RADM2 (Stockwell et al., 1990) and the Carbon Bond Mechanism, CB05 (Yarwood et al., 2005). These reduced chemical mechanisms were chosen as they are currently used by modelling groups in 3D regional and global models (Baklanov et al., 2014). Coates and Butler (2015) described the implementation of these chemical mechanisms in MECCA.

The chemical mechanisms use temperature-dependent rate constants,  $k(T)$ , to represent temperature-dependent chemical processes. These temperature-dependent chemical processes include the initial oxidation of VOC, peroxy nitrate ( $RO_2NO_2$ ) formation and destruction, and reactions between peroxy radicals and NO leading to alkyl nitrate ( $RONO_2$ ) formation. However, not all chemical mechanisms represent the same chemical processes by a temperature-dependent rate constant. For example, in CB05, the rate constant of reactions  $RONO_2$  formation during isoprene degradation is temperature dependent while  $RONO_2$  formation during alkane degradation is temperature independent. None of the chemical mechanisms represent the alkyl nitrate branching ratio as a temperature dependent process,  $RONO_2$  chemistry is not well known (Pusede et al., 2015) although experiments have shown the temperature dependence of this branching ratio (Atkinson et al., 1987). Pusede et al. (2015) described the influence that the temperature dependence of the  $RONO_2$  branching ratio may have on ozone production and more work is required to adequately represent this process within chemical mechanisms.

Model runs were repeated using a temperature-independent and temperature-dependent source of BVOC emissions to determine the relative importance of increased emissions of BVOC and faster reaction rates of chemical processes for the increase of ozone with temperature. Many types of VOCs are emitted from vegetation with isoprene and monoterpenes having globally the largest yearly emissions, 535 and 162 Tg yr<sup>-1</sup> (Guenther et al., 2012). Temperature-dependent emissions of these highly-reactive BVOC in urban areas, characterised by high NO<sub>x</sub> emissions, during the summer months in urban areas leads to high levels of ozone pollution during the summertime, for example the study of Wang et al. (2013) attributed high summertime levels of ozone in Taipei to increased isoprene emissions from vegetation during the hotter summer months. The influence of BVOCs in urban areas has additional complexity as many BVOCs also react with ozone and stomatal uptake of ozone by vegetation are also sinks of ozone, the review of Calfapietra et al. (2013) discusses the role of BVOCs in urban areas in more detail.

In all model runs, biogenic emissions of monoterpenes and isoprene are included with the model runs using a temperature-dependent source of BVOC emissions considered only the temperature-dependence of isoprene emissions as specified by MEGAN2.1 (Guenther et al., 2012), Sect. 2.3 provides further details. We focused on the temperature-dependence of isoprene emissions as isoprene is the most important BVOC on the global scale (Guenther et al., 2006), future work should assess the influence of temperature-dependent emissions of monoterpenes on ozone production especially to determine whether the increased emissions are a larger source or sink of ozone. In the temperature-dependent set of model simulations, only isoprene emissions were dependent on temperature and all other emissions were constant in all simulations. Our assumption that all anthropogenic VOC emissions are temperature dependent is not the case in reality where evaporative emissions increase with temperature (Rubin et al., 2006), temperature dependent emission factors of anthropogenic emissions require detailed local knowledge of traffic fleets for example and our modelling study was not meant to focus on a particular region. The influence of temperature dependent BVOC emissions on ozone production requires detailed knowledge of regional emissions and the strength of these sources, our box modelling study does not aim to characterise the influence of temperature dependent emis-

sions of a particular area and thus was kept in a general state. Furthermore, isoprene has been measured from vehicular exhausts in urban areas by Borbon et al. (2001) and once again since we are not characterising the influence of temperature dependent emissions of a particular region the detailed knowledge of the strength of this source of isoprene, that is not included in the TNO-MACC\_III emission inventory. Further modelling work assessing this influence on ozone production would be highly useful for mitigating ozone pollution in urban areas.

## 2.2 VOC Emissions

Emissions of urban AVOC over central Europe were taken from the TNO-MACC\_III emission inventory for the Benelux (Belgium, Netherlands and Luxembourg) region for the year 2011. TNO-MACC\_III is the updated TNO-MACC\_II emission inventory created using the same methodology as Kuenen et al. (2014) and based upon improvements to the existing emission inventory during AQMEII-2 (Pouliot et al., 2015).

Temperature-independent emissions of isoprene and monoterpenes from biogenic sources were calculated as a fraction of the total AVOC emissions from each country in the Benelux region. This data was obtained from the supplementary data available from the EMEP (European Monitoring and Evaluation Programme) model (Simpson et al., 2012). Temperature-dependent emissions of isoprene are described in Sect. 2.3.

Table 1 shows the quantity of VOC emissions from each source category and the temperature-independent BVOC emissions. These AVOC emissions were assigned to chemical species and groups based on the profiles provided by TNO. The NMVOC emissions were speciated to MCMv3.2 species as described by von Schneidemesser et al. (2016). For simulations done with other chemical mechanisms, the VOC emissions represented by the MCMv3.2 were mapped to the mechanism species representing VOC emissions in each reduced chemical mechanism based on the recommendations of the source literature and Carter (2015). The VOC emissions in the reduced chemical mechanisms were weighted by the carbon numbers of the MCMv3.2 species and the emitted mechanism species, thus keeping the amount of emitted reactive carbon constant between simulations. The supplementary data outlines the primary VOC and calculated emissions with each chemical mechanism.

## 2.3 Temperature Dependent Isoprene Emissions

Temperature-dependent emissions of isoprene were estimated using the MEGAN2.1 algorithm for calculating the emissions of VOC from vegetation (Guenther et al., 2012). Emissions from nature are dependent on many variables including temperature, radiation and age of vegetation but for the purpose of our study all variables except temperature were held constant. The idealised nature of our study was designed to be generally applicable to urban areas rather than performed detailed simulations representative of a particular region over a certain time frame. Thus to directly compare the effects of temperature-dependent emissions of isoprene to temperature-dependent chemistry, the MEGAN2.1 parameters used to calculate isoprene emissions online by the model were chosen to give similar isoprene mixing ratios at 20 °C to the temperature-independent emissions of isoprene in order to compare the effects of increased isoprene emissions with temperature. The estimated emissions of isoprene with MEGAN2.1 using these assumptions are illustrated in Fig. 1 and show the expected exponential increase in isoprene emissions with temperature (Guenther et al., 2006).

The estimated emissions of isoprene at 20 °C lead to 0.07 ppbv of isoprene in our simulations while at 30 °C, the increased emissions of isoprene using MEGAN2.1 estimations lead to 0.35 ppbv of isoprene in the model. A measurement campaign over Essen, Germany (Wagner and Kuttler, 2014) measured 0.1 ppbv of isoprene at temperature 20 °C and 0.3 ppbv of isoprene were measured at 30 °C. The similarity of the simulated and observed isoprene mixing ratios indicates that the MEGAN2.1 variables chosen for calculating the temperature-dependent emissions of isoprene were suitable for simulating urban conditions over central Europe.

### 3 Results and Discussion

#### 3.1 Relationship between Ozone, NO<sub>x</sub> and Temperature

Figure 2 depicts the peak mixing ratio from ozone of each simulation as a function of the total NO<sub>x</sub> emissions and temperature when using a temperature-independent and temperature-dependent source of isoprene emissions for each chemical mechanism. A non-linear relationship of ozone mixing ratios with NO<sub>x</sub> and temperature is produced by each chemical mechanism. This non-linear relationship is similar to that determined by Pusede et al. (2014) using an analytical model constrained to observational measurements over the San Joaquin Valley, California.

Higher peak ozone mixing ratios are produced when using a temperature-dependent source of isoprene emissions (Fig. 2). The highest mixing ratios of peak ozone are produced at high temperatures and moderate emissions of NO<sub>x</sub> regardless of the temperature dependence of isoprene emissions. Conversely, the least amount of peak ozone is produced with low emissions of NO<sub>x</sub> over the whole temperature range (15 – 40 °C) when using both a temperature-independent and temperature-dependent source of isoprene emissions.

The contours of ozone mixing ratios as a function of NO<sub>x</sub> and temperature can be split into three NO<sub>x</sub> regimes (Low-NO<sub>x</sub>, Maximal-O<sub>3</sub> and High-NO<sub>x</sub>), similar to the NO<sub>x</sub> regimes defined for the non-linear relationship of ozone with VOC and NO<sub>x</sub>. The Low-NO<sub>x</sub> regime corresponds with regions having little increase in ozone with temperature, also called the NO<sub>x</sub>-sensitive regime. The High-NO<sub>x</sub> (or NO<sub>x</sub>-saturated) regime is when ozone levels increase rapidly with temperature. The contour ridges correspond to regions of maximal ozone production; this is the Maximal-O<sub>3</sub> regime. Pusede et al. (2014) showed that temperature can be used as a proxy for VOC, thus we assigned the ozone mixing ratios from each box model simulation to a NO<sub>x</sub> regime based on the H<sub>2</sub>O<sub>2</sub>:HNO<sub>3</sub> ratio. This ratio was used by Sillman (1995) to designate ozone to NO<sub>x</sub> regimes based on NO<sub>x</sub> and VOC levels. The Low-NO<sub>x</sub> regime corresponds to H<sub>2</sub>O<sub>2</sub>:HNO<sub>3</sub> ratios less than 0.5, the High-NO<sub>x</sub> regime corresponds to ratios larger than 0.3 and ratios between 0.3 and 0.5 correspond to the Maximal-O<sub>3</sub> regime.

The peak ozone mixing ratio from each simulation was assigned to a NO<sub>x</sub> regime based on the H<sub>2</sub>O<sub>2</sub>:HNO<sub>3</sub> ratio of that simulation. The peak ozone mixing ratios assigned to each NO<sub>x</sub> regime at each temperature were averaged, and illustrated in Fig. 3 for each chemical mechanism and each type of isoprene emissions (temperature independent and temperature dependent). We define the absolute increase in ozone from 20 °C to 40 °C due to faster reaction rates as the difference between ozone mixing ratios from 20 °C to 40 °C when using a temperature-independent source of isoprene emissions. When using a temperature-dependent source of isoprene emissions, the difference in ozone mixing ratios from 20 °C to 40 °C minus the increase due to

faster reaction rates, gives the absolute increase in ozone mixing ratios from increased isoprene emissions. These differences are represented graphically in Fig. 3 and summarised in Table 2.

Table 2 shows that the absolute increase in ozone with temperature due to chemistry (i.e. faster reaction rates) is larger than the absolute increase in ozone due to increased isoprene emissions for each chemical mechanism and each  $\text{NO}_x$  regime. In all cases the absolute increase in ozone with temperature is largest under High- $\text{NO}_x$  conditions and lowest with Low- $\text{NO}_x$  conditions (Fig. 3 and Table 2). The increase in ozone mixing ratio from 20 °C to 40 °C due to faster reaction rates with High- $\text{NO}_x$  conditions is almost double that with Low- $\text{NO}_x$  conditions. In the Low- $\text{NO}_x$  regime, the increase of ozone with temperature using the reduced chemical mechanisms (CRIV2, MOZART-4, CB05 and RADM2) is similar to that from the MCMv3.2. Larger differences occur in the Maximal- $\text{O}_3$  and High- $\text{NO}_x$  regimes.

All reduced chemical mechanisms except RADM2 have similar increases in ozone due to increased isoprene emissions as the MCMv3.2 (Table 2). RADM2 produces 3 ppbv less ozone than the MCMv3.2 due to increased isoprene emissions in each  $\text{NO}_x$  regime, indicating that this difference is due the representation of isoprene degradation chemistry in RADM2.

Coates and Butler (2015) compared ozone production in different chemical mechanisms to the MCMv3.2 using the TOPP metric (Tagged Ozone Production Potential) as defined in Butler et al. (2011) and showed that less ozone is produced per molecule of isoprene emitted using RADM2 than with MCMv3.2. The degradation of isoprene has been extensively studied and it is well-known that methyl vinyl ketone (MVK) and methacrolein are signatures of isoprene degradation (Atkinson, 2000). All chemical mechanisms in our study except RADM2 explicitly represent MVK and methacrolein (or in the case of CB05, a lumped species representing both these secondary degradation products). RADM2 does not represent methacrolein and the mechanism species representing ketones (KET) is a mixture of acetone and methyl ethyl ketone (MEK) (Stockwell et al., 1990). Thus the secondary degradation of isoprene in RADM2 is unable to represent the ozone production from the further degradation of the signature secondary degradation products of isoprene, MVK and methacrolein. Updated versions of RADM2, RACM (Stockwell et al., 1997) and RACM2 (Goliff et al., 2013), sequentially included methacrolein and MVK and with these updates the ozone production from isoprene oxidation approached that of the MCMv3.2 (Coates and Butler, 2015).

The intermechanism differences between the ozone contours of Fig. 2 obtained from the chemical mechanisms are not significant, the largest differences are produced at the highest temperatures and High- $\text{NO}_x$  conditions. The maximum difference between ozone mixing ratios produced using the chemical mechanisms (at 40 °C) in the High- $\text{NO}_x$  state (Fig. 3) is 10 ppbv when using both a temperature-independent and temperature-dependent source of isoprene emissions. These differences in ozone mixing ratios between the chemical mechanisms are similar to that obtained in Coates and Butler (2015) (8 ppbv) and von Schneidemesser et al. (2016) (7 ppbv).

The Maximal- $\text{O}_3$  regime is reached with  $\sim 30$  % more  $\text{NO}_x$  emissions using CB05 and RADM2 than the detailed MCMv3.2 chemical mechanism throughout the temperature range of our study and for both a temperature-independent and -dependent source of isoprene emissions. The CRIV2 and MOZART-4 chemical mechanisms require very similar  $\text{NO}_x$  emissions to the MCMv3.2 at each temperature to reach the maximum ozone production. This indicates that the increase of ozone with temperature using CB05 and RADM2 is more sensitive to the faster reaction rates (temperature-independent emissions) and increased isoprene emissions (temperature-dependent emissions). The larger slope of the increase of ozone with temperature (Table 3)

obtained with CB05 and RADM2 compared to the other chemical mechanisms is consistent with the conclusion that both these chemical mechanisms produce a more NO<sub>x</sub>-saturated state. Thus when using CB05 and RADM2 for air quality studies, these chemical mechanisms would predict a higher sensitivity of ozone reduction to decreasing VOC-emissions. The study using the 3D WRF-Chem model of Mar et al. (2016) compared ozone levels produced using MOZART-4 and RADM2 and also noted that RADM2 is in a more NO<sub>x</sub>-saturated state than MOZART-4. These differences in the sensitivity of ozone to NO<sub>x</sub> and VOC emissions using different chemical mechanisms highlight the need for air quality studies to assess the chemical scheme used by the model by performing simulations using different chemical mechanisms.

To re-cap, a very similar non-linear relationship between ozone, temperature and NO<sub>x</sub> is produced by each chemical mechanism with the absolute increase in ozone with temperature due to temperature-dependent chemistry larger than the increase in ozone with temperature due to temperature-dependent isoprene emissions. In order to determine the chemical processes responsible for the increased ozone with temperature, we analyse the production and consumption budgets of ozone in Sect. 3.2.

### 3.2 Ozone Production and Consumption Budgets

In order to understand the temperature dependence of ozone production directly, we examine the modelled day-time production and consumption budgets of O<sub>x</sub> ( $\equiv$  O<sub>3</sub> + NO<sub>2</sub> + O(<sup>1</sup>D) + O) normalised by the total chemical loss rate of the emitted VOC (Fig. 4). Chemical reactions contributing to both production and consumption of O<sub>x</sub> have temperature-dependent rate constants, for example the production and decomposition of peroxy nitrates, thus to consider the effects of temperature-dependent reaction rates we analysed the production and consumption budgets of O<sub>x</sub>. The O<sub>x</sub> budgets are assigned to each NO<sub>x</sub> regime for each chemical mechanism and type of isoprene emissions.

The chemical reactions producing and consuming O<sub>x</sub> were determined and then allocated to major chemical categories in Fig. 4. Production of O<sub>x</sub> is through the reactions of peroxy radicals with NO and in Fig. 4 the peroxy radicals are subdivided into ‘HO2’, ‘RO2’, ‘ARO2’ representing the reactions of NO with HO2, alkyl peroxy radicals and acyl peroxy radicals respectively. Thus at each time step the total O<sub>x</sub> production rate is given by

$$PO_x = k_{HO_2+NO}[HO_2][NO] + \sum_i k_{RO_{2,i}+NO}[RO_{2,i}][NO] + \sum_j k_{ARO_{2,j}+NO}[ARO_{2,j}][NO] \quad (1)$$

for each alkyl peroxy radical  $i$  and acyl peroxy radical  $j$  represented by each chemical mechanism. The net contributions of peroxy nitrates, inorganic reactions and any other remaining organic reactions on the O<sub>x</sub> budget are represented by the ‘RO2NO2’, ‘Inorganic’ and ‘Other Organic’ categories in Fig. 4. The net contributions of these categories to the O<sub>x</sub> budget was calculated by subtracting the cumulative day-time consumption rate from the production rate of the reactions contributing to each category. For example, RO2NO2 produce O<sub>x</sub> when thermally decomposing or reacting with OH and consume O<sub>x</sub> when produced are produced from RO<sub>2</sub> reacting with NO<sub>2</sub>. Hence, at each time step the net contribution of RO2NO2 to the O<sub>x</sub> budget was calculated by

$$\sum_k k_{RO_2NO_{2,k}}[RO_2NO_{2,k}] + \sum_k k_{RO_2NO_{2,k}+OH}[RO_2NO_{2,k}][OH] - \sum_k k_{RO_{2,k}+NO_2}[RO_{2,k}][NO_2] \quad (2)$$



for each peroxy nitrate species  $k$  represented by each chemical mechanism. In Fig. 4 the cumulative day-time budgets were calculated by summing the net contributions of the reaction rates of each category over the day-time period and normalised by the total chemical loss rate of the emitted VOC. The absolute production and consumption budgets allocated to the major categories are displayed in Fig. 5a

- 5 Figure 4 also illustrates the net production or consumption of  $O_x$  in each case. The ratio of net ozone to net  $O_x$  production was practically constant with temperature in all cases showing that using  $O_x$  budgets as a proxy for ozone budgets was suitable at each temperature in our study. The absolute production and consumption budgets of  $O_x$  allocated to the same source categories as Fig. 4 are included in the supplementary material and further illustrate the increase in  $O_x$  production with temperature.
- 10 The net  $O_x$  production efficiency increases from 20 °C to 40 °C by  $\sim 0.25$  molecules of  $O_x$  per molecule of VOC oxidised with each  $NO_x$ -condition and type of isoprene emissions using the detailed MCMv3.2 chemical mechanism (Fig. 4). A lower increase in normalised net  $O_x$  production efficiency from 20 °C to 40 °C was obtained with the reduced chemical mechanisms ( $\sim 0.2$  molecules of  $O_x$  per molecule of VOC oxidised with CRIV2, CB05 and RADM2, and  $\sim 0.1$  molecules of  $O_x$  per molecule of VOC oxidised using MOZART-4). The increase in net  $O_x$  production efficiency is due to the increased contribution
- 15 with temperature of acyl peroxy radicals (ARO2) reacting with NO and the decreased net contribution with temperature of  $RO_2NO_2$  (peroxy nitrates) to the normalised  $O_x$  budgets.

The increased contribution of ARO2 to  $O_x$  production with temperature is linked to the decreased net contribution of  $RO_2NO_2$  with temperature to  $O_x$  budgets as peroxy nitrates are produced from the reactions of acyl peroxy radicals with  $NO_2$ . The decomposition rate of peroxy nitrates is strongly temperature dependent and at higher temperatures the faster decomposition rate of  $RO_2NO_2$  leads to faster release of acyl peroxy radicals and  $NO_2$ . Thus the equilibrium of  $RO_2NO_2$  shifts

20 towards thermal decomposition with increasing temperature leading to the increased contribution of ARO2 with temperature to  $O_x$  production (Fig. 4). Dawson et al. (2007) attributed the increase in maximum 8 h ozone mixing ratios with temperature during a modelling study over the eastern US to the decrease in PAN lifetime with temperature. Steiner et al. (2006) also recognised that the decrease in PAN lifetime with temperature contributed to the increase of ozone with temperature concluding that

25 the combined effects of increased oxidation rates of VOC and faster PAN decomposition increased the production of ozone with temperature.

As the production efficiency of  $O_x$  remains constant with temperature ( $\sim 2$  molecules of  $O_x$  per molecule of VOC oxidised, Fig. 4), the rate of  $O_x$  production is controlled by the oxidation of VOCs. Faster oxidation of VOCs with temperature speeds up the production of peroxy radicals increasing ozone production when peroxy radicals react with NO to produce  $NO_2$ . The reactivity of VOCs has been linked to ozone production (e.g. Kleinman (2005), Sadanaga et al. (2005)) and the review of Pusede

30 et al. (2015) acknowledged the importance of organic reactivity and radical production to the ozone-temperature relationship. Also, the modelling study of Steiner et al. (2006) noted that the increase in initial oxidation rates of VOCs with temperature leads to increased formaldehyde concentrations and in turn an increase of ozone as formaldehyde is an important source of  $HO_2$  radicals.

### 3.3 Comparison to Observations and 3D Model Simulations

This section compares the results from our idealised box model simulations to real-world observations and model output from a 3D model. Using the interpolated observations of the maximum daily 8 h mean (MDA8) of ozone from Schnell et al. (2015) and the meteorological observational data set of the ERA-Interim re-analysis, Otero et al. (2016) showed that over the summer (JJA) months, temperature is the main meteorological driver of ozone production over many regions of central Europe. Model output from the 3D WRF-Chem regional model using MOZART-4 chemistry set-up over the European domain for simulations of the year 2007 from Mar et al. (2016) was used to further compare the box model simulations to a model including more meteorological processes than our box model.

Figure 6 compares the observational and WRF-Chem data from summer 2007 averaged over central and eastern Germany, where summertime ozone values are driven by temperature (Otero et al., 2016), to the MDA8 values of ozone from the box model simulations for each chemical mechanism (solid lines). Despite a high bias in simulated ozone in WRF-Chem, the rate of change of ozone with temperature from the WRF-Chem simulations (2.05 ppbv/°C) is similar to the rate of change of ozone with temperature from the observed data (2.15 ppbv/°C). The differences in ozone production between the different chemical mechanisms with the box model are small compared to the spread of the observational and WRF-Chem data. A temperature-dependent source of isoprene with high-NO<sub>x</sub> conditions produces the highest ozone-temperature slope, but is still lower than the observed ozone-temperature slope by a factor of two. In particular, the box model simulations over-predict the ozone values at lower temperatures and under-predict the ozone values at higher temperatures compared to the observed data.

In our simulations, we focused on instantaneous production of ozone from a freshly-emitted source of VOC with mixing of clean air from the free troposphere due to a growing boundary layer. This box model setup did not consider stagnant atmospheric conditions characterised by low wind speeds slowing the transport of ozone and its precursors away from sources. Stagnant conditions have been correlated to high-ozone episodes in the summer over eastern US (Jacob et al., 1993).

In order to investigate the sensitivity of ozone production to mixing, box model simulations were performed without vertical mixing to approximate stagnant conditions that favour accumulation of secondary VOC oxidation products. The ozone-temperature relationship obtained with each chemical mechanism, using both a temperature-independent and temperature-dependent source of isoprene emissions and the different NO<sub>x</sub> conditions are displayed in Fig. 6 (dotted lines). Table 3 summarises the slopes ( $m_{O_3-T}$ ) of the linear fits of the box model simulations displayed in Fig. 6 in ppbv of ozone per °C determining the rate of increase of ozone with temperature, for both case: with, and without mixing.

For all chemical mechanisms, the rate of increase of ozone with temperature increased in the box model simulations without mixing. The  $m_{O_3-T}$  calculated from the box model simulations without mixing using a temperature-dependent source of isoprene and with Maximal-O<sub>3</sub> conditions (ranging between 2.0 and 2.4 ppbv/°C) are very similar to the slopes of the observational and WRF-Chem results (2.1 and 2.2 ppbv/°C, respectively). The differences in  $m_{O_3-T}$  when not including mixing in the box model compared to the differences in  $m_{O_3-T}$  between chemical mechanisms in Table 3 show that the ozone-temperature relationship using our box model setup is more sensitive to mixing than the choice of chemical mechanism.

An analysis similar to that presented in Fig. 4 shows no appreciable difference between the cases with and without mixing (not shown). The chemical production of  $O_x$  in each chemical mechanism normalised by the chemical loss rate of VOC remains unchanged. Furthermore, the supplementary material includes the absolute production and consumption budgets of  $O_x$  which also show the increased  $O_x$  production with temperature for the simulations performed without mixing. From this we conclude  
5 that the increased ozone production seen in the box model simulations with reduced mixing is due to enhanced OH reactivity from secondary VOC oxidation products.

## 4 Conclusions

In this study, we determined the effects of temperature on ozone production using a box model over a range of temperatures and  $NO_x$  conditions with a temperature-independent and temperature-dependent source of isoprene emissions. These simulations  
10 were repeated using reduced chemical mechanism schemes (CRIV2, MOZART-4, CB05 and RADM2) typically used in 3D models and compared to the near-explicit MCMv3.2 chemical mechanism.

Each chemical mechanism produced a non-linear relationship of ozone with temperature and  $NO_x$  with the most ozone produced at high temperatures and moderate emissions of  $NO_x$ . Conversely, lower  $NO_x$  levels led to a minimal increase of ozone with temperature. Thus air quality in a future with higher temperatures would benefit from reductions in  $NO_x$  emissions.  
15 Faster reaction rates at higher temperatures were responsible for a greater absolute increase in ozone than increased isoprene emissions. In our simulations, ozone production was controlled by the increased rate of VOC oxidation with temperature. The net influence of peroxy nitrates increased the net production of  $O_x$  per molecule of emitted VOC oxidised with temperature.

The rate of increase of ozone with temperature using observational data over Europe was twice as high as the rate of increase of ozone with temperature when using the box model. This was consistent with our box model setup not representing stagnant  
20 atmospheric conditions that are inherently included in observational data and models including meteorology, such as WRF-Chem. In model simulations without mixing the rate of increase of ozone with temperature was faster than the simulations including mixing. The simulations without mixing and a maximal ozone production chemical regime led to very similar rates of increase of ozone with temperature to the observational and WRF-Chem data. Furthermore, the ozone-temperature relationship was more sensitive to mixing than the choice of chemical mechanism.

25 *Author contributions.* T. M. Butler and J. Coates designed the experiment, J. Coates performed box model simulations and analysis. K. A. Mar and N. Ojha performed WRF-Chem model runs and provided this data. J. Coates prepared the manuscript with comments from all co-authors.

*Acknowledgements.* The authors would like to thank Noelia Otero Felipe for providing the ERA-Interim data.


## References

- Atkinson, R.: Atmospheric chemistry of VOCs and NO<sub>x</sub>, *Atmospheric Environment*, 34, 2063–2101, 2000.
- Atkinson, R., Aschmann, S. M., and Winer, A. M.: Alkyl nitrate formation from the reaction of a series of branched RO<sub>2</sub> radicals with NO as a function of temperature and pressure, *Journal of Atmospheric Chemistry*, 5, 91–102, 1987.
- 5 Baklanov, A., Schlünzen, K., Suppan, P., Baldasano, J., Brunner, D., Aksoyoglu, S., Carmichael, G., Douros, J., Flemming, J., Forkel, R., Galmarini, S., Gauss, M., Grell, G., Hirtl, M., Joffre, S., Jorba, O., Kaas, E., Kaasik, M., Kallos, G., Kong, X., Korsholm, U., Kurganskiy, A., Kushta, J., Lohmann, U., Mahura, A., Manders-Groot, A., Maurizi, A., Moussiopoulos, N., Rao, S. T., Savage, N., Seigneur, C., Sokhi, R. S., Solazzo, E., Solomos, S., Sørensen, B., Tsegas, G., Vignati, E., Vogel, B., and Zhang, Y.: Online coupled regional meteorology chemistry models in Europe: current status and prospects, *Atmospheric Chemistry and Physics*, 14, 317–398, 2014.
- 10 Bonn, B., von Schneidemesser, E., Andrich, D., Quedenau, J., Gerwig, H., Lüdecke, A., Kura, J., Pietsch, A., Ehlers, C., Klemp, D., Kofahl, C., Nothard, R., Kerschbaumer, A., Junkermann, W., Grote, R., Pohl, T., Weber, K., Lode, B., Schönberger, P., Churkina, G., Butler, T. M., and Lawrence, M. G.: BAERLIN2014 – the influence of land surface types on and the horizontal heterogeneity of air pollutant levels in Berlin, *Atmospheric Chemistry and Physics*, 16, 7785–7811, 2016.
- Borbon, A., Fontaine, H., Veillerot, M., Locoge, N., Galloo, J., and Guillermo, R.: An investigation into the traffic-related fraction of isoprene at an urban location, *Atmospheric Environment*, 35, 3749 – 3760, 2001.
- 15 Butler, T., Lawrence, M., Taraborrelli, D., and Lelieveld, J.: Multi-day ozone production potential of volatile organic compounds calculated with a tagging approach, *Atmospheric Environment*, 45, 4082 – 4090, 2011.
- Calfapietra, C., Fares, S., Manes, F., Morani, A., Sgrigna, G., and Loreto, F.: Role of Biogenic Volatile Organic Compounds (BVOC) emitted by urban trees on ozone concentration in cities: A review, *Environmental Pollution*, 183, 71 – 80, 2013.
- 20 Carter, W. P. L.: Development of a Database for Chemical Mechanism Assignments for Volatile Organic Emissions, *Journal of the Air & Waste Management Association*, 0, 2015.
- Carter, W. P. L., Winer, A. M., Darnall, K. R., and Jr., J. N. P.: Smog chamber studies of temperature effects in photochemical smog, *Environmental Science & Technology*, 13, 1094–1100, 1979.
- Coates, J. and Butler, T. M.: A comparison of chemical mechanisms using tagged ozone production potential (TOPP) analysis, *Atmospheric Chemistry and Physics*, 15, 8795–8808, 2015.
- 25 Dawson, J. P., Adams, P. J., and Pandis, S. N.: Sensitivity of ozone to summertime climate in the eastern USA: A modeling case study , *Atmospheric Environment*, 41, 1494 – 1511, 2007.
- Emmerson, K. M. and Evans, M. J.: Comparison of tropospheric gas-phase chemistry schemes for use within global models, *Atmospheric Chemistry and Physics*, 9, 1831–1845, 2009.
- 30 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43–67, 2010.
- Goliff, W. S., Stockwell, W. R., and Lawson, C. V.: The regional atmospheric chemistry mechanism, version 2, *Atmospheric Environment*, 68, 174 – 185, 2013.
- 35 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, 2006.

- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, 2012.
- Hatakeyama, S., Akimoto, H., and Washida, N.: Effect of temperature on the formation of photochemical ozone in a propene-nitrogen oxide (NOx)-air-irradiation system, *Environmental Science & Technology*, 25, 1884–1890, 1991.
- Jacob, D. J. and Winner, D. A.: Effect of climate change on air quality, *Atmospheric Environment*, 43, 51 – 63, 2009.
- Jacob, D. J., Logan, J. A., Gardner, G. M., Yevich, R. M., Spivakovsky, C. M., Wofsy, S. C., Sillman, S., and Prather, M. J.: Factors regulating ozone over the United States and its export to the global atmosphere, *Journal of Geophysical Research*, 98, 1993.
- Jenkin, M., Watson, L., Utembe, S., and Shallcross, D.: A Common Representative Intermediates (CRI) mechanism for VOC degradation. Part 1: Gas phase mechanism development, *Atmospheric Environment*, 42, 7185 – 7195, 2008.
- Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmospheric Environment*, 31, 81 – 104, 1997.
- Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, *Atmospheric Chemistry and Physics*, 3, 181–193, 2003.
- Karl, T. R. and Trenberth, K. E.: Modern Global Climate Change, *Science*, 302, 1719–1723, 2003.
- Kleinman, L. I.: The dependence of tropospheric ozone production rate on ozone precursors, *Atmospheric Environment*, 39, 575 – 586, 2005.
- Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M., and Denier van der Gon, H. A. C.: TNO-MACC\_II emission inventory; a multi-year (2003–2009) consistent high-resolution European emission inventory for air quality modelling, *Atmospheric Chemistry and Physics*, 14, 10963–10976, 2014.
- Lourens, A. S. M., Butler, T. M., Beukes, J. P., van Zyl, P. G., Fourie, G. D., and Lawrence, M. G.: Investigating atmospheric photochemistry in the Johannesburg-Pretoria megacity using a box model, *South African Journal of Science*, 112, 2016.
- Mar, K. A., Ojha, N., Pozzer, A., and Butler, T. M.: Ozone air quality simulations with WRF-Chem (v3.5.1) over Europe: Model evaluation and chemical mechanism comparison, *Geoscientific Model Development Discussions*, 2016, 1–50, 2016.
- Otero, N., Sillmann, J., Schnell, J. L., Rust, H. W., and Butler, T.: Synoptic and meteorological drivers of extreme ozone concentrations over Europe, *Environmental Research Letters*, 11, 024005, 2016.
- Pouliot, G., van der Gon, H. A. D., Kuenen, J., Zhang, J., Moran, M. D., and Makar, P. A.: Analysis of the emission inventories and model-ready emission datasets of Europe and North America for phase 2 of the AQMEII project, *Atmospheric Environment*, 115, 345–360, 2015.
- Pusede, S. E., Gentner, D. R., Wooldridge, P. J., Browne, E. C., Rollins, A. W., Min, K.-E., Russell, A. R., Thomas, J., Zhang, L., Brune, W. H., Henry, S. B., DiGangi, J. P., Keutsch, F. N., Harrold, S. A., Thornton, J. A., Beaver, M. R., St. Clair, J. M., Wennberg, P. O., Sanders, J., Ren, X., VandenBoer, T. C., Markovic, M. Z., Guha, A., Weber, R., Goldstein, A. H., and Cohen, R. C.: On the temperature dependence of organic reactivity, nitrogen oxides, ozone production, and the impact of emission controls in San Joaquin Valley, California, *Atmospheric Chemistry and Physics*, 14, 3373–3395, 2014.
- Pusede, S. E., Steiner, A. L., and Cohen, R. C.: Temperature and Recent Trends in the Chemistry of Continental Surface Ozone, *Chemical Reviews*, 115, 3898–3918, 2015.
- Rasmussen, D. J., Hu, J., Mahmud, A., and Kleeman, M. J.: The Ozone–Climate Penalty: Past, Present, and Future, *Environmental Science & Technology*, 47, 14258–14266, 2013.

- Rickard, A., Young, J., Pilling, M. J., Jenkin, M. E., Pascoe, S., and Saunders, S. M.: The Master Chemical Mechanism Version MCM v3.2, <http://mcm.leeds.ac.uk/MCMv3.2/>, [Online; accessed 25-March-2015], 2015.
- Rubin, J. I., Kean, A. J., Harley, R. A., Millet, D. B., and Goldstein, A. H.: Temperature dependence of volatile organic compound evaporative emissions from motor vehicles, *Journal of Geophysical Research: Atmospheres*, 111, d03305, 2006.
- 5 Sadanaga, Y., Yoshino, A., Kato, S., , and Kajii, Y.: Measurements of OH Reactivity and Photochemical Ozone Production in the Urban Atmosphere, *Environmental Science & Technology*, 39, 8847–8852, pMID: 16323785, 2005.
- Sander, R., Kerkweg, A., Jöckel, P., and Lelieveld, J.: Technical note: The new comprehensive atmospheric chemistry module MECCA, *Atmospheric Chemistry and Physics*, 5, 445–450, 2005.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM  
10 v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmospheric Chemistry and Physics*, 3, 161–180, 2003.
- Schnell, J. L., Prather, M. J., Josse, B., Naik, V., Horowitz, L. W., Cameron-Smith, P., Bergmann, D., Zeng, G., Plummer, D. A., Sudo, K., Nagashima, T., Shindell, D. T., Faluvegi, G., and Strode, S. A.: Use of North American and European air quality networks to evaluate global chemistry–climate modeling of surface ozone, *Atmospheric Chemistry and Physics*, 15, 10 581–10 596, 2015.
- 15 Sillman, S.: The use of NO<sub>y</sub>, H<sub>2</sub>O<sub>2</sub>, and HNO<sub>3</sub> as indicators for ozone-NO<sub>x</sub>-hydrocarbon sensitivity in urban locations, *Journal of Geophysical Research: Atmospheres*, 100, 14 175–14 188, 1995.
- Sillman, S.: The relation between ozone, NO<sub>x</sub> and hydrocarbons in urban and polluted rural environments, *Atmospheric Environment*, 33, 1821 – 1845, 1999.
- Sillman, S. and Samson, P. J.: Impact of temperature on oxidant photochemistry in urban, polluted rural and remote environments, *Journal*  
20 *of Geophysical Research: Atmospheres*, 100, 11 497–11 508, 1995.
- Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Flechard, C. R., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena, V. S., Tsyro, S., Tuovinen, J.-P., Valdebenito, Á., and Wind, P.: The EMEP MSC-W chemical transport model – technical description, *Atmospheric Chemistry and Physics*, 12, 7825–7865, 2012.
- Solberg, S., Hov, Ø., Søvde, A., Isaksen, I. S. A., Coddeville, P., De Backer, H., Forster, C., Orsolini, Y., and Uhse, K.: European surface  
25 ozone in the extreme summer 2003, *Journal of Geophysical Research: Atmospheres*, 113, D07307, 2008.
- Steiner, A. L., Tonse, S., Cohen, R. C., Goldstein, A. H., and Harley, R. A.: Influence of future climate and emissions on regional air quality in California, *Journal of Geophysical Research: Atmospheres*, 111, d18303, 2006.
- Stockwell, W. R., Middleton, P., Chang, J. S., and Tang, X.: The second generation regional acid deposition model chemical mechanism for regional air quality modeling, *Journal of Geophysical Research: Atmospheres*, 95, 16 343–16 367, 1990.
- 30 Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S.: A new mechanism for regional atmospheric chemistry modeling, *Journal of Geophysical Research: Atmospheres*, 102, 25 847–25 879, 1997.
- Vautard, R., Honoré, C., Beekmann, M., and Rouil, L.: Simulation of ozone during the August 2003 heat wave and emission control scenarios, *Atmospheric Environment*, 39, 2957 – 2967, 2005.
- von Schneidemesser, E., Monks, P. S., Allan, J. D., Bruhwiler, L., Forster, P., Fowler, D., Lauer, A., Morgan, W. T., Paasonen, P., Righi, M., Sindelarova, K., and Sutton, M. A.: Chemistry and the Linkages between Air Quality and Climate Change, *Chemical Reviews*, pMID: 25926133, 2015.
- 40 von Schneidemesser, E., Coates, J., Visschedijk, A. J. H., Denier van der Gon, H. A. C., and Butler, T. M.: Variation of the NMVOC speciation in the solvent sector and the sensitivity of modelled tropospheric ozone, *Atmospheric Environment*, Submitted for Publication, 2016.

**Figure 1.** The estimated isoprene emissions (molecules isoprene cm<sup>-2</sup> s<sup>-1</sup>) using MEGAN2.1 at each temperature used in the study.



**Table 1.** Total AVOC emissions in 2011 in tonnes from each anthropogenic source category assigned from TNO-MACC\_III emission inventory and temperature-independent BVOC emissions in tonnes from Benelux region assigned from EMEP. The allocation of these emissions to MCMv3.2, CRIV2, CB05, MOZART-4 and RADM2 species are found in the supplementary material.

Source Category	Total Emissions	Source Category	Total Emissions
Public Power	13755	Road Transport: Diesel	6727
Residential Combustion	21251	Road Transport: Others	1433
Industry	62648	Road Transport: Evaporation	2327
Fossil Fuel	15542	Non-road Transport	17158
Solvent Use	100826	Waste	1342
Road Transport: Gasoline	24921	BVOC	10702

Wagner, P. and Kuttler, W.: Biogenic and anthropogenic isoprene in the near-surface urban atmosphere — A case study in Essen, Germany, Science of The Total Environment, 475, 104 – 115, 2014.

Wang, J., Chew, C., Chang, C.-Y., Liao, W.-C., Lung, S.-C. C., Chen, W.-N., Lee, P.-J., Lin, P.-H., and Chang, C.-C.: Biogenic isoprene in subtropical urban settings and implications for air quality, Atmospheric Environment, 79, 369–379, 2013.

5 Yarwood, G., Rao, S., Yocke, M., and Whitten, G. Z.: Updates to the Carbon Bond Chemical Mechanism: CB05, Tech. rep., U. S Environmental Protection Agency, 2005.

**Table 2.** Increase in mean ozone mixing ratio (ppbv) due to chemistry (i.e. faster reaction rates) and temperature-dependent isoprene emissions from 20 °C to 40 °C in the NO<sub>x</sub>-regimes of Fig. 3.


Chemical	Source of	Increase in Ozone from 20 °C to 40 °C (ppbv)		
Mechanism	Difference	Low-NO <sub>x</sub>	Maximal-O <sub>3</sub>	High-NO <sub>x</sub>
MCMv3.2	Isoprene Emissions	4.6	7.7	10.6
	Chemistry	6.8	12.5	15.2
CRIV2	Isoprene Emissions	4.8	7.9	10.8
	Chemistry	6.0	11.1	13.7
MOZART-4	Isoprene Emissions	4.1	6.7	10.0
	Chemistry	6.0	10.2	12.3
CB05	Isoprene Emissions	4.6	7.4	9.8
	Chemistry	9.3	16.0	19.9
RADM2	Isoprene Emissions	3.8	5.7	7.8
	Chemistry	8.6	14.1	17.3

**Table 3.** Slopes ( $m_{O_3-T}$ , ppbv per °C) of the linear fit to MDA8 values of ozone and temperature correlations in Fig. 6, indicating the increase of MDA8 in ppbv of ozone per °C. The slope of the observational data is 2.15 ppbv/°C and the slope of the WRF-Chem output is 2.05 ppbv/°C.


Mechanism	Isoprene Emissions	Low-NO <sub>x</sub>		Maximal-O <sub>3</sub>		High-NO <sub>x</sub>	
		Mixing	No Mixing	Mixing	No Mixing	Mixing	No Mixing
MCMv3.2	Temperature Independent	0.28	1.01	0.51	1.36	0.59	0.96
	Temperature Dependent	0.42	1.48	0.74	2.16	0.93	2.63
CRIV2	Temperature Independent	0.25	0.93	0.47	1.27	0.55	0.88
	Temperature Dependent	0.40	1.44	0.71	2.09	0.90	2.52
MOZART-4	Temperature Independent	0.25	0.97	0.44	1.21	0.49	0.59
	Temperature Dependent	0.38	1.43	0.65	1.98	0.81	2.05
CB05	Temperature Independent	0.39	1.30	0.67	1.72	0.79	1.45
	Temperature Dependent	0.52	1.72	0.89	2.44	1.12	2.94
RADM2	Temperature Independent	0.37	1.31	0.61	1.64	0.70	1.28
	Temperature Dependent	0.48	1.68	0.79	2.22	0.97	2.49




**Figure 2.** Contours of peak ozone mixing ratios (ppbv) as a function of the total  $\text{NO}_x$  emissions and temperature for each chemical mechanism using a temperature-dependent and temperature-independent source of isoprene emissions. The contours can be split into three separate regimes: High- $\text{NO}_x$ , Maximal- $\text{O}_3$  and Low- $\text{NO}_x$  indicated in the figure.



**Figure 3.** Mean ozone mixing ratios (ppbv) at each temperature after allocation to the different NO<sub>x</sub>-regimes of Fig. 2. The differences in ozone mixing ratios due to chemistry (solid line) and isoprene emissions (dotted line) are represented graphically for MOZART-4 with High-NO<sub>x</sub> conditions. Table 2 details the differences for each chemical mechanism and NO<sub>x</sub>-condition.




**Figure 4.** Day-time budgets of  $O_x$  normalised by the total loss rate of emitted VOC in the  $NO_x$ -regimes of Fig. 3. The white line indicates net production or consumption of  $O_x$ . The net contribution of reactions to  $O_x$  budgets are allocated to categories of inorganic reactions, peroxy nitrates (RO<sub>2</sub>NO<sub>2</sub>), reactions of NO with HO<sub>2</sub>, alkyl peroxy radicals (RO<sub>2</sub>) and acyl peroxy radicals (ARO<sub>2</sub>). All other reactions are allocated to the ‘Other Organic’ category.




**Figure 5.** Day-time budgets of  $O_x$  allocated to the  $NO_x$ -regimes allocated to the net contribution of reactions to  $O_x$  budgets are allocated to categories of inorganic reactions, peroxy nitrates ( $RO_2NO_2$ ), reactions of  $NO$  with  $HO_2$ , alkyl peroxy radicals ( $RO_2$ ) and acyl peroxy radicals ( $ARO_2$ ). All other reactions are allocated to the 'Other Organic' category.

(a)  $O_x$  production and consumption budgets with box model setup including mixing.



(b)  $O_x$  production and consumption budgets with box model setup without including mixing.



**Figure 6.** MDA8 values of ozone from the box model simulations allocated to the different  $\text{NO}_x$  regimes for each chemical mechanism with mixing (solid lines) and without mixing (dashed lines). The box model ozone-temperature correlation is compared to the summer 2007 observational data (black circles) and WRF-Chem output (purple boxes).

