Institute for Advanced Sustainability Studies (IASS)

Understanding Ozone Pollution:

A Comparison of Chemical Mechanisms

Jane Coates and Tim Butler

Motivation

- Importance of O_3 production chemistry representation – future emission scenarios.
- Compare different O_3 chemistry representations used in chemical transport models.
- Determine effects on O_3 production by comparing treatment of Volatile Organic Compounds (VOCs) degradation products.

Approach

- Tagged Ozone Production Potentials (TOPPs) [1] calculated over 7 days for VOCs common to urban environments.
- Following mechanisms are compared to near-explicit MCM v3.2.

MCM v3.1	CRI v2	CBM-IV	CB05
RADM2	RACM	RACM2	MOZART-4

- O_x (= $O_3 + NO_2$) production allocated to emitted VOC by 'tagging' its organic degradation products.

Results







Conclusions

- More explicit mechanisms produce more O_x than less explicit mechanisms.
- VOCs broken down into smaller fragments quicker in less-explicit mechanisms resulting in less O_x production.
- First day O_x production from VOCs similar between many mechanisms, larger differences over time.
- Differences in VOC degradation treatments impacts on O_x production – RACM aromatic chemistry.

References

[1] T. M. Butler, M. G. Lawrence, D. Taraborrelli, and J. Lelieveld. Multi-day ozone production potential of volatile organic compounds calculated with a tagging approach. Atmospheric Environment, 45(24):4082–4090, 2011.

The IASS is sponsored by

SPONSORED BY THE
Federal Ministry of Education and Research
FONA Research for Sustainable Development
BMBF
LAND BRANDENBURG
Ministerium für Wissenschaft, Forschung und Kultur