

Corrected proposed-versus-control decisions by evidence scope

		OPSD lead 1 fixed	OPSD lead 24 fixed	OPSD lead 24 rolling	SimBench lead 1 fixed	SimBench lead 24 fixed	Ausgrid lead 24 OLS
	MLP	loss Holm $p = 0.03477$	win Holm $p = 0.008531$	not run	n.s. Holm $p = 1$	n.s. Holm $p = 0.08412$	n.s. Holm $p = 0.6894$
	Target-self context	not run	not run	n.s. Holm $p = 0.9844$	not run	not run	not run
	TemporalOnly (smaller head)	n.s. Holm $p = 0.1882$	win Holm $p = 0.001096$	not run	n.s. Holm $p = 0.2709$	n.s. Holm $p = 1$	n.s. Holm $p = 1$
	Uniform/equal-weight neighbors	n.s. Holm $p = 1$	n.s. Holm $p = 1$	n.s. Holm $p = 0.9844$	n.s. Holm $p = 1$	n.s. Holm $p = 1$	n.s. Holm $p = 1$
	Euclidean weights	n.s. Holm $p = 1$	n.s. Holm $p = 1$	n.s. Holm $p = 0.9844$	n.s. Holm $p = 1$	n.s. Holm $p = 1$	n.s. Holm $p = 1$
	Fixed distance scale	n.s. Holm $p = 1$	n.s. Holm $p = 1$	n.s. Holm $p = 0.0625$	n.s. Holm $p = 1$	n.s. Holm $p = 1$	n.s. Holm $p = 1$

Fixed-split cells use seed-level Mann-Whitney tests; rolling cells use exact sign-flip tests over 8 origin means.