

Note:- Formula

i) $(A')' = A$

ii) $(kA)' = kA'$

iii) $(A \pm B)' = A' \pm B'$

iv) $(AB)' = B'A'$

3.5.1 Properties of transpose of the matrices

We now state the following properties of transpose of matrices without proof. These may be verified by taking suitable examples.

For any matrices A and B of suitable orders, we have

- (i) $(A')' = A$, (ii) $(kA)' = kA'$ (where k is any constant)
- (iii) $(A + B)' = A' + B'$ (iv) $(AB)' = B'A'$

Example 20 If $A = \begin{bmatrix} 3 & \sqrt{3} & 2 \\ 4 & 2 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & -1 & 2 \\ 1 & 2 & 4 \end{bmatrix}$, verify that

- (i) $(A')' = A$, (ii) $(A + B)' = A' + B'$,
- (iii) $(kB)' = kB'$, where k is any constant.

$$\begin{aligned} (A - B)' &= (A + (-B))' \\ &= A' + (-B)' \\ &= A' + (-1)B' \\ &= A' - B' \end{aligned}$$

Solution

- (i) We have

Symmetric matrix

$$B = B'$$

we say B is
a symmetric matrix.

$$B'A' = \begin{bmatrix} 1 \\ 3 \\ -6 \end{bmatrix} \begin{bmatrix} -2 & 4 & 5 \end{bmatrix} = \begin{bmatrix} -2 & 4 & 5 \\ -6 & 12 & 15 \\ 12 & -24 & -30 \end{bmatrix} = (AB)'$$

Clearly $(AB)' = B'A'$

3.6 Symmetric and Skew Symmetric Matrices

Definition 4 A square matrix $A = [a_{ij}]$ is said to be *symmetric* if $A' = A$, that is, $[a_{ij}] = [a_{ji}]$ for all possible values of i and j .

For example $A = \begin{bmatrix} \sqrt{3} & 2 & 3 \\ 2 & -1.5 & -1 \\ 3 & -1 & 1 \end{bmatrix}$ is a symmetric matrix as $A' = A$

Definition 5 A square matrix $A = [a_{ij}]$ is said to be *skew symmetric* matrix if $A' = -A$, that is $a_{ji} = -a_{ij}$ for all possible values of i and j . Now, if we put $i = j$, we have $a_{ii} = -a_{ii}$. Therefore $2a_{ii} = 0$ or $a_{ii} = 0$ for all i 's.

This means that all the diagonal elements of a skew symmetric matrix are zero.

Reprint 2025-26

Skew-symmetric matrix

$$B' = -B$$

$B \rightarrow$ skew-symm.

Note ① Skew-symm

$$B' = -B$$

$$\text{Let } B = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$B' = \begin{bmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \end{bmatrix}$$

$$B' = -B$$

$$\begin{bmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \end{bmatrix} = \begin{bmatrix} -a_{11} & -a_{12} \\ -a_{21} & -a_{22} \end{bmatrix}$$

$$a_{11} = -a_{11} \quad \& \quad a_{22} = -a_{22}$$

3.6 Symmetric and Skew Symmetric Matrices

Definition 4 A square matrix $A = [a_{ij}]$ is said to be *symmetric* if $A' = A$, that is, $[a_{ij}] = [a_{ji}]$ for all possible values of i and j .

For example $A = \begin{bmatrix} \sqrt{3} & 2 & 3 \\ 2 & -1.5 & -1 \\ 3 & -1 & 1 \end{bmatrix}$ is a symmetric matrix as $A' = A$

Definition 5 A square matrix $A = [a_{ij}]$ is said to be *skew symmetric* matrix if $A' = -A$, that is $a_{ji} = -a_{ij}$ for all possible values of i and j . Now, if we put $i = j$, we have $a_{ii} = -a_{ii}$. Therefore $2a_{ii} = 0$ or $a_{ii} = 0$ for all i 's.

This means that all the diagonal elements of a skew symmetric matrix are zero.

Reprint 2025-26

$$\begin{bmatrix} -a_{11} & -a_{12} \\ -a_{21} & -a_{22} \end{bmatrix}$$

MATHEMATICS

For example, the matrix $B = \begin{bmatrix} 0 & e & f \\ -e & 0 & g \\ -f & -g & 0 \end{bmatrix}$ is a skew symmetric matrix as $B' = -B$

$$2a_{11} = 0 \quad \& \quad 2a_{22} = 0$$

$$a_{11} = 0 \quad \& \quad a_{22} = 0$$

$$B = \begin{bmatrix} 0 & a_{12} \\ a_{21} & 0 \end{bmatrix}$$

diagonal elements

diagonal elements of a skew-symm matrix are zero



Search



ENG IN

7:50 PM
5/4/2026

Note 2

$A \rightarrow$ Square matrix

$A + A' \rightarrow$ Symmetric matrix

$A - A' \rightarrow$ Skew-symmetric matrix

Now, we are going to prove some results on symmetric and skew-symmetric matrices.

Theorem 1 For any square matrix A with real number entries, $A + A'$ is a symmetric matrix and $A - A'$ is a skew symmetric matrix.

Proof Let $B = A + A'$, then

$$\begin{aligned} B' &= (A + A')' \\ &= A' + (A')' \text{ (as } (A + B)' = A' + B') \\ &= A' + A \text{ (as } (A')' = A) \\ &= A + A' \text{ (as } A + B = B + A) \\ &= B \end{aligned}$$

Therefore

$B = A + A'$ is a symmetric matrix

Now let

$$C = A - A'$$

$$\begin{aligned} C' &= (A - A')' = A' - (A')' \quad (\text{Why?}) \\ &= A' - A \quad (\text{Why?}) \\ &= -(A - A') = -C \end{aligned}$$

Therefore

$C = A - A'$ is a skew symmetric matrix.

Theorem 2 Any square matrix can be expressed as the sum of a symmetric and a skew symmetric matrix.

Proof Let A be a square matrix, then we can write

$$A = \frac{1}{2}(A + A') + \frac{1}{2}(A - A')$$

From the Theorem 1, we know that $(A + A')$ is a symmetric matrix and $(A - A')$ is a skew symmetric matrix. Since for any matrix A , $(kA)' = kA'$, it follows that $\frac{1}{2}(A + A')$

Note 3 Any square matrix A can be represented as the sum of symmetric and skew-symmetric matrices

Let $A = B + C$ where $B \rightarrow \text{symmetric} \quad (B' = B)$
 $C \rightarrow \text{skew-symmetric} \quad (C' = -C)$

$$B = \frac{1}{2}(A + A') \quad \text{and} \quad C = \frac{1}{2}(A - A')$$

$$B' = \frac{1}{2}(A + A')'$$

$$B' = \frac{1}{2}(A' + (A')')$$

$$B' = \frac{1}{2}(A' + A)$$

$$B' = B$$

$\Rightarrow B \rightarrow \text{symmetric matrix}$

$$C' = \frac{1}{2}(A' - (A')')$$

$$C' = \frac{1}{2}(A' - A) = -\frac{1}{2}(A - A')$$

$$C' = -C$$

$C \rightarrow \text{skew-symmetric matrix}$

$$\underline{A} = B + C$$

Here, $B = \frac{1}{2}(A + A')$ $\rightarrow$ Symmetric

$$C = \frac{1}{2}(A - A') \rightarrow \text{Skew-Symmetric}$$

Example 22 Express the matrix $B = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$ as the sum of a symmetric and a skew symmetric matrix.

sol

$$B = P + Q$$

where

$$P = \frac{1}{2} (B + B')$$

→ Symmetric

$$Q = \frac{1}{2} (B - B')$$

→ Skew-Symmetric

$$B' = \begin{bmatrix} 2 & -1 & 1 \\ -2 & 3 & -2 \\ -4 & 4 & -3 \end{bmatrix}$$

$$P = \frac{1}{2} \left(\begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix} + \begin{bmatrix} 2 & -1 & 1 \\ -2 & 3 & -2 \\ -4 & 4 & -3 \end{bmatrix} \right) = \frac{1}{2} \begin{bmatrix} 4 & -3 & -3 \\ -3 & 6 & 2 \\ -3 & 2 & -6 \end{bmatrix}$$

$$P = \begin{bmatrix} 2 & -3/2 & -3/2 \\ -3/2 & 3 & 1 \\ -3/2 & 1 & -3 \end{bmatrix}$$

$$Q = \frac{1}{2} (B - B')$$

$$P + Q = \begin{bmatrix} \quad \end{bmatrix} + \begin{bmatrix} \quad \end{bmatrix} = B$$



Edit Convert E-Sign

Find text or tools Share Ask AI Assistant

Q6

$$A' = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

$$A' A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \alpha + \sin^2 \alpha & \sin \alpha \cos \alpha - \sin \alpha \cos \alpha \\ \sin \alpha \cos \alpha - \sin \alpha \cos \alpha & \sin^2 \alpha + \cos^2 \alpha \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

Hand proved

4. If $A' = \begin{bmatrix} -2 & 3 \\ 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix}$, then find $(A + 2B)'$

5. For the matrices A and B, verify that $(AB)' = B'A'$, where

(i) $A = \begin{bmatrix} 1 \\ -4 \\ 3 \end{bmatrix}$, $B = \begin{bmatrix} -1 & 2 & 1 \end{bmatrix}$ (ii) $A = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 5 & 7 \end{bmatrix}$

6. If (i) $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, then verify that $A' A = I$

(ii) If $A = \begin{bmatrix} \sin \alpha & \cos \alpha \\ -\cos \alpha & \sin \alpha \end{bmatrix}$, then verify that $A' A = I$

7. (i) Show that the matrix $A = \begin{bmatrix} 1 & -1 & 5 \\ -1 & 2 & 1 \\ 5 & 1 & 3 \end{bmatrix}$ is a symmetric matrix.

(ii) Show that the matrix $A = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$ is a skew symmetric matrix.

8. For the matrix $A = \begin{bmatrix} 1 & 5 \\ 6 & 7 \end{bmatrix}$, verify that

- (i) $(A + A')$ is a symmetric matrix
- (ii) $(A - A')$ is a skew symmetric matrix



Edit Convert E-Sign

Find text or tools



Share

Ask AI Assistant

Q7 (ii)

$$A' = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

$$-A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

$$A' = -A$$

$A \rightarrow$ Skew-symm

If $A' = \begin{bmatrix} -2 & 3 \\ 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix}$, then find $(A + 2B)'$

For the matrices A and B, verify that $(AB)' = B'A'$, where

(i) $A = \begin{bmatrix} 1 \\ -4 \\ 3 \end{bmatrix}$, $B = \begin{bmatrix} -1 & 2 & 1 \end{bmatrix}$ (ii) $A = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 5 & 7 \end{bmatrix}$

6. If (i) $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, then verify that $A' A = I$

(ii) If $A = \begin{bmatrix} \sin \alpha & \cos \alpha \\ -\cos \alpha & \sin \alpha \end{bmatrix}$, then verify that $A' A = I$

7. (i) Show that the matrix $A = \begin{bmatrix} 1 & -1 & 5 \\ -1 & 2 & 1 \\ 5 & 1 & 3 \end{bmatrix}$ is a symmetric matrix.

(ii) Show that the matrix $A = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$ is a skew symmetric matrix.

8. For the matrix $A = \begin{bmatrix} 1 & 5 \\ 6 & 7 \end{bmatrix}$, verify that

(i) $(A + A')$ is a symmetric matrix

(ii) $(A - A')$ is a skew symmetric matrix

Q11

$$A' = A$$

$$B' = B$$

} → given

$$(iii) \begin{bmatrix} 3 & 3 & -1 \\ -2 & -2 & 1 \\ -4 & -5 & 2 \end{bmatrix}$$

$$(iv) \begin{bmatrix} 1 & 5 \\ -1 & 2 \end{bmatrix}$$

Choose the correct answer in the Exercises 11 and 12.

11. If A, B are symmetric matrices of same order, then AB - BA is a

(A) Skew symmetric matrix

(B) Symmetric matrix

(C) Zero matrix

(D) Identity matrix

12. If $A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$, and $A + A' = I$, then the value of α is

(A) $\frac{\pi}{6}$

(B) $\frac{\pi}{3}$

(C) π

(D) $\frac{3\pi}{2}$

find about this matrix

$$(AB - BA)' = (AB)' - (BA)'$$

$$= B'A' - A'B'$$

$$= BA - AB$$

$$(AB - BA)' = -(AB - BA)$$

3.7 Invertible Matrices

Definition 6 If A is a square matrix of order m , and if there exists another square matrix B of the same order m , such that $AB = BA = I$, then B is called the *inverse* matrix of A and it is denoted by A^{-1} . In that case A is said to be invertible.

For example, let

$$A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} \text{ be two matrices.}$$

$$AB = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 4-3 & -6+6 \\ 2-2 & -3+4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

Now

Edit Convert E-Sign

Find text or tools

Share

Ask AI Assistant

Invertible matrix

$A, B \rightarrow$ square matrix of same order

such that

$$AB = BA = I$$

then

$$B = A^{-1} \neq \frac{1}{A}$$

or

$$A = B^{-1}$$

(A) $\frac{\pi}{6}$

(B) $\frac{\pi}{3}$

(C) π

(D) $\frac{3\pi}{2}$

3.7 Invertible Matrices

Definition 6 If A is a square matrix of order m , and if there exists another square matrix B of the same order m , such that $AB = BA = I$, then B is called the inverse matrix of A and it is denoted by A^{-1} . In that case A is said to be invertible.

For example, let

$$A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} \text{ be two matrices.}$$

Now

$$\begin{aligned} AB &= \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 4-3 & -6+6 \\ 2-2 & -3+4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I \end{aligned}$$

$$BA = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I. \text{ Thus } B \text{ is the inverse of } A, \text{ in other}$$

Also words $B = A^{-1}$ and A is inverse of B , i.e., $A = B^{-1}$

Q2

$$B'AB \rightarrow \text{Symm}$$

$$A \rightarrow \text{Symm}$$

(a) $A' = A$

$$(B'AB)' = B'AB$$

Miscellaneous Exercise on Chapter 3

1. If A and B are symmetric matrices, prove that $AB - BA$ is a skew symmetric matrix.
2. Show that the matrix $B'AB$ is symmetric or skew symmetric according as A is symmetric or skew symmetric.

3. Find the values of x, y, z if the matrix $A = \begin{bmatrix} 0 & 2y & z \\ x & y & -z \\ x & -y & z \end{bmatrix}$ satisfy the equation

$$A'A = I$$

4. For what values of x: $\begin{bmatrix} 1 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \\ x \end{bmatrix} = O$?

5. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = O$.

6. Find x, if $\begin{bmatrix} x & -5 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ 1 \end{bmatrix} = O$

7. A manufacturer produces three products x, y, z which he sells in two markets. Annual sales are indicated below:

(b) $A' = -A$

$$(B'AB)' = -B'AB$$

(b) $A' = -A$ (given)

To show $(B'AB)' = -B'AB$

Proof

$$\begin{aligned}(B'AB)' &= (B'(AB))' \\ &= (AB)'(B')' \\ &= B'A'B \\ &= B'(-A)B \\ (B'AB)' &= -B'AB\end{aligned}$$

Hence Proved

Formula

$$(AB)' = B'A'$$

$$AB \neq BA$$

$$A(BC) = (AB)C$$


$$(a) \quad A' = A \text{ (given)}$$

To Prove

$$(B' A B)' = B' A B$$

Proof

$$(B' \underbrace{AB})' = \underbrace{(B')}_I (\underbrace{AB})_{II}'$$

$$= (AB)' (B')'$$

$$= B' A' B$$

$$(B' A B)' = B' A B$$

$$B' A B \rightarrow \text{symmetric}$$



Edit Convert E-Sign

Find text or tools

Share

Ask AI Assistant

Q10

$$A' = A$$

$$A' = -A$$

$$2A' = 0$$

$$A' = 0$$

$$A = 0$$

(a) If unit sale prices of x , y and z are ₹ 2.50, ₹ 1.50 and ₹ 1.00, respectively, find the total revenue in each market with the help of matrix algebra.

(b) If the unit costs of the above three commodities are ₹ 2.00, ₹ 1.00 and 50 paise respectively. Find the gross profit.

8. Find the matrix X so that $X \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} = \begin{bmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{bmatrix}$

Choose the correct answer in the following questions:

9. If $A = \begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ is such that $A^2 = I$, then

(A) $1 + \alpha^2 + \beta\gamma = 0$

(B) $1 - \alpha^2 + \beta\gamma = 0$

(C) $1 - \alpha^2 - \beta\gamma = 0$

(D) $1 + \alpha^2 - \beta\gamma = 0$

10. If the matrix A is both symmetric and skew symmetric, then

(A) A is a diagonal matrix

(B) A is a zero matrix

(C) A is a square matrix

(D) None of these

11. If A is square matrix such that $A^2 = A$, then $(I + A)^3 - 7A$ is equal to

(A) A

(B) $I - A$

(C) I

(D) $3A$

Summary

- ◆ A matrix is an ordered rectangular array of numbers or functions.
- ◆ A matrix having m rows and n columns is called a matrix of order $m \times n$.
- ◆ $[a_{ij}]_{m \times 1}$ is a column matrix.
- ◆ $[a_{ij}]_{1 \times n}$ is a row matrix.
- ◆ An $m \times n$ matrix is a square matrix if $m = n$.
- ◆ $A = [a_{ij}]_{m \times m}$ is a diagonal matrix if $a_{ij} = 0$, when $i \neq j$.
- ◆ $A = [a_{ij}]_{n \times n}$ is a scalar matrix if $a_{ij} = 0$, when $i \neq j$, $a_{ii} = k$, (k is some

Q11

$$A^2 = A$$

$$(I + A)^3 - 7A$$

$$I^3 + A^3 + 3I^2A + 3IA^2 - 7A$$

$$I^2 = I^3 = I^4 = I$$

$$I + A \cdot A^2 + 3IA + 3IA - 7A$$

$$I + A \cdot A + 3A + 3A - 7A$$

$$I + A^2 - A$$

$$I + A - A = I$$

8. Find the matrix X so that $X \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} = \begin{bmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{bmatrix}$

Sol

2×2

2×3

2×3

X $\rightarrow$ 2×2 matrix

$$X = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} = \begin{bmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{bmatrix}$$

$$\begin{bmatrix} a+4b & 2a+5b & 3a+6b \\ c+4d & 2c+5d & 3c+6d \end{bmatrix} = \begin{bmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{bmatrix}$$

$$a + 4b = -7$$

$$2a + 5b = -8$$

$$c + 4d = 2$$

$$2c + 5d = 4$$

64 PM
/2026

Example 24 If A and B are symmetric matrices of the same order, then show that AB is symmetric if and only if A and B commute, that is $AB = BA$.

Sol Given

$$A' = A$$

$$B' = B$$

$$AB = BA$$

To Prove

$$(AB)' = AB$$

Proof

$$(AB)' = B' A'$$

$$= BA$$

$$(AB)' = AB$$

1. If A and B are symmetric matrices, prove that $AB - BA$ is a skew symmetric matrix.

Mis
Q1

Given

$$A' = A$$

$$B' = B$$

To prove

$$(AB - BA)' = -(AB - BA)$$

Proof

$$(AB - BA)' = (AB)' - (BA)'$$

$$= B'A' - A'B'$$

$$= BA - AB$$

$$(AB - BA)' = -(AB - BA) \Rightarrow$$

$AB - BA \rightarrow$ Skew symm.

Example 23 If $A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$, then prove that $A^n = \begin{bmatrix} \cos n\theta & \sin n\theta \\ -\sin n\theta & \cos n\theta \end{bmatrix}$, $n \in \mathbb{N}$.

Sol

$$A^n = \begin{bmatrix} \cos n\theta & \sin n\theta \\ -\sin n\theta & \cos n\theta \end{bmatrix}$$

Put $n=1$

$$A^1 = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

It is true for $n=1$

Let us assume it is true for $n=k$

$$A^k = \begin{bmatrix} \cos k\theta & \sin k\theta \\ -\sin k\theta & \cos k\theta \end{bmatrix}$$

we shall prove it for $n=k+1$

$$A^{k+1} = \begin{bmatrix} \cos (k+1)\theta & \sin (k+1)\theta \\ -\sin (k+1)\theta & \cos (k+1)\theta \end{bmatrix}$$

Proof

L.H.S

A^{k+1}

$A^k \cdot A$

$$\begin{bmatrix} \cos k\theta & \sin k\theta \\ -\sin k\theta & \cos k\theta \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$\begin{bmatrix} \cos K\theta & \sin K\theta \\ -\sin K\theta & \cos K\theta \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$\begin{bmatrix} \cos K\theta \cos \theta - \sin K\theta \sin \theta \\ -\sin K\theta \cos \theta - \sin \theta \cos K\theta \\ -(\sin K\theta \cos \theta + \sin \theta \cos K\theta) \end{bmatrix}$$

$$\begin{bmatrix} \cos (K\theta + \theta) \\ -\sin (K\theta + \theta) \end{bmatrix}$$

$$\begin{bmatrix} \cos (k+1)\theta \\ -\sin (k+1)\theta \end{bmatrix}$$

$$\begin{bmatrix} \cos K\theta \sin \theta + \sin K\theta \cos \theta \\ -\sin K\theta \sin \theta + \cos K\theta \cos \theta \end{bmatrix}$$

$$\begin{bmatrix} \sin (\theta + K\theta) \\ \cos (K\theta + \theta) \end{bmatrix}$$

$$\begin{bmatrix} \sin (k+1)\theta \\ \cos (k+1)\theta \end{bmatrix}$$

= R.H.S

Hence, it is true for $n \in \mathbb{N}$.

$$\begin{aligned} \cos (A+B) \\ = \cos A \cos B - \sin A \sin B \end{aligned}$$

$$\begin{aligned} \sin (A+B) \\ = \sin A \cos B + \cos A \sin B \end{aligned}$$

H.W

Ex 3.3 → Q4, Q6, Q7, Q10 (ii), (iii), Q11, Q12

Ex 3.4 → Q1

Example → 23, 24, 25

Misc. Exercise → Q1 to Q11

By wed

respectively. Similarly, in the second arrangement, the entries in the first row represent the number of notebooks possessed by Radha, Fauzia and Simran, respectively. The entries in the second row represent the number of pens possessed by Radha, Fauzia and Simran, respectively. An arrangement or display of the above kind is called a *matrix*. Formally, we define matrix as:

Definition 1 A *matrix* is an ordered rectangular array of numbers or functions. The numbers or functions are called the elements or the entries of the matrix.

We denote matrices by capital letters. The following are some examples of matrices:

a_{ij} row column

$$A = \begin{bmatrix} -2 & 5 \\ 0 & \sqrt{5} \\ 3 & 6 \end{bmatrix}, B = \begin{bmatrix} 2+i & 3 & -\frac{1}{2} \\ 3.5 & -1 & 2 \\ \sqrt{3} & 5 & \frac{5}{7} \end{bmatrix}, C = \begin{bmatrix} 1+x & x^3 & 3 \\ \cos x & \sin x + 2 & \tan x \end{bmatrix}$$

In the above examples, the horizontal lines of elements are said to constitute, *rows* of the matrix and the vertical lines of elements are said to constitute, *columns* of the matrix. Thus A has 3 rows and 2 columns, B has 3 rows and 3 columns while C has 2 rows and 3 columns.

3.2.1 Order of a matrix

A matrix having m rows and n columns is called a matrix of *order* $m \times n$ or simply $m \times n$ matrix (read as an m by n matrix). So referring to the above examples of matrices, we have A as 3×2 matrix, B as 3×3 matrix and C as 2×3 matrix. We observe that A has $3 \times 2 = 6$ elements, B and C have 9 and 6 elements, respectively.

$$a_{11} = -2$$

$$a_{12} = 5$$

$$a_{21} = 0$$

$$a_{22} = \sqrt{5}$$

$$a_{13} = \text{DNE}$$

$$a_{31} = 3$$

$$a_{32} = 6$$



Edit Convert E-Sign

Find text or tools Share Ask AI Assistant

$$A = [a_{ij}]_{m \times n}$$

a_{ij} = element of i^{th} row

and j^{th} column

$$1 \leq i \leq m$$

and

$$1 \leq j \leq n$$

of the matrix and the vertical lines of elements are said to constitute, columns of the matrix. Thus A has 3 rows and 2 columns, B has 3 rows and 3 columns while C has 2 rows and 3 columns.

3.2.1 Order of a matrix

A matrix having m rows and n columns is called a matrix of order $m \times n$ or simply $m \times n$ matrix (read as an m by n matrix). So referring to the above examples of matrices, we have A as 3×2 matrix, B as 3×3 matrix and C as 2×3 matrix. We observe that A has $3 \times 2 = 6$ elements, B and C have 9 and 6 elements, respectively.

In general, an $m \times n$ matrix has the following rectangular array:

1st row $\leftarrow$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2j} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & & \vdots & & \vdots \\ a_{i1} & a_{i2} & a_{i3} & \dots & a_{ij} & \dots & a_{in} \\ \vdots & \vdots & \vdots & & \vdots & & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}_{m \times n}$$

m^{th} row $\rightarrow$

or $A = [a_{ij}]_{m \times n}, 1 \leq i \leq m, 1 \leq j \leq n, i, j \in \mathbb{N}$

Thus the i^{th} row consists of the elements $a_{i1}, a_{i2}, a_{i3}, \dots, a_{in}$, while the j^{th} column consists of the elements $a_{1j}, a_{2j}, a_{3j}, \dots, a_{mj}$.

In general a_{ij} is an element lying in the i^{th} row and j^{th} column. We can also call it as the $(i, j)^{\text{th}}$ element of A. The number of elements in an $m \times n$ matrix will be equal to mn .

Total num of elements in matrix A = mn

$$i, j \in \mathbb{N}$$

$m \times n$ order
 m = no of rows
 n = no of column

Imp Note.

Reprint 2025-26

$A = \begin{bmatrix} 0 & 2 & 3 & 2 \end{bmatrix}_{2 \times 4}$ or $I = C \begin{bmatrix} 1 & 3 \\ -1 & 2 \end{bmatrix}_{4 \times 2}$

Thus, matrices can be used as representation of vertices of geometrical figures in a plane.
Now, let us consider some examples.

Example 1 Consider the following information regarding the number of men and women workers in three factories I, II and III

	Men workers	Women workers
I	30	25
II	25	31
III	27	26

Represent the above information in the form of a 3×2 matrix. What does the entry in the third row and second column represent?

Reprint 2025-26

Solution The information is represented in the form of a 3×2 matrix as follows:

$\begin{bmatrix} 30 & 25 \end{bmatrix}$

Eg1

$A = \begin{bmatrix} 30 \\ 25 \\ 27 \end{bmatrix}$
↑
men worker

$\begin{bmatrix} 25 \\ 31 \\ 26 \end{bmatrix}$
↑
women
 3×2

2×3 matrix

$A = \begin{bmatrix} 30 & 25 & 27 \\ 25 & 31 & 26 \end{bmatrix}$
↑ ↑ ↑
I II III
← men
← women
 2×3

Eg²

$$mn = 8$$

m	n	Total	
1	8	8	1×8
2	4	8	2×4
4	2	8	4×2
8	1	8	8×1

$$A = \begin{bmatrix} 30 & 25 \\ 25 & 31 \\ 27 & 26 \end{bmatrix}$$

The entry in the third row and second column represents the number of women workers in factory III.

Example 2 If a matrix has 8 elements, what are the possible orders it can have?

Solution: We know that if a matrix is of order $m \times n$, it has mn elements. Thus, to find all possible orders of a matrix with 8 elements, we will find all ordered pairs of natural numbers, whose product is 8.

Thus, all possible ordered pairs are $(1, 8), (8, 1), (4, 2), (2, 4)$

Hence, possible orders are $1 \times 8, 8 \times 1, 4 \times 2, 2 \times 4$

Example 3 Construct a 3×2 matrix whose elements are given by $a_{ij} = \frac{1}{2}|i - 3j|$.

Solution In general a 3×2 matrix is given by $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$

$$a_{ij} = \frac{1}{2} |i - 3j|, i = 1, 2, 3 \text{ and } j = 1, 2.$$

Therefore $a_{11} = \frac{1}{2} |1 - 3 \times 1| = 1$ $a_{12} = \frac{1}{2} |1 - 3 \times 2| = \frac{5}{2}$

$$a_{21} = \frac{1}{2} |2 - 3 \times 1| = \frac{1}{2} \quad a_{22} = \frac{1}{2} |2 - 3 \times 2| = 2$$

$$a_{31} = \frac{1}{2} |3 - 3 \times 1| = 0 \quad a_{32} = \frac{1}{2} |3 - 3 \times 2| = \frac{3}{2}$$

1. In the matrix $A = \begin{bmatrix} 35 & -2 & \frac{5}{2} & 12 \\ \sqrt{3} & 1 & -5 & 17 \end{bmatrix}$, write:

(i) The order of the matrix, (ii) The number of elements,

(iii) Write the elements $a_{13}, a_{21}, a_{33}, a_{24}, a_{23}$.

2. If a matrix has 24 elements, what are the possible orders it can have? What, if it has 13 elements?

3. If a matrix has 18 elements, what are the possible orders it can have? What, if it has 5 elements?

4. Construct a 2×2 matrix, $A = [a_{ij}]$, whose elements are given by:

(i) $a_{ij} = \frac{(i+j)^2}{2}$ (ii) $a_{ij} = \frac{i}{j}$ (iii) $a_{ij} = \frac{(i+2j)^2}{2}$

5. Construct a 3×4 matrix, whose elements are given by:

(i) $a_{ij} = \frac{1}{2}|-3i+j|$ (ii) $a_{ij} = 2i-j$

6. Find the values of x, y and z from the following equations:

(i) $\begin{bmatrix} 4 & 3 \\ x & 5 \end{bmatrix} = \begin{bmatrix} y & z \\ 1 & 5 \end{bmatrix}$ (ii) $\begin{bmatrix} x+y & 2 \\ 5+z & xy \end{bmatrix} = \begin{bmatrix} 6 & 2 \\ 5 & 8 \end{bmatrix}$ (iii) $\begin{bmatrix} x+y+z \\ x+z \\ y+z \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \\ 7 \end{bmatrix}$

7. Find the value of a, b, c and d from the equation:

$$\begin{bmatrix} a-b & 2a+c \\ 2a-b & 3c+d \end{bmatrix} = \begin{bmatrix} -1 & 5 \\ 0 & 13 \end{bmatrix}$$

Q2

1×24

2×12

3×8

4×6

6×4

8×3

12×2

24×1

1×13

13×1



Edit Convert E-Sign

Find text or tools 🔍 | 🗂️ | 📄 | 🖨️ | Share | Ask AI Assistant

Eg 3

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

↑ ↑
3 × 2

$$a_{ij} = \frac{1}{2} |i - 3j|$$

$$a_{11} = \frac{1}{2} |1 - 3(1)|$$

$$= \frac{1}{2} |2| = \frac{2}{2} = 1$$

Solution We know that if a matrix is of order $m \times n$, it has mn elements. Thus, to find all possible orders of a matrix with 8 elements, we will find all ordered pairs of natural numbers, whose product is 8.

Thus, all possible ordered pairs are (1, 8), (8, 1), (4, 2), (2, 4)

Hence, possible orders are $1 \times 8, 8 \times 1, 4 \times 2, 2 \times 4$

Example 3 Construct a 3×2 matrix whose elements are given by $a_{ij} = \frac{1}{2} |i - 3j|$.

Solution In general a 3×2 matrix is given by $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$


Now $a_{ij} = \frac{1}{2} |i - 3j|, i = 1, 2, 3$ and $j = 1, 2$.

Therefore $a_{11} = \frac{1}{2} |1 - 3 \times 1| = 1$ $a_{12} = \frac{1}{2} |1 - 3 \times 2| = \frac{5}{2}$

$a_{21} = \frac{1}{2} |2 - 3 \times 1| = \frac{1}{2}$ $a_{22} = \frac{1}{2} |2 - 3 \times 2| = 2$

$a_{31} = \frac{1}{2} |3 - 3 \times 1| = 0$ $a_{32} = \frac{1}{2} |3 - 3 \times 2| = \frac{3}{2}$

Hence the required matrix is given by $A = \begin{bmatrix} 1 & \frac{5}{2} \\ \frac{1}{2} & 2 \\ 0 & \frac{3}{2} \end{bmatrix}$ ✓



Edit Convert E-Sign

Find text or tools



Share



Ask AI Assistant



8

42

Eg

$$x + 3 = 0$$

$$\Rightarrow \boxed{x = -3}$$

$$z + 4 = 6$$

$$\boxed{z = 2}$$

$$2y - 7 = 3y - 2$$

$$-7 + 2 = 3y - 2y$$

$$\boxed{y = -5}$$

A matrix is said to be *zero matrix* or *null matrix* if all its elements are zero.

For example, $[0]$, $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $[0, 0]$ are all zero matrices. We denote zero matrix by O . Its order will be clear from the context.

3.3.1 Equality of matrices

Definition 2 Two matrices $A = [a_{ij}]$ and $B = [b_{ij}]$ are said to be equal if

- (i) they are of the same order
- (ii) each element of A is equal to the corresponding element of B , that is $a_{ij} = b_{ij}$ for all i and j .

For example, $\begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix}$ are equal matrices but $\begin{bmatrix} 3 & 2 \\ 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix}$ are not equal matrices. Symbolically, if two matrices A and B are equal, we write $A = B$.

If $\begin{bmatrix} x & y \\ z & a \\ b & c \end{bmatrix} = \begin{bmatrix} -1.5 & 0 \\ 2 & \sqrt{6} \\ 3 & 2 \end{bmatrix}$, then $x = -1.5$, $y = 0$, $z = 2$, $a = \sqrt{6}$, $b = 3$, $c = 2$

Example 4 If $\begin{bmatrix} x+3 & z+4 & 2y-7 \\ -6 & a-1 & 0 \\ b-3 & -21 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 6 & 3y-2 \\ -6 & -3 & 2c+2 \\ 2b+4 & -21 & 0 \end{bmatrix}$

Find the values of a , b , c , x , y and z .

Solution As the given matrices are equal, therefore, their corresponding elements must be equal. Comparing the corresponding elements, we get

$$\begin{array}{lll} x+3=0, & z+4=6, & 2y-7=3y-2 \\ a-1=-3, & 0=2c+2 & b-3=2b+4, \end{array}$$

$$a-1 = -3$$

$$\boxed{a = -2}$$

$$0 = 2c + 2$$

$$\boxed{c = -1}$$

$$b-3 = 2b+4$$

$$-3-4 = 2b-b$$

$$\boxed{b = -7}$$

3

Search

ENG
IN3:47 PM
4/26/2026

Q9

$$3x + 7 = 0$$

$$x = -7/3$$

$$2 - 3x = 4$$

$$-3x = 2$$

$$x = -2/3$$

8. $A = [a_{ij}]_{m \times n}$ is a square matrix, if
 (A) $m < n$ (B) $m > n$ (C) $m = n$ (D) None of these
9. Which of the given values of x and y make the following pair of matrices equal

$$\begin{bmatrix} 3x+7 & 5 \\ y+1 & 2-3x \end{bmatrix}, \begin{bmatrix} 0 & y-2 \\ 8 & 4 \end{bmatrix}$$

 (A) $x = \frac{-1}{3}, y = 7$ (B) Not possible to find
 (C) $y = 7, x = \frac{-2}{3}$ (D) $x = \frac{-1}{3}, y = \frac{-2}{3}$
10. The number of all possible matrices of order 3×3 with each entry 0 or 1 is:
 (A) 27 (B) 18 (C) 81 (D) 512

3.4 Operations on Matrices

In this section, we shall introduce certain operations on matrices, namely, addition of matrices, multiplication of a matrix by a scalar, difference and multiplication of matrices.

3.4.1 Addition of matrices

Suppose Fatima has two factories at places A and B. Each factory produces sport shoes for boys and girls in three different price categories labelled 1, 2 and 3. The quantities produced by each factory are represented as matrices given below:

10. The number of all possible matrices of order 3×3 with each entry 0 or 1 is:
(A) 27 (B) 18 (C) 81 (D) 512

89

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3}$$

$$2 \times 2 \times 2 \times \dots \text{9 times}$$
$$2^9 = 512$$

$$\begin{aligned} a_{11} &\rightarrow 0 \text{ or } 1 \\ a_{12} &\rightarrow 0 \text{ or } 1 \\ &\vdots \\ a_{33} &\rightarrow 0 \text{ or } 1 \end{aligned}$$

$$(1) \quad a_{ij} = \frac{-1}{2} | -5l + j | \quad (11) \quad a_{ij} = 2l - j$$

6. Find the values of x , y and z from the following equations:

$$\begin{array}{lll} \text{(i)} \quad \begin{bmatrix} 4 & 3 \\ x & 5 \end{bmatrix} = \begin{bmatrix} y & z \\ 1 & 5 \end{bmatrix} & \text{(ii)} \quad \begin{bmatrix} x+y & 2 \\ 5+z & xy \end{bmatrix} = \begin{bmatrix} 6 & 2 \\ 5 & 8 \end{bmatrix} & \text{(iii)} \quad \begin{bmatrix} x+y+z \\ x+z \\ y+z \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \\ 7 \end{bmatrix} \end{array}$$

7. Find the value of a , b , c and d from the equation:

$$\begin{bmatrix} a-b & 2a+c \\ 2a-b & 3c+d \end{bmatrix} = \begin{bmatrix} -1 & 5 \\ 0 & 13 \end{bmatrix}$$

$$2a + c = 5$$

$$2(1) + c = 5$$

$$C = 3$$

$$-a = -1$$
$$a = 1$$

$$1 - b = -1$$
$$1 + 1 = b \Rightarrow \boxed{b = 2}$$

8. $A = [a_{ij}]_{m \times n}$ is a square matrix, if

H.W

Example - 4 and 5

+

Exercise 3.1 → Q2, Q3, Q5, Q6, Q7, Q9, Q10
└────────→ Tuesday (6:30)

+

You need to solve C & D Test again
└────────────────────────→ Friday

Q7 Q10

$$2x + 3y = \begin{bmatrix} 2 & 3 \\ 4 & 0 \end{bmatrix} \quad \text{--- ①}$$

$$3x + 2y = \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \quad \text{--- ②}$$

multiply ① with 3 and ② with 2

$$6x + 9y = \begin{bmatrix} 6 & 9 \\ 12 & 0 \end{bmatrix}$$

5. If $A = \begin{bmatrix} 1 & 2 & 4 \\ 3 & 3 & 3 \\ 7 & 2 & 2 \\ 3 & 3 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 4 \\ 5 & 5 & 5 \\ 7 & 6 & 2 \\ 5 & 5 & 5 \end{bmatrix}$, then compute $3A - 5B$.

6. Simplify $\cos \theta \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} + \sin \theta \begin{bmatrix} \sin \theta & -\cos \theta \\ \cos \theta & \sin \theta \end{bmatrix}$

7. Find X and Y, if

(i) $X + Y = \begin{bmatrix} 7 & 0 \\ 2 & 5 \end{bmatrix}$ and $X - Y = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$

(ii) $2X + 5Y = \begin{bmatrix} 2 & 3 \\ 4 & 0 \end{bmatrix}$ and $3X + 2Y = \begin{bmatrix} 2 & -2 \\ -1 & 5 \end{bmatrix}$

8. Find X, if $Y = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$ and $2X + Y = \begin{bmatrix} 1 & 0 \\ -3 & 2 \end{bmatrix}$

9. Find x and y, if $2 \begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$

10. Solve the equation for x, y, z and t, if $2 \begin{bmatrix} x & z \\ y & t \end{bmatrix} + 3 \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} = 3 \begin{bmatrix} 3 & 5 \\ 4 & 6 \end{bmatrix}$

11. If $x \begin{bmatrix} 2 \\ 3 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 10 \\ 5 \end{bmatrix}$, find the values of x and y.


12. Given $3 \begin{bmatrix} x & y \\ z & w \end{bmatrix} = \begin{bmatrix} x & 6 \\ -1 & 2w \end{bmatrix} + \begin{bmatrix} 4 & x+y \\ z+w & 3 \end{bmatrix}$, find the values of x, y, z and w.

meet.google.com

Online Maths Classes (You)

Raphael Rusten

Emmanuel Thomas



same
no. of column of
first matrix = no. of
rows of
second matrix

Then E → Result matrix
will have order 2x2

52 MATHEMATICS

and is given by $CD = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & 4 \end{bmatrix} \begin{bmatrix} 2 & 7 \\ -1 & 1 \\ 5 & -4 \end{bmatrix}$. This is a 2×2 matrix in which each entry is the sum of the products across some row of C with the corresponding entries down some column of D. These four computations are

Entry in first row first column $\rightarrow \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix} = \begin{bmatrix} (1)(2) + (-1)(-1) + (2)(5) \\ ? \end{bmatrix}$

Entry in first row second column $\rightarrow \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & 4 \end{bmatrix} \begin{bmatrix} 7 \\ 1 \\ -4 \end{bmatrix} = \begin{bmatrix} (1)(7) + (-1)(1) + 2(-4) \\ ? \end{bmatrix}$

Entry in second row first column $\rightarrow \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix} = \begin{bmatrix} 13 \\ 0(2) + 3(-1) + 4(5) \end{bmatrix}$

Entry in second row second column $\rightarrow \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & 4 \end{bmatrix} \begin{bmatrix} 7 \\ 1 \\ -4 \end{bmatrix} = \begin{bmatrix} 13 \\ 0(7) + 3(1) + 4(-4) \end{bmatrix}$

Thus $CD = \begin{bmatrix} 13 & -2 \\ 17 & -13 \end{bmatrix}$

$\begin{bmatrix} 13 & -2 \\ 17 & -13 \end{bmatrix}$

Rough

$$\begin{array}{r} 2 + 1 + 10 \\ = 13 \\ \hline 7 - 1 - 8 = -2 \\ \hline 0 - 3 + 20 = 17 \\ \hline 0 + 3 - 16 = -13 \end{array}$$

meet.google.com

Online Maths Classes

H.W

Example - 8, 10, 9, 14, 17, 18

Exercice - 3.2 ; Q2 (ii) & (iv)

Q3 (iv), (vi)

Q5, Q6, Q7 (ii)

Q10, Q11, Q12, Q15, Q16, Q17

+ CSD question Paper by Friday.

Q7 (ii)

$$\begin{aligned} 3A - 5B &= \begin{bmatrix} 2 & 3 & 5 \\ 1 & 2 & 4 \\ 7 & 6 & 2 \end{bmatrix} - \begin{bmatrix} 2 & 3 & 5 \\ 1 & 2 & 4 \\ 7 & 6 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$
$$\begin{aligned} 6. \cos \theta &\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} + \sin \theta \begin{bmatrix} \sin \theta & -\cos \theta \\ \cos \theta & \sin \theta \end{bmatrix} \\ &= \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ -\sin \theta \cos \theta & \cos^2 \theta \end{bmatrix} + \begin{bmatrix} \sin^2 \theta & -\sin \theta \cos \theta \\ \sin \theta \cos \theta & \sin^2 \theta \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$
$$\begin{aligned} 7(i) \quad 2x + 3y &= \begin{bmatrix} 2 & 3 \\ 4 & 0 \end{bmatrix} \quad (1) \\ 3x + 2y &= \begin{bmatrix} 2 & -2 \\ -1 & 5 \end{bmatrix} \quad (2) \end{aligned}$$

Multiply (1) by -2 and (2) by 3

$$\begin{aligned} -4x - 6y &= \begin{bmatrix} -4 & -6 \\ -8 & 0 \end{bmatrix} \quad (3) \\ 9x + 6y &= \begin{bmatrix} 6 & -6 \\ -3 & 15 \end{bmatrix} \quad (4) \end{aligned}$$
$$5x = \begin{bmatrix} 2 & -12 \\ -11 & -15 \end{bmatrix}$$

$$X = \begin{bmatrix} 2/5 & -12/5 \\ -11/5 & 3 \end{bmatrix}$$

$\rightarrow +15$

$$A^2 = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$$

$$A^3 = A^2 \cdot A = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix} \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$$

Rough

$$5A = 5 \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 10 & 0 & 5 \\ 10 & 5 & 15 \\ 5 & -5 & 0 \end{bmatrix}$$

$$6I = \begin{bmatrix} 6 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{bmatrix}$$

$$A^2 - 5A + 6I,$$

$$\begin{bmatrix} 5 & -1 & 3 \\ 9 & -2 & 5 \\ 0 & -1 & -2 \end{bmatrix} - \begin{bmatrix} 10 & 0 & 5 \\ 10 & 5 & 15 \\ 5 & -5 & 0 \end{bmatrix} + \begin{bmatrix} 6 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & -1 & -2 \\ -1 & -7 & -10 \\ -5 & 4 & 4 \end{bmatrix}$$

$$16. A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$$

$$A^3 = A^2 \cdot A$$

$$= \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1(1)+0(2)+2(2) & 1(0)+0(2)+2(0) & 1(2)+0(1)+2(3) \\ 0(1)+2(2)+1(2) & 0(0)+2(2)+1(0) & 0(2)+2(1)+1(3) \\ 2(1)+0(3)+3(2) & 2(0)+0(2)+3(0) & 2(2)+0(1)+3(3) \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 0 & 8 \\ 6 & 4 & 5 \\ 8 & 0 & 13 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 5 & 0 & 8 \\ 4 & 4 & 5 \\ 0 & 0 & 13 \end{bmatrix}$$

✓. prob

13. If $F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$, show that $F(x) F(y) = F(x+y)$. ✓

$$F(x+y) = \begin{bmatrix} \cos(x+y) & -\sin(x+y) & 0 \\ \sin(x+y) & \cos(x+y) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

sol

$$F(y) = \begin{bmatrix} \cos y & -\sin y & 0 \\ \sin y & \cos y & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$F(x) F(y) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos y & -\sin y & 0 \\ \sin y & \cos y & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$F(x) F(y) = \begin{bmatrix} \cos x \cos y - \sin x \sin y & -\cos x \sin y - \sin x \cos y & 0 \\ \sin x \cos y + \cos x \sin y & -\sin x \sin y + \cos x \cos y & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$F(x) F(y) = \begin{bmatrix} \cos x \cos y - \sin x \sin y & -\cos x \sin y - \sin x \cos y & 0 \\ \sin x \cos y + \cos x \sin y & -\sin x \sin y + \cos x \cos y & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$F(x)F(y) = \begin{bmatrix} \cos(x+y) & -[\cos x \sin y + \sin x \cos y] & 0 \\ \sin(x+y) & \cos(x+y) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$\nearrow \sin(x+y)$

$$= \begin{bmatrix} \cos(\underline{x+y}) & -\sin(\underline{x+y}) & 0 \\ \sin(\underline{x+y}) & \cos(\underline{x+y}) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$F(x) F(y) = F(x+y)$$

$$\begin{aligned} \sin(A+B) &= \sin A \cos B + \cos A \sin B \end{aligned}$$

$$\begin{aligned} \cos(A+B) &= \cos A \cos B - \sin A \sin B \end{aligned}$$

18. If $A = \begin{bmatrix} 0 & -\tan \frac{\alpha}{2} \\ \tan \frac{\alpha}{2} & 0 \end{bmatrix}$ and I is the identity matrix of order 2, show that

$$I + A = (I - A) \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

sol
L.H.S

$$I + A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & -\tan \alpha/2 \\ \tan \alpha/2 & 0 \end{bmatrix} = \begin{bmatrix} 1 & -\tan \alpha/2 \\ \tan \alpha/2 & 1 \end{bmatrix} \quad \underline{\underline{=}}$$

R.H.S

$$(I - A) \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

R.H.S

$$= \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & -\tan \alpha/2 \\ \tan \alpha/2 & 0 \end{bmatrix} \right) \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

$$= \begin{bmatrix} 1 & \tan \alpha/2 \\ -\tan \alpha/2 & 1 \end{bmatrix} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

R.H.S

$$= \begin{bmatrix} 1 & \tan \alpha/2 \\ -\tan \alpha/2 & 1 \end{bmatrix} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

R.H.S

$$\begin{bmatrix} 1 & \tan \alpha/2 \\ -\tan \alpha/2 & 1 \end{bmatrix} \begin{bmatrix} \frac{1 - \tan^2 \alpha/2}{1 + \tan^2 \alpha/2} & -\frac{2 \tan \alpha/2}{1 + \tan^2 \alpha/2} \\ \frac{2 \tan \alpha/2}{1 + \tan^2 \alpha/2} & \frac{1 - \tan^2 \alpha/2}{1 + \tan^2 \alpha/2} \end{bmatrix}$$

let $\tan \alpha/2 = t$

$$\begin{bmatrix} 1 & t \\ -t & 1 \end{bmatrix} \begin{bmatrix} \frac{1-t^2}{1+t^2} & \frac{-2t}{1+t^2} \\ \frac{2t}{1+t^2} & \frac{1-t^2}{1+t^2} \end{bmatrix}$$

$$\begin{bmatrix} \frac{1-t^2}{1+t^2} + \frac{2t^2}{1+t^2} & \frac{-2t}{1+t^2} + \frac{t(1-t^2)}{1+t^2} \\ \frac{-t(1-t^2)}{1+t^2} + \frac{2t}{1+t^2} & \frac{2t^2}{1+t^2} + \frac{1-t^2}{1+t^2} \end{bmatrix}$$

$$\sin 2A = \frac{2 \tan A}{1 + \tan^2 A}$$

$$\cos 2A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$$

RHS

$$\left[\begin{array}{cc} \frac{1-t^2}{1+t^2} + \frac{2t^2}{1+t^2} & \frac{-2t}{1+t^2} + \frac{t(1-t^2)}{1+t^2} \\ \frac{-t(1-t^2)}{1+t^2} + \frac{2t}{1+t^2} & \frac{2t^2}{1+t^2} + \frac{1-t^2}{1+t^2} \end{array} \right]$$

$$\left[\begin{array}{cc} 1 & \frac{-t(1+t^2)}{1+t^2} \\ \frac{t(1+t^2)}{1+t^2} & 1 \end{array} \right]$$

$$\left[\begin{array}{cc} \frac{1+t^2}{1+t^2} & \frac{-2t+t-t^3}{1+t^2} \\ \frac{-t+t^3+2t}{1+t^2} & \frac{2t^2+1-t^2}{1+t^2} \end{array} \right]$$

$$\left[\begin{array}{cc} 1 & -t \\ t & 1 \end{array} \right]$$

$$t = \tan \alpha/2$$

$$\left[\begin{array}{cc} 1 & \frac{-t-t^3}{1+t^2} \\ \frac{t+t^3}{1+t^2} & \frac{t^2+1}{1+t^2} \end{array} \right]$$

$$\left[\begin{array}{cc} 1 & -\tan \alpha/2 \\ \tan \alpha/2 & 1 \end{array} \right]$$

$$L.H.S = R.H.S$$

hence proved

$$1800 = \frac{5x + 7 \times 30000 - 7x}{100}$$

$$180000 = -2x + 210000$$

$$2x = 30000$$

$$x = 15,000$$

Type I bond investment = $x = \text{Rs. } 15,000$

Type II bond investment = $30000 - x = \text{Rs. } 15000$

(b)

$$2000 = \frac{5x}{100} + \frac{(3000 - x)7}{100}$$

20. The bookshop of a particular school has 10 dozen chemistry books, 8 dozen physics books, 10 dozen economics books. Their selling prices are ₹ 80, ₹ 60 and ₹ 40 each respectively. Find the total amount the bookshop will receive from selling all the books using matrix algebra.

Sol

$$\text{No. of books} = \begin{bmatrix} 120 & 96 & 120 \end{bmatrix}_{1 \times 3}$$

$\uparrow$ chem $\uparrow$ Phy $\uparrow$ Eco

$$\text{s.p} = \begin{bmatrix} 80 \\ 60 \\ 40 \end{bmatrix}_{3 \times 1}$$

$\leftarrow$ chem
 $\leftarrow$ Phy
 $\leftarrow$ Eco

$$\text{Total amount} = \begin{bmatrix} 120 & 96 & 120 \end{bmatrix}_{1 \times 3} \begin{bmatrix} 80 \\ 60 \\ 40 \end{bmatrix}_{3 \times 1} = \begin{bmatrix} \quad \quad \quad \end{bmatrix}_{1 \times 1}$$

~~$\begin{bmatrix} 80 \\ 60 \\ 40 \end{bmatrix}_{3 \times 1} \begin{bmatrix} \quad \quad \quad \end{bmatrix}_{1 \times 3}$
 $\begin{bmatrix} \quad \quad \quad \end{bmatrix}_{3 \times 3}$~~

$\swarrow$ No. of books $\nwarrow$ s.p

021

P Y

$P \times K$ $3 \times K$

$K = 3$

P Y

$P \times 3$ 3×3

order of PY = $P \times 3$

MATRICES 61

20. The bookshop of a particular school has 10 dozen chemistry books, 8 dozen physics books, 10 dozen economics books. Their selling prices are ₹ 80, ₹ 60 and ₹ 40 each respectively. Find the total amount the bookshop will receive from selling all the books using matrix algebra.

Assume X, Y, Z, W and P are matrices of order $2 \times n$, $3 \times k$, $2 \times p$, $n \times 3$ and $p \times k$, respectively. Choose the correct answer in Exercises 21 and 22.

21. The restriction on n , k and p so that $PY + WY$ will be defined are:

(A) $k = 3, p = n$ (B) k is arbitrary, $p = 2$
 (C) p is arbitrary, $k = 3$ (D) $k = 2, p = 3$

22. If $n = p$, then the order of the matrix $7X - 5Z$ is:

(A) $p \times 2$ (B) $2 \times n$ (C) $n \times 3$ (D) $p \times n$

3.5. Transpose of a Matrix

In this section, we shall learn about transpose of a matrix and special types of matrices such as symmetric and skew symmetric matrices.

Definition 3 If $A = [a_{ij}]$ be an $m \times n$ matrix, then the matrix obtained by interchanging the rows and columns of A is called the *transpose* of A . Transpose of the matrix A is denoted by A' or (A^T) . In other words, if $A = [a_{ij}]_{m \times n}$, then $A' = [a_{ji}]_{n \times m}$. For example,

if $A = \begin{bmatrix} 3 & 5 \\ \sqrt{3} & 1 \\ 0 & -\frac{1}{5} \end{bmatrix}_{3 \times 2}$, then $A' = \begin{bmatrix} 3 & \sqrt{3} & 0 \\ 5 & 1 & -\frac{1}{5} \end{bmatrix}_{2 \times 3}$

3.5.1 Properties of transpose of the matrices

W Y

$n \times 3$ 3×3

order of $WY = n \times 3$

for addition of matrices
 PY & WY
 $P \times 3 = n \times 3$
 $\Rightarrow P = n$

20. The bookshop of a particular school has 10 dozen chemistry books, 8 dozen physics books, 10 dozen economics books. Their selling prices are ₹ 80, ₹ 60 and ₹ 40 each respectively. Find the total amount the bookshop will receive from selling all the books using matrix algebra.

Assume X, Y, Z, W and P are matrices of order $2 \times n$, $3 \times k$, $2 \times p$, $n \times 3$ and $p \times k$, respectively. Choose the correct answer in Exercises 21 and 22.

21. The restriction on n , k and p so that $PY + WY$ will be defined are:

- (A) $k = 3, p = n$
- (B) k is arbitrary, $p = 2$
- (C) p is arbitrary, $k = 3$
- (D) $k = 2, p = 3$

22. If $n = p$, then the order of the matrix $7X - 5Z$ is:

- (A) $p \times 2$
- (B) $2 \times n$
- (C) $n \times 3$
- (D) $p \times n$

3.5. Transpose of a Matrix

In this section, we shall learn about transpose of a matrix and special types of matrices such as symmetric and skew symmetric matrices.

Definition 3 If $A = [a_{ij}]$ be an $m \times n$ matrix, then the matrix obtained by interchanging the rows and columns of A is called the *transpose* of A . Transpose of the matrix A is denoted by A' or (A^T) . In other words, if $A = [a_{ij}]_{m \times n}$, then $A' = [a_{ji}]_{n \times m}$. For example,

if $A = \begin{bmatrix} 3 & 5 \\ \sqrt{3} & 1 \\ 0 & -\frac{1}{5} \end{bmatrix}_{3 \times 2}$, then $A' = \begin{bmatrix} 3 & \sqrt{3} & 0 \\ 5 & 1 & -\frac{1}{5} \end{bmatrix}_{2 \times 3}$

3.5.1 Properties of transpose of the matrices

Q22
 $n = p$

$7X - 5Z$
 $\downarrow \quad \downarrow$
 $2 \times n \quad 2 \times p$

$2 \times n = 2 \times p$

order of final matrix $7X - 5Z$ $= 2 \times n = 2 \times p$

H.W

EXERCISE 3.2 $\rightarrow$ Q22, Q21, Q20, Q19, Q18, Q13

Example - 19, 11

Sunday
1 PM