# Homework 1

## Air Quality Dataset Analysis

Gary Walsh - use19210998

## Introduction

The airquality dataset from the datasets package in R contains daily air quality measurements in New York, from May to September 1973. The dataset includes measurements of Ozone, Solar Radiation, Wind, and Temperature.

In this document, we will plot the relationship between Ozone levels and Wind speed. The data will be color-coded based on the month to provide additional insights.

### **Approach**

- 1. Dataset: We use the airquality dataset from the datasets package.
- 2. **Plot Type:** Scatter plot with color-coding for months and a smoother line to show the trend.
- 3. Descriptive Statistics: Mean Ozone levels and mean Wind speed.

#### **Descriptive Statistics**

Table 1 below presents some descriptive statistics for the airquality dataset.

Table 1: Descriptive Statistics for Ozone and Wind

Statistic	Ozone	Wind
Mean	42.12931	9.957516
Median	31.50000	9.700000
Std Dev	32.98788	3.523001
Min	1.00000	1.700000
Max	168.00000	20.700000

#### **Summary**


Ozone levels exhibit high variability with extreme values, suggesting significant fluctuations in air quality. Wind speeds are more stable, with measurements closely clustering around the mean.

#### **Plot**

The scatter plot in fig 1 below shows the relationship between Ozone levels and Wind speed. The points are color-coded based on the month, and a smoother line is added to illustrate the trend.

# **Ozone Levels vs Wind Speed**

Color-coded by Month



#### Interpretation of the Plot

1. Inverse Relationship: Ozone levels generally decrease as Wind speed increases, suggesting higher wind speeds help disperse Ozone.

#### 2. Seasonal Variation:

- May and June: Lower Ozone levels across various wind speeds.
- July and August: Higher Ozone levels at lower wind speeds, indicating severe Ozone pollution in mid-summer.
- September: Mixed data but overall lower Ozone levels as fall approaches.

# 2. Data Dispersion:

- High variability in Ozone levels at lower wind speeds, especially in July and August.
- Reduced variability and lower Ozone levels at higher wind speeds.