

Improving Catch Estimation Methods in Sparsely Sampled Mixed-Stock Fisheries.

Nick Grunloh, Edward Dick, Don Pearson, John Field, Marc Mangel

Abstract

Introduction

Context

Data

- Collection issues
 - funding => nature of sparsity
- Lay down goal modeling goal
 - mean
 - uncertainty

Methods

Data Generating Model

Something something heirarchical poisson model. Something something (Shelton, 2012).

For the purposes of accurately modeling not only species composition means, but also higher moments of the data, such as species composition variances, it is necessary to recognize model limitations with respect to over-dispersed data. Among the simplest models for count data are the poisson and binomial models. Both models are typically specified with a single degree of freedom for modeling all of the moments of the data, and thus they rely heavily on their respective data generating processes to accurately represent higher moments in the data. McCullagh and Nelder (1989, pg. 124) commiserate about the prevalence of


over-dispersed data in cluster sampling, and explain the numerous ways in which cluster sampling may result in over-dispersion.

Extending the Poisson and binomial models to deal with over-dispersion, typically involves adding additional parameters for the purpose of modeling higher moments of the data. The negative binomial distribution is often used as an over-dispersed extension of the poisson model, since it can be expressly written as an infite mixture of poisson distributions. While the beta-binomial model is typically used to as an over-dispersed extension of the binomial model.


An Example

To discern between these models we consider a small scale example of the Poisson, binomial, negative binomial, and beta-binomial models fit to the port sampling integer weight data from market category 250, in the Monterey port complex trawl fishery in 1990. (*anywillwork*) This stratum was visited 38 times by port samplers, collecting a total of 67 cluster samples, resulting in 344 model observations across 21 (*atleast;URCK*) unique species. For brevity we only consider for the top six species here (CLPR, BCAC, WDOW, BLGL, ARRA, BANK). These data are serially fit by each of the Poisson, binomial, negative binomial, and beta-binomial models, predictive species compositions are then derived from each model, plotted along side the observed species compositions, and metrics of model fits are compared.

95% Predictive HDI Model Comparison



Beta-Binomial Posterior Predictive Species Compositions



	Poisson	Binomial	Negative-Binomial	Beta-binomial
DIC WAIC	5675.25	6759.86	1301.51 1302.19	1261.00
$\log p(y M)$	5840.56	6939.74	-688.19	1261.30
	-2864.01	-3406.01		-650.49

- describe picture
- notice overdispersion
- improper variance model biases mean, beta-binomial is flexible to disentangle these effects.
- maths stuffs
 - mean function
 - variance; introduce ρ
- justify linear predictor/transition to priors

Heirarchical Priors

Prediction

Model Exploration & Averaging

Results

- General Products
- Degree of smoothing (heirarchical parameters)
- Posterior v. Current
 - Report degree of similarity
- Prediction v. Data
 - Report predictive accuracy

Conclusions

- General Math/Science
- Database Stuff
- Looking Forward
 - forecasting/hindcasting
 - * simple

- * timeseries models
- more computation faster
- * broader model exploration
- * broader spatial expansion

References

- [1] Gelman, A., Carlin, J. B., Stern, H. S., & Rubin, D. B. (2014). Bayesian data analysis (Vol. 2). Boca Raton, FL, USA: Chapman & Hall/CRC.
- [2] Hoeting, J. A., Madigan, D., Raftery, A. E., & Volinsky, C. T. (1999). Bayesian model averaging: a tutorial. *Statistical science*, 382-401.
- [3] McCullagh P. & Nelder, J.A. (1989). *Generalized Linear Models*, 2nd ed. London: Chapman and Hall.
- [4] Pearson, D.E., and Erwin, B. (1997). Documentation of California’s commercial market sampling data entry and expansion programs. NOAA Tech Memo. NOAA-TM-NMFS-SWFSC-240.
- [5] R Core Team (2015). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- [6] Rue H., Martino S., Lindgren F., Simpson D., Riebler A. (2013). R-INLA: Approximate Bayesian Inference using Integrated Nested Laplace Approximations. Trondheim, Norway. URL <http://www.r-inla.org/>.
- [7] Rue, H., Martino, S., & Chopin, N. (2009). Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the royal statistical society: Series b (statistical methodology)*, 71(2), 319-392.
- [8] Sen, A.R. (1984). Sampling commercial rockfish landings in California. NOAA Tech Memo. NOAA-TM-NMFS-SWFSC-45.
- [9] Sen AR. (1986). Methodological problems in sampling commercial rockfish landings. *Fish Bull.* 84: 409-421 .
- [10] Shelton, A. O., Dick, E. J., Pearson, D. E., Ralston, S., & Mangel, M. (2012). Estimating species composition and quantifying uncertainty in multispecies fisheries: hierarchical Bayesian models for stratified sampling protocols with missing data. *Canadian Journal of Fisheries and Aquatic Sciences*, 69(2), 231-246.