

₁ Chapter 1

₂ The Pella-Tomlinson Model

Abstract

4 Stock assessments often assume a two-parameter functional form (e.g., Beverton-Holt or
5 Ricker) for the expected recruitment produced by a given level of spawning output. [Mangel](#)
6 [et al. \(2013\)](#) and others have shown that biological reference points such as $\frac{F^*}{M}$ and $\frac{B^*}{B(0)}$
7 are largely determined by a single parameter (steepness) when using two-parameter rela-
8 tionships. These functions introduce strong correlations between reference points that are
9 pre-determined by the functional form, rather than a biological characteristic of the stock.
10 Mangel et al. note that use of a three-parameter stock-recruitment relationship allows for
11 independent estimation of these reference points. This research seeks to understand the
12 nature of biases in reference points resulting from fitting a two-parameter functional form
13 when the true relationship follows a three-parameter stock-recruitment relationship. This
14 work demonstrates the useful limits of misspecified two-parameter models, and suggests the
15 mechanisms of model failure which arise from mapping a three-dimensional parameter space
16 into two dimensions.

1 Introduction

The most fundamental model in modern fisheries management is the surplus-production model. These models focus on modeling population growth via nonlinear parametric ordinary differential equations (ODE). Key management quantities called reference points (RPs) are commonly derived from the ODE equilibrium equations and depend upon the parameterization of biomass production. Two-parameter forms of the production function have been shown to limit the theoretical domain of RPs (Mangel et al., 2013). The limited RP-space of two parameter models are a major source of model misspecification for RPs and thus induce bias in RP estimation. The behavior of RP estimation bias is not well understood and as a result often underappreciated. A metamodeling approach is developed here to describe RP biases and explore mechanisms of model failure under the most common two parameter models.

Data for a typical surplus-production model comes in the form of an index of abundance through time which is assumed to be proportional to the reproducing biomass for the population of interest. The index is often observed alongside a variety of other known quantities, but at a minimum, each observed index will be observed in the presence of some known catch for the period. Figure (3.13) shows the classic Namibian Hake dataset exemplifying the form.

Indices are assumed to have multiplicative log-normal errors, and thus the following observation model arises naturally,

$$I_t = qB_te^\epsilon \quad \epsilon \sim N(0, \sigma^2). \quad (1.1)$$

Above q is often referred to as the “catchability parameter”; it serves as the proportionality constant mapping between the observed index of abundance and biomass. σ^2 models residual variation. Biologically speaking q and σ^2 are often treated as nuisance parameters with the “biological parameters” entering the model through a process model on biomass.

Biomass is assumed to evolve as an ODE; in this case I focus on the following form

$$\frac{dB}{dt} = P(B(t); \boldsymbol{\theta}) - Z(t)B(t). \quad (1.2)$$




Figure 1.1: *left*: An index of abundance data, catch per unit effort (CPUE), for Namibian Hake from 1965 to 1987 (Hilborn & Mangel, 1997). *right*: The associated catch data for Namibian Hake over the same time period.

Here biomass is assumed to change in time by two processes, net production of biomass into the population, $P(B)$, and various sources of biomass removal, Z , from the population.

Firstly, the population grows through a production function, $P(B)$. Production in this setting is defined as the net biomass increase due to all reproduction and maturation processes. The production function is assumed to be a parametric (generally non-linear) function relating the current biomass of the population to an aggregate production of biomass.

Secondly, the population decreases as biomass is removed by various sources that are assumed to remove biomass linearly with biomass. Above, $Z(t)$, is an aggregate rate of removal. When the fishing rate, $F(t)$, is the only source of removal $Z(t) = F(t)$, however often models will also included other linear terms in $Z(t)$. Commonly the rate of “natural mortality”, M , is also included as an additional term so that $Z(t) = M + F(t)$.

From a management perspective a major goal of modeling is to accurately infer a quantity known as *maximum sustainable yield* (MSY). One could maximize simple yield at a particular moment in time (and only for that moment) by fishing all available biomass in that moment. This strategy is penny-wise but pound-foolish (not to mention ecologically devastating) since it doesn’t leave biomass in the population to reproduce in the future. We seek to fish in a way that allows (or even encourages) future productivity in the population. This is accomplished by maximizing the equilibrium level of catch over time. Equilibrium yield is considered by

replacing the steady state biomass (\bar{B}) in the assumed form for catch, so that $\bar{Y} = F\bar{B}(F)$, where $\bar{\cdot}$ indicates a value at steady state. MSY is found by maximizing $\bar{Y}(F)$ with respect to F , and F^* is the fishing rate at MSY. Going forward let $*$ decorate any value derived under the condition of MSY.

Fisheries are very often managed based upon reference points which serve as simplified heuristic measures of population behavior. The mathematical form of RPs depends upon the model assumptions through the production function. While a number of different RPs exist which describe the population in different (but related) ways, the most common RPs revolve around the concept of MSY (or robust ways of measuring MSY (Hilborn, 2010; Punt et al., 2016)). Here the focus is primarily on the RPs $\frac{B^*}{B(0)}$ and F^* ($\frac{F^*}{M}$ when appropriate) for their pervasive use in modern fisheries (Punt & Cope, 2019).

F^* is the afore mentioned fishing rate which results in MSY. $\frac{B^*}{B(0)}$ is the depletion of the stock at MSY. That is to say $\frac{B^*}{B(0)}$ describes the fraction of the unfished population biomass that will remain in the equilibrium at MSY. In general $F^* \in \mathbb{R}^+$ and $\frac{B^*}{B(0)} \in (0, 1)$, however under the under the assumption of a two parameter production function production models will be structurally unable to capture the full theoretical range of RPs.

Many of the most commonly used production functions depend only on two parameters. For example, the Schaefer model depends only on the biological parameters r and K , and limits RP inference so that under the Schaefer model $\left(F^*, \frac{B^*}{B(0)}\right) \in \left(\mathbb{R}^+, \frac{1}{2}\right)$. The two parameter Fox model (Fox Jr., 1970) limits $\left(F^*, \frac{B^*}{B(0)}\right) \in \left(\mathbb{R}^+, \frac{1}{e}\right)$. Similarly the two parameter Cushing (Cushing, 1971), Beverton-Holt (Beverton & Holt, 1957, BH) and Ricker (Ricker, 1954) production functions do not model the full theoretical space of RPs (Mangel et al., 2013; Yeakel & Mangel, 2015).

The bias-variance trade-off (Ramasubramanian & Singh, 2017) makes it clear that the addition of a third parameter in the production function will necessarily reduce estimation bias. However the utility of this bias reduction is still under debate because the particular mechanisms and behavior (direction and magnitude) of these biases for key management quantities are not fully understood or described. Lee et al. (2012) provides some evidence that estimation of productivity parameters are dependent on biomass contrast as well as model specification. Conn et al. (2010) comes to similar conclusions via calibration modeling

88 techniques. These studies indicate important factors that contribute to inferential failure.
 89 However they do not offer mechanisms of model failure, nor do their experimental designs
 90 allow for the control of different types of model misspecification.

91 In this study I consider the behavior of inference when index data are simulated from
 92 three parameter PT and Schnute production models, but the simulated data are fit using
 93 intentionally misspecified two parameter logistic or BH production models. The work begins
 94 with a derivation of RPs under the three parameter models. A method is then presented
 95 for generating simulation designs based on the parametric form of RPs which serves as a
 96 control on the nature of simulated model misspecification. Finally a Gaussian Process (GP)
 97 metamodel ([Gramacy, 2020](#)) is constructed for exploration and analysis of RP biases.

98 A key insight of this approach is that bias is considered broadly across RP-space to
 99 uncover patterns and correlations between RPs. The GP metamodel is explicit about trade-
 100 offs between RPs so as to inform the full utility of reducing bias, as well as to suggest
 101 mechanisms for understanding what causes bias. Further, the effect of contrast on estimation
 102 is considered together with model misspecification.

103 2 Methods

104 2.1 Pella-Tomlinson Model

The three parameter Pella-Tomlinson (PT) family has a convenient form that includes, among others ([Fox Jr., 1970](#); [Rankin & Lemos, 2015](#)), the logistic production function as a special case. PT production function is parameterized so that $\boldsymbol{\theta} = [r, K, \gamma]$ and the family takes the following form,

$$P_p(B; [r, K, \gamma]) = \frac{rB}{\gamma - 1} \left(1 - \left(\frac{B}{K} \right)^{(\gamma-1)} \right). \quad (1.3)$$

γ is a parameter which breaks PT out of the restrictive symmetry of the logistic curve. In general $\gamma \in (1, \infty)$, with the logistic model appearing in the special case of $\gamma = 2$, and the Fox model appearing as a limiting case as $\gamma \rightarrow 1$. The parameter r controls the maximum reproductive rate of the population in the absence of competition for resources (i.e. the slope of production function at the origin). K is the so called "carrying capacity" of the population. In this context the carrying capacity can be formally stated as steady state biomass in the absence of fishing (i.e. $\bar{B}(0) = K$). In Figure (3.14) PT recruitment is shown for a range of parameter values so as to demonstrate the various recruitment shapes that can be achieved by PT recruitment.

While the form of the PT curve produces some limitations (Fletcher, 1978), importantly the introduction of a third parameter allows enough flexibility to fully describe the space of reference points used in management. To see this, the reference points are analytically derived for the PT model below.

PT Reference Points

With $B(t)$ representing biomass at time t , under PT production, the dynamics of biomass are defined by the following ODE,

$$\frac{dB}{dt} = \frac{rB}{\gamma - 1} \left(1 - \left(\frac{B}{K} \right)^{\gamma-1} \right) - FB. \quad (1.4)$$

An expression for the equilibrium biomass is attained by setting Eq (3.16) equal to zero, and rearranging the resulting equation to solve for B . Thinking of the result as a function




Figure 1.2: The Pella-Tomlinson production function plotted across a variety of parameter values. The special cases of Logistic production is shown in black, and the left-leaning and right-leaning regimes are shown in blue and red respectively.

of F gives,

$$\bar{B}(F) = K \left(1 - \frac{F(\gamma - 1)}{r} \right)^{\frac{1}{\gamma-1}}. \quad (1.5)$$

At this point it is convenient to notice that $\bar{B}(0) = K$. The expression for B^* is given by evaluating Eq (3.17) at F^* . To get an expression for F^* , the equilibrium yield is maximized with respect to F ,

$$F^* = \operatorname{argmax}_F F \bar{B}(F). \quad (1.6)$$

In the case of PT production this maximization can be done analytically, by differentiating the equilibrium yield with respect to F as follows,

$$\frac{d\bar{Y}}{dF} = \bar{B}(F) + F \frac{d\bar{B}}{dF} \quad (1.7)$$

$$\frac{d\bar{B}}{dF} = -\frac{K}{r} \left(1 - \frac{F(\gamma - 1)}{r} \right)^{\frac{1}{\gamma-1}-1}. \quad (1.8)$$

Setting Eq (3.19) equal to 0, substituting $\bar{B}(F)$ and $\frac{d\bar{B}}{dF}$ by Equations (3.17) and (3.20) respectively, and solving for F produces the following expression for the fishing rate required to produce MSY,

$$F^* = \frac{r}{\gamma} \quad (1.9)$$

Plugging the above expression for F^* back into Eq (3.17) gives the following expression for biomass at MSY,

$$B^* = K \left(\frac{1}{\gamma} \right)^{\frac{1}{\gamma-1}}. \quad (1.10)$$

The above derived expressions for $\bar{B}(0)$, B^* , and F^* can then be used to build a specific analytical form for the biological reference points in terms of only productivity parameters.

$$F^* = \frac{r}{\gamma} \qquad \frac{B^*}{\bar{B}(0)} = \left(\frac{1}{\gamma} \right)^{\frac{1}{\gamma-1}} \quad (1.11)$$

132 Simulation

Generating simulated indices of abundance from the PT model requires inverting the relationship between $\left(F^*, \frac{B^*}{B(0)}\right)$, and (r, γ) . It is not generally possible to analytically invert this relationship for many three parameter production functions (Punt & Cope, 2019; J. T. Schnute & Richards, 1998). Most three parameter production functions lead to RPs that require expensive numerical methods to invert; more over the numerical inversion procedure can often be unstable. That said, for the case of PT this relationship is analytically invertible, and leads to the following relationship

$$r = \gamma F^* \quad \gamma = \frac{W\left(\frac{B^*}{B(0)} \log\left(\frac{B^*}{B(0)}\right)\right)}{\log\left(\frac{B^*}{B(0)}\right)}. \quad (1.12)$$

133 Above W is the Lambert product logarithm function. More details about this derivation,
134 and the Lambert product logarithm, are given in Appendix (7).

135 Using Eq. (3.24) to obtain production parameters, a PT production model can be fully
136 defined for any combination of the RPs F^* and $\frac{B^*}{B(0)}$. Since K does not enter the RP
137 calculation its value is fixed arbitrarily at 10000.

138 Indices of abundance are simulated from the three parameter PT production model
139 broadly over the space of F^* and $\frac{B^*}{B(0)}$ via a space filling design as described in Section
140 (5 .3). A small amount of residual variation, $\sigma = 0.01$, is added to the simulated index, and
141 these data are then fit with a Schaefer model, at various degrees of misspecification, so as to
142 observe the effect of productivity model misspecification upon RP inference.

143 PT Design

144 Letting \mathcal{F} and \mathcal{B} be regular grids, of size $n = 100$, on $F^* \in (0.1, 0.7)$ and $\frac{B^*}{B_0} \in (0.2, 0.6)$
145 respectively, a LHS design of size 100 is collected among the cells produced by $\mathcal{F} \times \mathcal{B}$.

146 Each of the sampled LHS design locations represent a unique PT model with the sampled
147 RP values. Since the relationship mapping RPs analytically to productivity parameters can
148 be found for the PT model, LHS designs the the PT model are computed directly in RP
149 space and Eq. (3.24) is used to map the sampled RP design locations to PT productivity
150 parameters.

2.2 Gaussian Process Metamodel

At its core, a metamodel is simply a model of some mapping of inputs to outputs (the mapping itself is typically defined by a computer model). By modeling the mapping with a statistical model (that explicitly defines the relevant features of the mapping) a metamodel defines a specific ontology for the mapping. By simulating examples of the mapping, the inferential infrastructure of the statistical model is used to empirically learn an effective emulation of the mapping within the ontology defined by the statistical model. The predictive infrastructure of the statistical model is then useful as an approximate abstraction of the system itself to better understand the system through further data collection, cheap approximation of the mapping, and/or study of the mapping itself.

In this setting, the aim of metamodeling is to study how well RPs are inferred when typical two parameter models of productivity (Logistic and BH) are misspecified for populations that are actually driven by more complicated dynamics. The simulation design, \mathbf{X} , provides a sample of different population dynamics that are driven by three parameter production functions broadly in RP space. By simulating index of abundance data from the three parameter model, and fitting those data with the two parameter production model, we observe particular instances of how well RPs are inferred at the given misspecification of the two parameter model relative to the true three parameter production model. By gathering all of the simulated instances of how RPs are inferred (under the two parameter model), we form a set of example mappings to train a metamodel which represents the mapping of true RPs (under the three parameter model) to estimates of RPs under the misspecified two parameter production model. The metamodel is essentially a surrogate for inference under the misspecified two parameter production model that controls for the specific degree of model misspecification.

A flexible GP model is assumed for the structure of the metamodel to describe the mapping of RPs under misspecified two parameter models of productivity. A GP is a stochastic process generalizing the multivariate normal distribution to an infinite dimensional analog. GP models are often specified primarily through the choice of a covariance (or correlation) function which defines the relationship between locations in the input space. Typically corre-

180 lation functions are specified so that points closely related in space result in correlated effects
 181 in the model. In this setting the inputs to the GP metamodel are the space of reference points
 182 which define the simulated three parameter production models.

While index of abundance data are generated from three parameter models, at each design location of the simulation, fitting the restricted two parameter model results in a maximum likelihood estimate (MLE; and associated estimation uncertainty) of each of the productivity parameters (i.e. Schaefer: $[\log(r), \log(K)]$, BH: $[\log(\alpha), \log(\beta)]$). To simplify the specification of the metamodel, let \mathbf{y} be a vector collecting the fitted MLEs for one of the productivity parameters, and let $\boldsymbol{\omega}$ be a vector of estimates of the estimator variances (via the inverted Fisher information) at each \mathbf{y} . Each of the fitted productivity parameter estimates are then modeled using independent instances of the following GP metamodel.

$$\begin{aligned}\mathbf{y} &= \beta_0 + \mathbf{X}\boldsymbol{\beta} + \mathbf{v} + \boldsymbol{\epsilon} \\ \mathbf{v} &\sim N_n(\mathbf{0}, \tau^2 \mathbf{R}_\ell) \\ \boldsymbol{\epsilon} &\sim N_n(\mathbf{0}, \boldsymbol{\omega}' \mathbf{I})\end{aligned}\tag{1.13}$$

183 \mathbf{X} is the $n \times 2$ LHS design matrix of RPs for each simulated three parameter data
 184 generating model as described in Section (5.3). ϵ models independent normally distributed
 185 error, which provides an ideal mechanism for propagating uncertainty from inference in the
 186 simulation step into the metamodel. By matching each \mathbf{y}_i with an observed ω_i variance term,
 187 ϵ serves to down weight the influence of each \mathbf{y}_i in proportion to the inferred production model
 188 sampling distribution uncertainty. This has the effect of smoothing the GP model in a way
 189 similar to the nugget effect (Gramacy & Lee, 2012), although the application here models
 190 this effect heterogeneously.

The term, \mathbf{v} , contains spatially correlated GP effects. The correlation matrix, \mathbf{R}_ℓ describes how RPs close together in the simulation design are more correlated than those that are far away. This spatial effect is modeled with a squared exponential correlation function,

$$R(\mathbf{x}, \tilde{\mathbf{x}}) = \exp \left(\sum_{i=1}^2 \frac{-(x_i - \tilde{x}_i)^2}{2\ell_j^2} \right).\tag{1.14}$$

R has an anisotropic separable form which allows for differing length scales, ℓ_1 and ℓ_2 , in the different RP axes. The flexibility to model correlations separately in the different RP axes is key due to the differences in the extent of the RP domains marginally. The metamodel parameters β_0 , $\boldsymbol{\beta}$, τ^2 , ℓ_1 and ℓ_2 are fit via MLE against the observations \mathbf{y} , \mathbf{X} , and $\boldsymbol{\omega}$ from simulation fits.

Fitting the metamodel allows for a full predictive description of inference under the misspecified restricted models. Predictive estimates are obtained via kriging (Cressie, 2015)

$$\hat{y}(\mathbf{x}) = \beta_0 + \mathbf{x}\boldsymbol{\beta} + \mathbf{r}(\mathbf{x})'\mathbf{R}_\ell^{-1}\left(\mathbf{y} - (\beta_0 + \mathbf{X}\boldsymbol{\beta})\right) \quad (1.15)$$

$\hat{y}(\mathbf{x})$ is the predicted value of the modeled productivity parameter MLE under the two parameter production model, when the index of abundance is generated from the three parameter production model at RP location \mathbf{x} . $\mathbf{r}(\mathbf{x})$ is a vector-valued function of correlation function evaluations for the predictive location \mathbf{x} against all observations in \mathbf{X} (i.e. $\mathbf{r}(\mathbf{x}) = \mathbf{R}(\mathbf{x}, \mathbf{x}_i) \forall \mathbf{x}_i \in \mathbf{X}$).

While metamodeling occurs on the inferred productivity parameters of the restricted production model, the metamodel can also be used to build estimates of major biological RPs. For the BH model the relevant transformations for relating productivity parameters with RPs are given in Eqs. (3.29, 3.32) with γ fixed to -1; for the Schaefer model $\hat{B}^* = \frac{\hat{K}}{2}$ and $\hat{F}^* = \frac{\hat{r}}{2}$. Applying the metamodel predictive surfaces on the scale of RP estimates allows for the quantification of estimation bias that is induced by fitting a misspecified two parameter production model to indices of abundance generated under three parameter productivity.

2.3 Catch

It is known that contrast in the observed index and catch time series can effect inference on the productivity parameters (Hilborn & Walters, 1992). In this setting contrast refers to changes in the long term trends of index data. Figure (3.19, *right*) demonstrates an example of biomass that includes contrast induced by catch. It is not well understood how contrast may factor into inferential failure induced by model misspecification. Thus catch is parameterized so as to allow for a spectrum of possible contrast simulation settings.

217 Catch is parameterized so that $F(t)$ can be controlled with respect to F^* . Recall that
 218 catch is assumed to be proportional to biomass, so that $C(t) = F(t)B(t)$. To control $F(t)$
 219 with respect to F^* , $C(t)$ is specified by defining the quantity $\frac{F(t)}{F^*}$ as the relative fishing rate.
 220 $B(t)$ is defined by the solution of the ODE, and F^* is defined by the biological parameters
 221 of the model. By defining $\frac{F(t)}{F^*}$, catch can then be written as $C(t) = F^* \left(\frac{F(t)}{F^*} \right) B(t)$.

222 Intuitively $\frac{F(t)}{F^*}$ describes the fraction of F^* that $F(t)$ is specified to for the current $B(t)$.
 223 When $\frac{F(t)}{F^*} = 1$, $F(t)$ will be held at F^* , and the solution of the ODE brings $B(t)$ into
 224 equilibrium at B^* . When $\frac{F(t)}{F^*}$ is held constant in time biomass comes to equilibrium as an
 225 exponential decay from K approaching B^* . When $\frac{F(t)}{F^*} < 1$, $F(t)$ is lower than F^* and $B(t)$ is
 226 pushed toward $\bar{B} > B^*$. Contrarily, when $\frac{F(t)}{F^*} > 1$, $F(t)$ is higher than F^* and $B(t)$ is pushed
 227 toward $\bar{B} < B^*$; the precise values of \bar{B} can be calculated from the steady state biomass
 228 equations provided above and depend upon the specific form of the production function.

For the simulations presented here, a family of fishing behaviors are considered where the fishing rate accelerates as technology and fishing techniques improve rapidly until management practices are applied, which ultimately brings fishing into equilibrium at F^* . This is parameterized as three distinct phases, over a total of 45 units of time, with each phase lasting 15 time units. The specific form is given below.

$$\frac{F(t)}{F^*} = ae^{bt}\mathbf{1}_{0 \leq t < 15} + (d - ct)\mathbf{1}_{15 \leq t < 30} + \mathbf{1}_{30 \leq t \leq 45} \quad (1.16)$$

The first term of Eq(3.41) is an exponential increase in fishing, the second term is a linear decline in relative fishing as initial management practices are applied, and the third term, $\mathbf{1}_{30 \leq t \leq 45}$, simply holds the fishing rate at F^* there after. These three phases are controlled by the four parameters a , b , c , and d . By enforcing that the interface of the phases meet at χ_{max} and 1 respectively the relative fishing series is reduced to a two parameter family.

$$a = e^{\log(\chi_{max}) - 15b} \quad b = \frac{1}{t - 15} \log \left(\frac{\chi_{min}}{\chi_{max}} \right) \quad (1.17)$$

$$c = \frac{\chi_{max} - 1}{15 - 1} \quad d = 15c + \chi_{max} \quad (1.18)$$

229 By further specifying $\chi_{max} = 1.6^x$ and $\chi_{min} = 0.4^x$ the two parameters χ_{max} , and χ_{min}

can be reduced to the single parameter χ . The tuning parameter χ then singularly controls contrast that appears in time series data.




Figure 1.3: (left) Relative fishing with low, medium, and high contrast. (right) Population biomass and catch at each associated level of contrast.

When $\chi = 0$, the relative fishing rate is a constant at 1 to create a low contrast simulation environment. As χ increases Eq (3.41) induces more and more contrast in the observed index and catch time series until $\chi = 1$ which produces a high contrast simulation environment. Figure (3.19) demonstrates a spectrum of contrast simulation environments as well as the time series data they induce in the solution of the production model ODE.

2.4 Two Parameter Production Model Inference

The simulated mapping results from fitting an intentionally misspecified two parameter production model to index of abundance data that are generated from a more complex three parameter model of productivity. Thus, let I_t be an index of abundance simulated from the three parameter PT or Schnute production models at time $t \in \{1, 2, 3, \dots, T\}$. However the fitted model is specified to be intentionally misspecified so that the fitted model is driven by a two parameter Schaefer, or BH production model respectively.

The observation model for the fitted model is log-normal such that,

$$I_t|q, \sigma^2, \boldsymbol{\theta} \sim LN(qB_t(\boldsymbol{\theta}), \sigma^2). \quad (1.19)$$

$B_t(\boldsymbol{\theta})$ is defined by the solution of the ODEs defined by the Schaefer, or BH models. For

the Schaefer model $\boldsymbol{\theta} = [r, K]$, and for the BH model $\boldsymbol{\theta} = [\alpha, \beta]$. From the perspective of the fitted model, the observed I_t are assumed independent conditional on q , σ^2 , r , K and the two parameter ODE model for biomass. Thus the log likelihood can be written as

$$\log \mathcal{L}(q, \sigma^2, \boldsymbol{\theta}; I) = -\frac{T}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_t \log \left(\frac{I_t}{qB_t(\boldsymbol{\theta})} \right)^2. \quad (1.20)$$

In this setting, q is fixed at the true value of 0.0005 to focus on the inferential effects of model misspecification on biological parameters. σ^2 and $\boldsymbol{\theta}$ are reparameterized to the log scale and fit via MLE. Reparameterizing the parameters to the log scale improves the reliability of optimization, in addition to facilitating the use of Hessian information for estimating MLE standard errors.

Given that the biological parameters enter the likelihood via a nonlinear ODE, and further the parameters themselves are related to each other nonlinearly, the likelihood function can often be difficult to optimize. A hybrid optimization scheme is used to maximize the log likelihood to ensure that a global MLE solution is found. The R package GA ([Scrucca, 2013, 2017](#)) is used to run a genetic algorithm to explore parameter space globally. Optimization periodically jumps into the L-BFGS-B local optimizer to refine optima within a local mode. The scheme functions by searching globally, with the genetic algorithm, across many initial values for starting the local gradient-based optimizer. The genetic algorithm serves to iteratively improve hot starts for the local gradient-based optimizer. Additionally, optimization is only considered to be converged when the optimum results in an invertible Hessian at the found MLE.

2.5 Continuous model formulation

An important (and often overlooked) implementation detail is the solution to the ODE which defines the progression of biomass through time. As a statistical model it is of paramount importance that this ODE not only have a solution, but also that the solution be unique. Of primary concern, uniqueness of the ODE solution is necessary for well conditioned inference.

If the form of $\frac{dB}{dt}$ is at least Lipschitz continuous, then the Cauchy-Lipschitz-Picard theorem provides local existence and uniqueness of $B(t)$. Recall from Eq(3.14) that $\frac{dB}{dt}$ is

separated into a term for biomass production, $P(B)$, and a term for removals, $Z(t)B(t)$. For determining Lipschitz continuity of $\frac{dB}{dt}$, the smallest Lipschitz constant of $\frac{dB}{dt}$ will be the sum of the constants for each of the terms $P(B)$ and $Z(t)B(t)$ separately. Typically any choice of $P(B)$ will be continuously differentiable, which implies Lipschitz continuity. At a minimum $Z(t)$ typically contains fishing mortality as a function of time $F(t)$ to model catch in time as $C(t) = F(t)B(t)$. $Z(t)$ may or may not contain M , but typically M is modeled as stationary in time and does not pose a continuity issue, unlike some potential assumptions for $C(t)$.

In practice $C(t)$ is determined by a series of observed, assumed known, catches. Catch observations are typically observed on a quarterly basis, but in practice may not be complete for every quarter (or year) of the modeled period. It is overwhelmingly common to discretize the ODE in time via Euler's method with integration step sizes to match the observation frequency of the modeled data. This is often computationally convenient when the underlying species dynamics are reasonably well behaved, however when the dynamics model is used as a statistical model, with the goal of inferring the behavior of the underlying species dynamics, the regularity of the dynamics are not guaranteed. An implicit assumption of continuity of catch in time provides the necessary regularity for the statistical model. Furthermore a continuous handling of the dynamics provides improved accuracy in evaluating the ODE, particularly when inferring productivity parameters which largely control the regularity of the dynamics.

While there are many ways to handle catch continuity, here I assume that catches accrue linearly between observed catches. This assumption defines the catch function as a piecewise linear function of time, with the smallest Lipschitz constant for the catch term defined by the steepest time segment of the catch function. This assumption represents one of the simplest ways of handling catch, while retaining Lipschitz continuity overall. Furthermore linearly interpolated catch is adequately parsimonious for the typical handling of catches.

Integration and Stiffness

As previously mentioned, the overwhelming majority of implementations of stock assessment models discretized the ODE using Euler's method with the integration step sized fixed so as to match the observation frequency. In this setting we explore model parameterizations that

explore the full extent of biologically relevant reference points. This exercise produces some combinations of parameters that result in numerically stiff ODEs.

The concept of stiffness in ODEs is hard to precisely characterize. [Wanner and Hairer \(1996, p.2\)](#) describe stiffness in the following pragmatic sense, “Stiff equations are problems for which explicit methods don’t work”. It is hard to make this definition more mathematically precise, but this is a consistent issue for models of fast growing species in the low contrast simulation. Euler’s method, as often implemented, is particularly poorly suited for these stiff regions of parameter space. In these stiff regions it is necessary to integrate the ODE with an implicit integration method.

Several of the most common implicit methods were tried including the Livermore Solver for ODEs (lsode), and the Variable Coefficient ODE Solver (vode) as implemented in the deSolve package of R ([Soetaert et al., 2010](#)). The difference between implicit solvers is negligible, while explicit methods result in wildly varying solutions to the ODE in stiff regions of parameter space. Results shown here are computed using the lsode integration since it runs relatively quickly and has a relatively smaller footprint in system memory.

3 Results

3.1 PT/Schaefer

An MSY -Optimal Catch History

When $F(t)$ is held constant at F^* , as it is in the "low contrast" simulation setting, $B(t)$ comes to equilibrium as an exponential decay from K to B^* . Understanding model misspecification bias is simplified in this setting due to the relative simplicity that this induces in $B(t)$. However this simplicity is known to poorly inform estimates of r , and thus F^* , due to the limited range of the production function that is observed (Hilborn & Walters, 1992).

Figure (3.20) shows four of the most misspecified example production function fits as compared to the true data generating PT production functions. The rug plots below each set of curves show how the observed biomasses decay exponentially from K to B^* in each case. In particular, notice how observations only exist where the PT biomass is greater than B^* . Due to the leaning of the true PT curves, and the symmetry of the logistic parabola, the logistic curve only observes information about its slope at the origin from data observed on the right portion of the PT curves. The top two panels of Figure (3.20) shows PT data generated such that $\frac{B^*}{B(0)} > 0.5$; in these cases PT is steeper to the right of B^* than it is on the left, and so the the logistic curve over-estimates r , and consequently also over-estimates F^* . The

bottom two panels of Figure (3.20) show PT data generated with $\frac{B^*}{B(0)} < 0.5$ and where the vice versa phenomena occurs. PT is shallower to the right of B^* than it is on the left




Figure 1.4: A comparison of the true PT production function (in black) and the estimated logistic curve (in red) with 95% CI shown. The examples shown represent the four corners of maximum model misspecification in the simulated RP-space. Observed biomasses are plotted in the rug plots below the curves.

and so the logistic parabola estimate tends to under estimate F^* .

Metamodeled Trends

Each point in the space of the RPs F^* and $\frac{B^*}{B(0)}$ uniquely identifies a complete PT model with different combinations of parameters values. Recall that when $\gamma = 2$ for the PT model, the PT curve becomes a parabola and is equivalent to the logistic curve of the Schaefer model. Since the logistic curve is symmetric about B^* , the Schaefer model must fix the value of $\frac{B^*}{B(0)}$ at the constant 0.5 for any value of F^* . So the line through RP space defined by $\frac{B^*}{B(0)} = 0.5 \quad \forall \quad F^*$, defines the subset of RP space where $\gamma = 2$ and where the PT model is equivalent to the Schaefer model. For brevity this subset of RP where $\frac{B^*}{B(0)} = 0.5$ will be referred to as the ‘‘Schaefer set’’. Thus simulated data that are generated along the Schaefer set will be the only data that are not misspecified relative to the Schaefer model; as PT data are simulated farther and farther away from this line at $\frac{B^*}{B(0)} = 0.5$ model misspecification of the Schaefer model becomes worse and worse.

While Figure (3.20) demonstrates a real trend in simulation results, individual simulation runs will at best show jittery trends due to the stochastic nature of statistical inference. The GP process metamodel accounts for this stochasticity to focus analysis on the signal in the simulation results. Recall that metamodeling occurs on the scale of the inferred productivity parameters of the restricted production model, by transforming metamodel predictions via Eq. (3.23), metamodeled predictions are obtained for Schaefer RPs. By further subtracting the true data generating PT RPs from the predicted Schaefer RPs at each point in RP space a pattern of inferential RP bias, induced by model misspecification of the Schaefer model, can be seen.

Figure (3.21) shows the pattern of biases the Schaefer model creates when fit to PT data generated at each point of RP space. An equivalent way to think of Figure (3.21) is that since the Schaefer model must estimate RPs in the Schaefer set, the metamodel arrows indicate the mapping that is created by inferring RPs under a misspecified Schaefer model fit to PT data generated at each point over the pictured region.

Since $\frac{B^*}{B_0}$ must be 0.5 under the Schaefer model, biases in the $\frac{B^*}{B_0}$ direction must simply map vertically onto the Schaefer set. Due to this simplified RP geometry under the Schaefer




Figure 1.5: Joint bias direction for $(F^*, \frac{B^*}{B_0})$ estimates under the misspecified Schaefer Model. The intensity of color represents the excess bias relative to the shortest possible mapping. Results in the low contrast setting are shown *left*, and the high contrast setting is shown *right*.

model, the degree of bias in $\frac{B^*}{B_0}$ estimation is defined solely by the degree of model misspecification irrespective of F^* . Furthermore, the closest possible point along the Schaefer set that Schaefer model inference could map RPs would be the perfectly vertical mapping. This pattern only contains the strictly necessary bias present in $\frac{B^*}{B_0}$, and zero bias in F^* . Any deviation from this minimal bias pattern is necessarily due to added bias in F^* .

The two simulation settings shown in Figure (3.21) are identical except for the amount of contrast present in the simulated index. The left panel of Figure (3.21) shows RP biases in the low contrast setting, while the right panel shows the high contrast setting. Notice that in the low contrast setting the RP bias pattern is far from the minimum distance mapping, however when contrast is added the mapping becomes much closer to a minimal bias mapping. In the low contrast setting the observed bias is consistent with the pattern and mechanism described in Figure (3.20), where F^* is underestimated for data generated below the Schaefer line and overestimated above the Schaefer set. In the high contrast simulation the mapping is nearly minimal distance with the exception of PT data generated with simultaneously low $\frac{B^*}{B_0}$ and high F^* .

Figure (6 .1) demonstrates how bias in F^* estimation decreases as contrast is added to




Figure 1.6: Bias in F^* under the Schaefer model when PT data are generated with increasing contrast so that F^* and $\frac{B^*}{B_0}$ are fixed at 0.699 and 0.201 respectively.

PT data as generated in the low $\frac{B^*}{B_0}$ and high F^* regime. By including additional contrast F^* bias is decreased, however parameterizing contrast so as to fully extinguish F^* bias may require a more complex model of fishing.

4 Discussion

Results presented here generally agree with what is known about estimating growth rate parameters (Lee et al., 2012; Conn et al., 2010; Magnusson & Hilborn, 2007). These study's appreciate the role of contrast for estimating growth rates, however they struggle to make generally extensible conclusions since they focus only on a handful of stocks that fall short of forming a random sample of the greater population of possible stock behaviors. The LHS design methods presented here are designed specifically to simulate a representative sample of stocks broadly across the space of possible RPs. Furthermore, the simulation design, taken together with the GP metamodel of productivity parameter estimates, allows this study to control the degree of model misspecification and generalize conclusions about the behavior of productivity estimation within the production model setting presented.

In the presence of contrast, F^* estimation can enjoy very low bias even for a wide range of poorly specified models; conversely in the absence of contrast F^* estimation can suffer very large bias even for slightly misspecified models. This pattern is particularly true for inference

under the Schaefer model where the geometry of the restricted RP set isolates estimation failure of F^* from $\frac{B^*}{B(0)}$. While contrast has a similar impact on F^* estimation under the BH model, the geometry of the BH RP set correlates estimation bias of F^* and $\frac{B^*}{B(0)}$. The GP metamodeling approach reveals a more general pattern that highly informative data sets (high contrast) produces a nearly minimal distance mapping of RPs onto the constrained RP set.

In all cases when model misspecification is removed, even with weakly informative data, RP estimation is unbiased and well estimated. Thus contrast alone is not the only factor leading to inferential failure. Model misspecification is a necessary but not sufficient condition for inducing RP estimation bias. The particular RP bias present depends on the RP geometry of the fitted model and how that geometry is misspecified relative to the data. The RP mapping is then oriented to the RP geometry of the fitted model.

While the relative fishing rate parameterized in Section (5.5) captures a usefully broad spectrum of relevant fishing behaviors, it is still limiting in the amount of information that it can induce. Improved methods for quantifying contrast in fisheries data, and/or methods of discovering more informative fishing behavior, could improve this analysis. In the absence of a maximally informative dataset simulation methods will not fully describe how inference fails, but the methods presented here tell the most complete picture yet, with explicit control of the degree model misspecification, contrast, and a simulation design that allows for uniform representative data generation across biologically meaningful stocks. The results presented here suggest the conjecture that under a maximally informative dataset, RP inference with a two parameter production function will be biased in the direction a shortest distance map from the true RPs onto restricted set of RPs under the two parameter model.

Given the potential for model misspecification of RPs, a minimal distance mapping of RPs represents a best-case scenario where the total bias of RPs, when measured jointly, is minimized. That said, without recognizing the geometry of how two parameter models of productivity limit RP space this may lead to unintuitive implications in RP estimation. For example, due to the shape of the BH RP set a minimal distance mapping ensures that if there is bias in one of $\frac{B^*}{B_0}$ or F^* , there will necessarily be bias in the other RP. However under the Schaefer model, since the RP set is a constant in $\frac{B^*}{B_0}$, bias in F^* is not adulterated in the

same way by bias in $\frac{B^*}{B_0}$ estimation. While models with constant RPs, such as the logistic model $\frac{B^*}{B_0} = \frac{1}{2}$ or the Fox model $\frac{B^*}{B_0} = \frac{1}{e}$, are extremely limited, they can be valuable tools for developing intuition precisely because they isolate RP estimation in their free RPs from the correlated RP biases present in models like the BH or Ricker model.

When one considers the implications of RP bias, overestimation of RPs carries the severe implication of management recommendations potentially leading to overfishing, while underestimation of RP leads to overly conservative management. In this sense, when the true model is not known, the geometry of the BH set together with the metamodeled bias trends makes the BH model a naturally conservative estimator of RPs for most stocks. For most non-BH populations the BH model is likely to make conservative errors in its estimates of F^* and $\frac{B^*}{B_0}$. The one notable exception to the conservatism of the BH model stands for data generated in the Cushing-like regime of Schnute RPs. In this regime the BH model tends to be fairly unbiased overall, however the bias that is present for these populations tends to be overestimation in both RPs, leading to much more severe management consequences for those populations.

The RP bias trends of the Schaefer model demonstrate much less conservatism than the BH overall. For any population with $\frac{B^*}{B_0} < 0.5$, $\frac{B^*}{B_0}$ will be overestimated. When the population comes from the regime where $\frac{B^*}{B_0} > 0.5$, $\frac{B^*}{B_0}$ will be under estimated, but F^* is likely to be overestimated depending on the degree of contrast present in the data. So while the Schaefer model is an intuitive model, it tends to lead to much less conservative RP estimation.

While it is important to recognize these limitations of two parameter models of productivity, we should not solely accept conservatism as a rationale of choosing a BH model of productivity. Increasing the flexibility of the production function by moving toward three parameter models would release the underlying structural limitations (Mangel et al., 2013) that cause these RP biases in the first place. Punt and Cope (2019) considers a suite of possible three parameter curves which could be used instead of current two parameter curves. For all of their benefits, three parameter production functions have their own complicating factors, and the structure present in the Schnute model explored here makes it an intuitive bridge model for developing three parameter models going forward.

- show a schnute fit to data? (Yeakel & Mangel, 2015) Prior

- summary of σ over RP space comparing between models (PT, Schnute, Schnute DD) to show areas of model breakdown.

 - miss-identifying signal for noise.
 - It happens more as the dynamics get more complex.
 - point to the full age structured models.
- show the constrained BH space over a grid of $M, \kappa, \omega, W_\infty$
- Show that the constrained spaces vary only slightly as compared with the consequences of misspecifying the functional form.
- estimating these other quantities (while they can create quite different Biomass series) can only do so much to improve (expand) RP inference as compared with correctly modeling P .
- mapping distance as a function of contrast at (3.5, 0.5)
- for LHS grid locations show $\frac{B^*}{B_0}$ and F^* biases for grids in $M \in (0, 0.5)$ For sure in High Contrast, maybe also in Low??.

5 Appendix: Inverting $\frac{B^*}{B(0)}$ and γ for the PT Model

For brevity let $\zeta = \frac{B^*}{B(0)}$.

$$\begin{aligned}\zeta &= \left(\frac{1}{\gamma}\right)^{\frac{1}{\gamma-1}} \\ \zeta &= \gamma \zeta^\gamma \\ \zeta &= \gamma e^{\gamma \log(\zeta)} \\ \zeta \log(\zeta) &= \gamma \log(\zeta) e^{\gamma \log(\zeta)}\end{aligned}$$

The Lambert product logarithm, W , is defined as the inverse function of $z = xe^x$ such that $x = W(z)$. Applying this definition allows for the isolation of γ .

$$\begin{aligned}\gamma \log(\zeta) &= W(\zeta \log(\zeta)) \\ \gamma &= \frac{W(\zeta \log(\zeta))}{\log(\zeta)}\end{aligned}\tag{1.21}$$

The Lambert product logarithm is a multivalued function with a branch point at $-\frac{1}{e}$. The principal branch, $W_0(z)$, is defined on $z \in (-\frac{1}{e}, \infty)$, and the lower branch, $W_{-1}(z)$, is defined on $z \in (-\frac{1}{e}, 0)$. Taken individually, each respective branch is analytic, but cannot be expressed in terms of elementary functions.

When $\zeta \in (0, \frac{1}{e})$ the solution of interest in Eq. (3.24) comes from W_0 . When $\zeta \rightarrow \frac{1}{e}$, the Fox Model emerges as $\gamma \rightarrow 1$. When $\zeta \in (\frac{1}{e}, 1)$ the solution of interest comes from W_{-1} . For the use case presented here, Eq. (3.24) is to be interpreted as,

$$\gamma = \begin{cases} \frac{W_0(\zeta \log(\zeta))}{\log(\zeta)} & \zeta \in (0, \frac{1}{e}) \\ \frac{W_{-1}(\zeta \log(\zeta))}{\log(\zeta)} & \zeta \in (\frac{1}{e}, 1) \end{cases}.\tag{1.22}$$

Prager 2002, Figure(2).

<https://math.stackexchange.com/questions/3004835/is-the-lambert-w-function-analytic-if-not-everywhere-then-on-what-set-is-it-analytic> <https://researchportal.bath.ac.uk/en/publications/algebraic-properties-of-the-lambert-w-function-from-a-result-of-r>

490 Chapter 2

491 The Schnute Model

0.1 Schnute Model

The Schnute production function is a three parameter generalization of many of the most common two parameter production functions (Deriso, 1980; J. Schnute, 1985). It can be written in the following form, with parameters α , β , and γ ,

$$P_s(B; [\alpha, \beta, \gamma]) = \alpha B(1 - \beta\gamma B)^{\frac{1}{\gamma}}. \quad (2.1)$$

The BH and Logistic production functions arise when γ is fixed to -1 or 1 respectively. The Ricker model is a limiting case as $\gamma \rightarrow 0$. For $\gamma < -1$ a family of strictly increasing Cushing-like curves arise, culminating in linear production as $\gamma \rightarrow -\infty$. These special cases form natural regimes of similarly behaving production functions as seen in Figure (3.15).

The behavior of RP inference under the BH model is of particular interest due to the overwhelming popularity of the BH assumption in fisheries models. Since Schnute production models can represent a quantifiably wide variety of possible productivity behaviors, they present an ideal simulation environment for inquiry of the reliability of inference under the BH assumption.

Under Schnute production, biomass dynamics evolve according to the following ODE,

$$\frac{dB}{dt} = P_s(B; \theta) - (M + F)B. \quad (2.2)$$

This equation largely takes the same form as previously described, except that P_s is the Schnute production function and natural mortality, M , is modeled explicitly here. Natural




Figure 2.1: The Schnute production function plotted across a variety of parameter values. Regimes of similarly behaving curves are grouped by color.

512 mortality models the instantaneous rate of mortality from all causes outside of fishing. Ex-
 513 plicitly modeling natural mortality is not only a typical assumption of fisheries models, but
 514 is also key to the making RPs well defined over the relevant domain of γ .

The derivation of RPs under Eq. (3.26) follows a similar logic as under the PT model. An expression for equilibrium biomass is attained by setting $\frac{dB}{dt} = 0$ and rearranging the resulting expression to solve for B

$$\bar{B}(F) = \frac{1}{\gamma\beta} \left(1 - \left(\frac{M+F}{\alpha} \right)^\gamma \right). \quad (2.3)$$

The above expression quickly yields B_0 , B^* by evaluation at $F = 0$ and F^* respectively,

$$B_0 = \frac{1}{\gamma\beta} \left(1 - \left(\frac{M}{\alpha} \right)^\gamma \right) \quad (2.4)$$

$$\frac{B^*}{B_0} = \frac{1 - \left(\frac{M+F^*}{\alpha} \right)^\gamma}{1 - \left(\frac{M}{\alpha} \right)^\gamma}. \quad (2.5)$$

Attaining an expression for F^* requires maximization of equilibrium yield, $\bar{Y} = F\bar{B}(F)$, with respect to F . Analytically maximizing proceeds by differentiating \bar{Y} to produce

$$\frac{d\bar{Y}}{dF} = \bar{B}(F) + F \frac{d\bar{B}}{dF} \quad (2.6)$$

$$\frac{d\bar{B}}{dF} = -\frac{1}{\beta} \left(\frac{\left(\frac{M+F}{\alpha} \right)^\gamma}{F+M} \right). \quad (2.7)$$

Setting $\frac{d\bar{Y}}{dF} = 0$, filling in the expressions for $\bar{B}(F)$ and $\frac{d\bar{B}}{dF}$, then rearranging to solve for F^* is less yielding here than it was in the case of the PT model. This procedure falls short of providing an analytical solution for F^* directly in terms of θ , but rather shows that F^* must respect the following expression,

$$0 = \frac{1}{\gamma} - \left(\frac{1}{\gamma} + \frac{F^*}{F^* + M} \right) \left(\frac{F^* + M}{\alpha} \right)^\gamma. \quad (2.8)$$

515 The lack of an analytical solution here is understood. [J. T. Schnute and Richards \(1998,](#)
 516 [pg. 519\)](#) specifically points out that F^* cannot be expressed analytically in terms of produc-
 517 tivity parameters, but rather gives a partial analytical expression for the inverse relationship.

Although parameterized slightly differently, [J. T. Schnute and Richards \(1998\)](#) derives expressions for α and β as a function of RPs and γ .

Since RPs are left without a closed form expression, computing RPs from productivity parameters amounts to numerically solving the system formed by collecting the expressions (3.32), (3.28), and (3.29).

Simulation

For the purposed of simulation, it is not necessary to completely know the precise relationships mapping RPs $\mapsto \theta$ or $\theta \mapsto$ RPs. Simulation only requires enough knowledge of these mappings to gather a list of (α, β, γ) tuples, for data generation under the Schnute model, and the corresponding RPs in some reasonable space-filling design over RP space.

Similarly to [J. T. Schnute and Richards \(1998\)](#), expressions (3.32) and (3.28) are solved for α and β respectively. This leads to the partial mapping $(F^*, B_0) \mapsto (\alpha(\cdot, \gamma), \beta(\cdot, \gamma))$ in terms of RPs and γ . By further working with Eq. (3.29), to identify γ , the following system is obtained,

$$\begin{aligned}\alpha &= (M + F^*) \left(1 + \frac{\gamma F^*}{M + F^*} \right)^{1/\gamma} \\ \beta &= \frac{1}{\gamma B_0} \left(1 - \left(\frac{M}{\alpha} \right)^\gamma \right) \\ \frac{B^*}{B_0} &= \frac{1 - \left(\frac{M + F^*}{\alpha} \right)^\gamma}{1 - \left(\frac{M}{\alpha} \right)^\gamma}.\end{aligned}\tag{2.9}$$

For a population experiencing natural mortality M , by fixing F^* , B_0 , and $\frac{B^*}{B_0}$ the above system can fully specify α and β for a given γ . Notice for a given γ a cascade of closed form solutions for α and β can be obtained. First $\alpha(\gamma)$ can be computed, and then $\beta(\alpha(\gamma), \gamma)$ can be computed. If $\alpha(\gamma)$ is filled back into the expression for $\frac{B^*}{B_0}$, the system collapses into a single onerous expression for $\frac{B^*}{B_0}(\alpha(\gamma), \gamma)$. For brevity, define the function $\zeta(\gamma) = \frac{B^*}{B_0}(\alpha(\gamma), \gamma, F^*, M)$ based on Eq. (3.29).

Inverting $\zeta(\gamma)$ for γ , and computing the cascade of $\alpha(\gamma)$, and then $\beta(\alpha(\gamma), \gamma)$, fully defines the Schnute model for a given $(\frac{F^*}{M}, \frac{B^*}{B_0})$. However inverting ζ accurately is extremely difficult. Inverting ζ analytically is not feasible, and numerical methods for inverting ζ are unstable

and can be computationally expensive. Rather than numerically invert precise values of $\zeta(\gamma)$, γ is sampled so that the overall simulation design is space filling as described in Section (5.3).


Each design location defines a complete Schnute production model with the given RP values. Indices of abundance are simulated from the Schnute model at each design location, a small amount of residual variation, $\sigma = 0.01$, is added to the simulated index, and the data are then fit with a misspecified BH production model. The design at large captures various degrees of model misspecification relative to the BH model, so as to observe the effect of productivity model misspecification upon RP inference.

0.2 Latin Hypercube Sampling

The goal of space filling design in this setting is to extend the notion of the random sample (and its desirable parameter estimation properties) across the simulated RP domain so as to represent the simulated space as well as possible (Gramacy, 2020). The simple random sample is the classical approach to unbiased parameter estimation, however simple randomness is patchy, often sampling some regions of design space quite densely, while leaving other regions of design space empty. Space filling designs aim to preserve (or enhance) parameter estimation properties across the simulated domain (Devon Lin & Tang, 2015; Stein, 1987), while constraining samples to be spaced in some notion of spread over the entire space. Latin hypercube sampling (McKay et al., 2000, LHS) is among the most foundational of space filling designs used in computer experiments.

A LHS of size n , in the 2 dimensional space defined by RPs, distributes samples so as to spread points across a design region in a broadly representative way. A LHS design extends the notion of a univariate random uniform sample across multiple dimensions so that each margin of the design space enjoys a uniform distribution.

LHS designs achieve this notion of uni-



formity by first partitioning each dimension of the design space into regular grids of size n . By intersecting the grids of each dimension, cells are produced that evenly partition the design space. In two dimensions n^2 cells are produced, from which a total of n samples are taken. Crucially only one sample is taken from a given element of each grid in each dimension so as to reduce clumping of the n samples across the design space.

Schnute Design

Due to the lack of an analytical relationship mapping RPs $\mapsto \theta$, analogous to the PT model's Eq. (3.24), producing a LHS design over Schnute RPs requires a more tactful approach. The structured relationship between the RPs and productivity parameters, described in Section (5), allows an approximate LHS to be obtained by a careful navigation of the system of equations seen in Eq. (3.33).

Under the Schnute model, let \mathcal{F} and \mathcal{B} represent regular grids on $\frac{F^*}{M} \in (0.25, 4)$ and $\frac{B^*}{B_0} \in (0.15, 0.7)$ respectively which can serve as the scaffolding for computing an approximate LHS.

Since it is not practical to invert $\zeta(\gamma)$, a uniform sample in $\frac{B^*}{B_0}$ can be obtained by modeling γ as a random variable, with realization γ^* , and thinking of $\zeta(\gamma)$ as its cumulative distribution function (CDF). The aim is to model γ as an easily sampled random variable with a CDF that closely approximates ζ , so that $\zeta(\gamma^*) \sim U(\zeta_{min}, 1)$ as closely as possible. There may be many good models for the distribution of γ , but in this setting the

Given B_0 , M , and F^* :

- 1) Draw $\gamma^* \sim \gamma|F^*, M$.
- 2) Compute $\frac{B^*}{B_0} = \zeta(\gamma^*)$
- 3) Compute $\alpha^* = \alpha(\gamma^*, F^*, M)$
- 4) Compute $\beta^* = \beta(\alpha^*, \gamma^*, M, B_0)$

Figure 2.3: An outline of the sampling procedure for γ given B_0 , M , and F^* .

following distribution is very effective,

$$\gamma \sim \zeta_{min} \delta(\gamma_{min}) + t(\mu, \sigma, \nu) \mathbf{1}_{\gamma > \gamma_{min}}. \quad (2.10)$$

586 Above, t is the density of the three pa-
 587 rameter location-scale family Student's t dis-
 588 tribution with location μ , scale σ , and de-
 589 grees of freedom ν . $\mathbf{1}_{\gamma > \gamma_{min}}$ is an indica-
 590 tor function that serves to truncate the Stu-
 591 dent's t distribution at the lower bound γ_{min} .
 592 $\delta(\gamma_{min})$ is the Dirac delta function evaluated
 593 at γ_{min} , which is scaled by the known value
 594 ζ_{min} ; this places probability mass ζ_{min} at
 595 the point γ_{min} . Since sampling from a Stu-
 596 dent's t distribution is readily doable, sam-
 597 pling from a truncated Student's t mixture
 598 only requires slight modification.

Let T be the CDF of the modeled distri-
 bution of γ . Since the point $(\gamma_{min}, \zeta_{min})$ is
 known from the dynamics of the Schnute model at a given RP, full specification of Eq. (3.34)
 only requires determining the values for μ , σ , and ν which make T best approximate $\zeta(\gamma)$.
 Thus, the values of μ , σ , and ν are chosen by minimizing the L^2 distance between $T(\gamma)$ and
 $\zeta(\gamma)$.




Figure 2.4: $\zeta(\gamma)$ Plotted for $F^* = 0.1$ and $M = 0.2$. The point $(\gamma_{min}, \zeta_{min})$ shows the lowest biologically meaningful value of γ ; below which productivity is negative.

$$[\hat{\mu}, \hat{\sigma}, \hat{\nu}] = \arg \min_{[\mu, \sigma, \nu]} \int_{\Gamma} (T(\gamma; \mu, \sigma, \nu) - \zeta(\gamma))^2 d\gamma \quad (2.11)$$

The distribution $T(\gamma|\hat{\mu}, \hat{\sigma}, \hat{\nu})$ is fit for use in generating γ^* random variates at a specific F^* and M . This approximation releases the need to invert ζ w.r.t γ by using samples of γ^* values to generate approximately uniform samples of $\zeta(\gamma^*)$. By sampling approximately uniform $\zeta(\gamma^*)$ random variates in this way, and making use of the structure in Eq. (3.33), an approximate LHS sample can be collected via Algorithm (2).

$\frac{F^*}{M}$ is drawn uniformly from \mathcal{F}_i . Conditioning on the sample of F^* , and M , $T(\gamma|\hat{\mu}, \hat{\sigma}, \hat{\nu})$ is fit and γ^* is sampled. ζ^* is then computed and placed into the appropriate grid element \mathcal{B}_j . Given γ^* , the cascade $\alpha(\gamma^*)$, and $\beta(\alpha(\gamma^*), \gamma^*)$, can be computed.

The algorithm continues until all of the de-

sign elements, $(\frac{F^*}{M}, \zeta^*) \Leftrightarrow (\alpha^*, \beta^*, \gamma^*)$, have been computed for all $i \in [1, \dots, n]$.

Design Refinement

Since the behavior of RP inference, under misspecified models, will vary in yet-unknown ways, the exact sampling design density may be hard to know a priori. Several factors, including the particular level of observation uncertainty, high variance (i.e. hard to resolve) features of the response surface, or simply "gappy" instantiations of the initial LHS design may necessitate adaptive design refinement, to accurately describe RP biases. Given the temperamental relationship between RPs and productivity parameters in the Schnute model, a recursive refinement algorithm that makes use of the previously described LHS routine, is developed.

While LHS ensures uniformity in the design margins, and a certain degree of spread, it

Algorithm 1 LHS of size n on rectangle R .

```

1: procedure  $LHS_n(R)$ 
2:   Define  $n$ -grids  $\mathcal{F}, \mathcal{B} \in R$ 
3:   for each grid element  $i$  do
4:     Draw  $\frac{F^*}{M} \sim Unif(\mathcal{F}_i)$ 
5:     Compute  $[\hat{\mu}, \hat{\sigma}, \hat{\nu}]$  given  $F^*$  &  $M$ 
6:     while  $\mathcal{B}_j$  not sampled do
7:       Draw  $\gamma^* \sim T(\gamma|\hat{\mu}, \hat{\sigma}, \hat{\nu})$ 
8:       Compute  $\zeta^* = \zeta(\gamma^*)$ 
9:       Compute  $j$  such that  $\zeta^* \in \mathcal{B}_j$ 
10:    end while
11:    Compute  $\alpha^* = \alpha(\gamma^*, F^*, M)$ 
12:    Compute  $\beta^* = \beta(\alpha^*, \gamma^*, M, B_0)$ 
13:    Save  $(\frac{F^*}{M}, \zeta^*) \Leftrightarrow (\alpha^*, \beta^*, \gamma^*)$  in  $\mathcal{F}_i \times \mathcal{B}_j$ 
14:  end for
15: end procedure

```

is widely recognized that particular LHS instantiations may leave substantive gaps in the simulation design. To correct this, LHS is often paired with design elements of maximin design (Morris & Mitchell, 1995; Devon Lin & Tang, 2015). Maximin designs sample the design space by maximizing the minimum distance between sampled points. This has the advantage of definitionally filling holes in the design, however because no points are ever drawn outside of the design domain, samples tend to clump around edges (particularly corners) of the design domain. Since LHS ensures uniformity in the margins and maximin designs enjoys a certain sense of optimality in how they define and fill gaps (Johnson et al., 1990), the methods are quite complimentary when combined.

Making use of this complimentary relationship, holes in the existing LHS design of RPs are identified based on maximin design principles. New design points are collected based on areas of the RP design space which maximizes the minimum distance between all pairs of points in the current design, based on the following distance function

$$d(\mathbf{x}, \mathbf{x}') = \sqrt{(\mathbf{x} - \mathbf{x}')^T \mathbf{D}^{-1} (\mathbf{x} - \mathbf{x}')} \quad (2.12)$$

$$\mathbf{D} = \mathbf{diag} \left[\left(\max(\mathcal{F}) - \min(\mathcal{F}) \right)^2, \left(\max(\mathcal{B}) - \min(\mathcal{B}) \right)^2 \right].$$

Above, d is a scaled distance function that defines the distance between points in the differing scales of $\frac{B^*}{B_0}$ and $\frac{F^*}{M}$. \mathbf{D} is a diagonal matrix that measures the squared size of the domain in each axis of so as to normalize distances to a common scale.

If \mathbf{X}_n is the initial design, computed on R_{full} , let \mathbf{x}_a be the augmenting point which maximizes the minimum distance between all of the existing design points,

$$\mathbf{x}_a = \underset{\mathbf{x}'}{\operatorname{argmax}} \min \{ d(\mathbf{x}_i, \mathbf{x}') : i = 1, \dots, n \}. \quad (2.13)$$

The point \mathbf{x}_a is used as an anchor for augmenting \mathbf{X}_n . An additional $LHS_{n'}$ (via Algorithm (2)) is collected, adding n' design points, centered around \mathbf{x}_a , to the overall design. The augmenting region, $R_{(\mathbf{x}_a, d_a)}$, for collecting $LHS_{n'}$ is defined based on the square centered at \mathbf{x}_a with side length $2d_a$, where $d_a = \min \{ d(\mathbf{x}_i, \mathbf{x}_a) : i = 1, \dots, n \}$, in the space defined by the metric d .

Due to the tendency of maximin sampling to cluster augmenting points on the edges of the design space, $R_{(x_a, d_a)}$ is truncated by the outer most limits of R_{full} so as to focus design augmentation within the specified domain of the simulation. Furthermore, since the design space has a nonlinear constraint at low values of $\frac{B}{B_0}$, the calculation of x_a is further truncated based on a convex hull defined by the existing samples in the overall design.

Design refinement then proceeds as follows. An initial design is computed, $X_n = LHS_n(R_{full})$, based on an overall simulated region of RPs R_{full} . The maximin augmenting point, x_a , is computed at a maximin distance of d_a from the existing samples. An augmenting design $X_{n'} = LHS_{n'}(R_{(x_a, d_a)})$ is collected and added to X_n . Design refinement carries on recursively collecting augmenting designs in this way until the maximin distance falls below the desired level.

0.3 Gaussian Process Metamodel

At its core, a metamodel is simply a model of some mapping of inputs to outputs (the mapping itself is typically defined by a computer model). By modeling the mapping with a statistical model (that explicitly defines the relevant features of the mapping) a metamodel defines a specific ontology for the mapping. By simulating examples of the mapping, the inferential infrastructure of the statistical model is used to empirically learn an effective emulation of the mapping within the ontology defined by the statistical model. The predictive infrastructure of the statistical model is then useful as an approximate abstraction of the system itself to better understand the system through further data collection, cheap approximation of the mapping, and/or study of the mapping itself.

In this setting, the aim of metamodeling is to study how well RPs are inferred when typical two parameter models of productivity (Logistic and BH) are misspecified for populations that are actually driven by more complicated dynamics. The simulation design, \mathbf{X} , provides a sample of different population dynamics that are driven by three parameter production functions broadly in RP space. By simulating index of abundance data from the three parameter model, and fitting those data with the two parameter production model, we observe particular instances of how well RPs are inferred at the given misspecification of the two parameter model relative to the true three parameter production model. By gathering

all of the simulated instances of how RPs are inferred (under the two parameter model), we form a set of example mappings to train a metamodel which represents the mapping of true RPs (under the three parameter model) to estimates of RPs under the misspecified two parameter production model. The metamodel is essentially a surrogate for inference under the misspecified two parameter production model that controls for the specific degree of model misspecification.

A flexible GP model is assumed for the structure of the metamodel to describe the mapping of RPs under misspecified two parameter models of productivity. A GP is a stochastic process generalizing the multivariate normal distribution to an infinite dimensional analog. GP models are often specified primarily through the choice of a covariance (or correlation) function which defines the relationship between locations in the input space. Typically correlation functions are specified so that points closely related in space result in correlated effects in the model. In this setting the inputs to the GP metamodel are the space of reference points which define the simulated three parameter production models.

While index of abundance data are generated from three parameter models, at each design location of the simulation, fitting the restricted two parameter model results in a maximum likelihood estimate (MLE; and associated estimation uncertainty) of each of the productivity parameters (i.e. Schaefer: $[\log(r), \log(K)]$, BH: $[\log(\alpha), \log(\beta)]$). To simplify the specification of the metamodel, let \mathbf{y} be a vector collecting the fitted MLEs for one of the productivity parameters, and let $\boldsymbol{\omega}$ be a vector of estimates of the estimator variances (via the inverted Fisher information) at each \mathbf{y} . Each of the fitted productivity parameter estimates are then modeled using independent instances of the following GP metamodel.

$$\begin{aligned}\mathbf{y} &= \beta_0 + \mathbf{X}\boldsymbol{\beta} + \mathbf{v} + \boldsymbol{\epsilon} \\ \mathbf{v} &\sim N_n(\mathbf{0}, \tau^2 \mathbf{R}_\ell) \\ \boldsymbol{\epsilon} &\sim N_n(\mathbf{0}, \boldsymbol{\omega}' \mathbf{I})\end{aligned}\tag{2.14}$$

\mathbf{X} is the $n \times 2$ LHS design matrix of RPs for each simulated three parameter data generating model as described in Section (5 .3). ϵ models independent normally distributed error, which provides an ideal mechanism for propagating uncertainty from inference in the

simulation step into the metamodel. By matching each y_i with an observed ω_i variance term, ϵ serves to down weight the influence of each y_i in proportion to the inferred production model sampling distribution uncertainty. This has the effect of smoothing the GP model in a way similar to the nugget effect (Gramacy & Lee, 2012), although the application here models this effect heterogeneously.

The term, \mathbf{v} , contains spatially correlated GP effects. The correlation matrix, \mathbf{R}_ℓ describes how RPs close together in the simulation design are more correlated than those that are far away. This spatial effect is modeled with a squared exponential correlation function,

$$R(\mathbf{x}, \tilde{\mathbf{x}}) = \exp \left(\sum_{i=1}^2 \frac{-(x_i - \tilde{x}_i)^2}{2\ell_j^2} \right). \quad (2.15)$$

R has an anisotropic separable form which allows for differing length scales, ℓ_1 and ℓ_2 , in the different RP axes. The flexibility to model correlations separately in the different RP axes is key due to the differences in the extent of the RP domains marginally. The metamodel parameters β_0 , $\boldsymbol{\beta}$, τ^2 , ℓ_1 and ℓ_2 are fit via MLE against the observations \mathbf{y} , \mathbf{X} , and $\boldsymbol{\omega}$ from simulation fits.

Fitting the metamodel allows for a full predictive description of inference under the misspecified restricted models. Predictive estimates are obtained via kriging (Cressie, 2015)

$$\hat{y}(\mathbf{x}) = \beta_0 + \mathbf{x}\boldsymbol{\beta} + \mathbf{r}(\mathbf{x})'\mathbf{R}_\ell^{-1}(\mathbf{y} - (\beta_0 + \mathbf{X}\boldsymbol{\beta})) \quad (2.16)$$

$\hat{y}(\mathbf{x})$ is the predicted value of the modeled productivity parameter MLE under the two parameter production model, when the index of abundance is generated from the three parameter production model at RP location \mathbf{x} . $\mathbf{r}(\mathbf{x})$ is a vector-valued function of correlation function evaluations for the predictive location \mathbf{x} against all observations in \mathbf{X} (i.e. $\mathbf{r}(\mathbf{x}) = \mathbf{R}(\mathbf{x}, \mathbf{x}_i) \forall \mathbf{x}_i \in \mathbf{X}$).

While metamodeling occurs on the inferred productivity parameters of the restricted production model, the metamodel can also be used to build estimates of major biological RPs. For the BH model the relevant transformations for relating productivity parameters with RPs are given in Eqs. (3.29, 3.32) with γ fixed to -1; for the Schaefer model $\hat{B}^* = \frac{\hat{K}}{2}$ and

711 $\hat{F}^* = \frac{\hat{r}}{2}$. Applying the metamodel predictive surfaces on the scale of RP estimates allows for
712 the quantification of estimation bias that is induced by fitting a misspecified two parameter
713 production model to indices of abundance generated under three parameter productivity.

Results

1.1 Schnute/BH

Design

Algorithm (1) enforces uniform marginals in $\frac{F^*}{M}$ directly, as well as the adherence of the overall design to latin squares. Figure (2.5) shows a uniform Q-Q plot for sampled ζ , using Algorithm (1), against theoretical uniform quantiles. As evidence by the excellent coherence to the theoretical uniform quantiles, the approximation in Section (5.3) for sampling γ (and therefore $\zeta(\gamma)$), is very effective. Furthermore since numerical inversion of $\zeta(\gamma)$ is costly and unreliable, the relative speed and accuracy that this approximate LHS sampling method provides is pivotal for the rest of the work presented here.

Similarly to the PT model, the three parameter Schnute model is uniquely identified by each point in the space of $\frac{F^*}{M}$ and $\frac{B^*}{B_0}$ RPs. As seen in Figure (2.6), Schnute production has different behaviors in different ranges of RPs space, which are entirely defined by the value of γ (shown in Figure (3.15)). When $\gamma \geq 1$ the Schnute model produces a family of Logistic-like curves that are increasingly right leaning as γ increases. For $1 > \gamma \geq 0$, Schnute production takes a family of left leaning Ricker-like curves that all, at least, approach the x-axis. For $0 > \gamma > -1$ there are a family of BH-like curves that do not




Figure 2.5: Uniform Q-Q plot for ζ plotted for $F^* = 0.1$ and $M = 0.2$.




Figure 2.6: A Schnute RP design. Colors indicate different regimes of Schnute production. The black curve shows the BH set.

approach the x-axis but still have decreas-

ing productivity for large biomass stocks. When γ is exactly -1 Schnute reduces to BH production which has asymptoting production for large biomass. Finally when $-1 > \gamma$ Schnute produces a family of increasing Cushing-like curves that do not asymptote, and produces linear production as $\gamma \rightarrow -\infty$.

Modeling index data that are simulated broadly over the theoretical space of RPs with misspecified BH production greatly limits the range of possible RPs that can be inferred. Under BH production the full theoretical space of RPs are limited to the curve $\frac{B^*}{B_0} = \frac{1}{F^*/M+2}$. Define the “BH set” as the set of RPs defined by this limited space, i.e. the curve $\left\{ \left(\frac{F^*}{M}, \frac{B^*}{B_0} \right) \mid \frac{B^*}{B_0} = \frac{1}{F^*/M+2} \right\}$. as seen in the black curve in Figure (2.6). The farther away from this set that Schnute data are simulated, the worse the BH model is misspecified for those data.

Metamodeled Trends

Unlike the Schaefer model, the BH set is not a constant in $\frac{B^*}{B_0}$. Under the BH model, bias in $\frac{B^*}{B_0}$ is no longer entirely defined by the degree of model misspecification, but rather the structure of BH RPs allows bias in both $\frac{B^*}{B_0}$ and $\frac{F^*}{M}$ to interact as a function of contrast in the data.

High Contrast Figure (2.7) shows metamodeled RP bias surfaces for inference under the BH model in the high contrast setting. The (*left*) and (*bottom*) panels focus only on the $\frac{B^*}{B(0)}$ and $\frac{F^*}{M}$ components of bias respectively. In these panels bias is shown as relative bias, $\frac{\widehat{RP}-RP}{RP}$, similar to a percent error calculation. Where RP represents the true value of the three parameter RP, and \widehat{RP} refers to the metamodel estimate.

Figure (2.7, *top-right*) combines the components of bias to show the overall mapping of RPs under BH inference in the high contrast simulation setting. Unlike high contrast RP inference under the Schaefer model, the BH model does shows bias in both RPs here. Despite the bias in $\frac{B^*}{B(0)}$ and $\frac{F^*}{M}$ these results are similar to that of the Schaefer model in that the overall mapping of RPs is very nearly a minimal distance mapping onto the constrained set of RPs. The primary difference between Schaefer model and BH RP inference is the geometry of their limited RP spaces. Unlike the Schaefer model the BH set encourages bias







Figure 2.7: Heatplots showing the bias in RP estimation induced by model misspecification of the BH model in the high contrast simulation setting. In all cases the restricted RP-space of the BH set is shown as the black curve. (*left*) Relative bias in $\frac{B^*}{B(0)}$. (*top-right*) Bias in RP-space shown directionally. Arrows point from the location where data is generated, toward the location in the BH set where MLE projects estimated RPs. The intensity of color represents the excess bias relative to the shortest possible mapping. (*bottom*) Relative bias in F^* .



in both RPs for misspecified models even in very well informed setting.

Low Contrast Figure (2.8) shows the mapping of RPs in the low contrast simulation setting. Figures (2.8) and (2.7, *top-right*) share a common scale for the inten-



sity of color to facilitate comparison. In Figure (2.8) notice that the mildly misspecified area around the BH set produces mappings onto the BH set which resemble the minimal distance mapping seen in the high contrast setting. The primary difference in this low contrast setting, is the break point around $\frac{B^*}{B(0)} = 0.4$ above which $\frac{F^*}{M}$ is sharply underestimated.

The region of RPs where the BH model manages to recover the minimal distance mapping may be considered a “safe regime” of data types that are reasonably well modeled by a BH model. By comparison of Figure (2.8), with Figure (2.6), this safe regime of the BH model occurs for data generated for Cushing-like or BH-like production. While bias of the RPs can still become concerningly large, this region can be considered safe in the sense that even for low contrast data RP estimation under the the BH model recovers the minimal distance mapping.

Outside of this safe regime, RP estimation breaks from the minimal distance mapping at the interface between BH-Like and Ricker-Like regimes of the Schnute model (again see Figure (2.6)). The Ricker model lies along this regime interface, and represents the first model to approach the x-axis for large biomasses as γ increases. This markedly unBH-like productivity in the low information simulation setting breaks MLE inference from the minimal distance mapping and instead maps RPs to extremely low values of F^* ; consequently $\frac{B^*}{B(0)}$ is estimated

Estimated Yield Curves For Poorly Specified BH




Figure 2.9: Yield curves for data generated with $\frac{F^*}{M} = 3.48$ and $\frac{B^*}{B(0)} = 0.48$.

near the limiting value under the BH (i.e.

$\lim_{F^* \rightarrow 0} \frac{1}{F^*/M+2} = 0.5$). Similarly the set of Ricker RPs (as well as the Schaeffer set) include this trivial limiting point in common ($\frac{F^*}{M} = 0$, $\frac{B^*}{B(0)} = 0.5$).

Interestingly, in the high contrast setting this trivial mapping for highly misspecified BH models is not present. This suggests that, under a misspecified BH model, the presence of adequate information in the data to produce reasonable estimates of $\frac{F^*}{M}$, drives $\frac{B^*}{B(0)}$ below 0.5 in accordance with $\frac{B^*}{B(0)} = \frac{1}{F^*/M+2}$, even when the true $\frac{B^*}{B(0)} > 0.5$. This phenomena balances RP estimation within the constrained BH set as mediated by the information content of the data and the degree of model misspecification. When the information content in the data is too small to drive a compromised RP estimate, inference completely disregards accurate estimation of F^* in order to better estimate $\frac{B^*}{B(0)}$ by exploiting the common limiting behavior of the BH set and that of Ricker-like and Logistic-like models.

2 Discussion

Results presented here generally agree with what is known about estimating growth rate parameters (Lee et al., 2012; Conn et al., 2010; Magnusson & Hilborn, 2007). These study's appreciate the role of contrast for estimating growth rates, however they struggle to make generally extensible conclusions since they focus only on a handful of stocks that fall short of forming a random sample of the greater population of possible stock behaviors. The LHS design methods presented here are designed specifically to simulate a representative sample of stocks broadly across the space of possible RPs. Furthermore, the simulation design, taken together with the GP metamodel of productivity parameter estimates, allows this study to control the degree of model misspecification and generalize conclusions about the behavior of productivity estimation within the production model setting presented.

In the presence of contrast, F^* estimation can enjoy very low bias even for a wide range of poorly specified models; conversely in the absence of contrast F^* estimation can suffer very large bias even for slightly misspecified models. This pattern is particularly true for inference under the Schaeffer model where the geometry of the restricted RP set isolates estimation failure of F^* from $\frac{B^*}{B(0)}$. While contrast has a similar impact on F^* estimation under the BH model, the geometry of the BH RP set correlates estimation bias of F^* and $\frac{B^*}{B(0)}$. The

GP metamodeling approach reveals a more general pattern that highly informative data sets (high contrast) produces a nearly minimal distance mapping of RPs onto the constrained RP set.

In all cases when model misspecification is removed, even with weakly informative data, RP estimation is unbiased and well estimated. Thus contrast alone is not the only factor leading to inferential failure. Model misspecification is a necessary but not sufficient condition for inducing RP estimation bias. The particular RP bias present depends on the RP geometry of the fitted model and how that geometry is misspecified relative to the data. The RP mapping is then oriented to the RP geometry of the fitted model.

While the relative fishing rate parameterized in Section (5.5) captures a usefully broad spectrum of relevant fishing behaviors, it is still limiting in the amount of information that it can induce. Improved methods for quantifying contrast in fisheries data, and/or methods of discovering more informative fishing behavior, could improve this analysis. In the absence of a maximally informative dataset simulation methods will not fully describe how inference fails, but the methods presented here tell the most complete picture yet, with explicit control of the degree model misspecification, contrast, and a simulation design that allows for uniform representative data generation across biologically meaningful stocks. The results presented here suggest the conjecture that under a maximally informative dataset, RP inference with a two parameter production function will be biased in the direction a shortest distance map from the true RPs onto restricted set of RPs under the two parameter model.

Given the potential for model misspecification of RPs, a minimal distance mapping of RPs represents a best-case scenario where the total bias of RPs, when measured jointly, is minimized. That said, without recognizing the geometry of how two parameter models of productivity limit RP space this may lead to unintuitive implications in RP estimation. For example, due to the shape of the BH RP set a minimal distance mapping ensures that if there is bias in one of $\frac{B^*}{B_0}$ or F^* , there will necessarily be bias in the other RP. However under the Schaefer model, since the RP set is a constant in $\frac{B^*}{B_0}$, bias in F^* is not adulterated in the same way by bias in $\frac{B^*}{B_0}$ estimation. While models with constant RPs, such as the logistic model $\frac{B^*}{B_0} = \frac{1}{2}$ or the Fox model $\frac{B^*}{B_0} = \frac{1}{e}$, are extremely limited, they can be valuable tools for developing intuition precisely because they isolate RP estimation in their free RPs from

the correlated RP biases present in models like the BH or Ricker model.

When one considers the implications of RP bias, overestimation of RPs carries the severe implication of management recommendations potentially leading to overfishing, while underestimation of RP leads to overly conservative management. In this sense, when the true model is not known, the geometry of the BH set together with the metamodeled bias trends makes the BH model a naturally conservative estimator of RPs for most stocks. For most non-BH populations the BH model is likely to make conservative errors in its estimates of F^* and $\frac{B^*}{B_0}$. The one notable exception to the conservatism of the BH model stands for data generated in the Cushing-like regime of Schnute RPs. In this regime the BH model tends to be fairly unbiased overall, however the bias that is present for these populations tends to be overestimation in both RPs, leading to much more severe management consequences for those populations.

The RP bias trends of the Schaefer model demonstrate much less conservatism than the BH overall. For any population with $\frac{B^*}{B_0} < 0.5$, $\frac{B^*}{B_0}$ will be overestimated. When the population comes from the regime where $\frac{B^*}{B_0} > 0.5$, $\frac{B^*}{B_0}$ will be under estimated, but F^* is likely to be overestimated depending on the degree of contrast present in the data. So while the Schaefer model is an intuitive model, it tends to lead to much less conservative RP estimation.

While it is important to recognize these limitations of two parameter models of productivity, we should not solely accept conservatism as a rationale of choosing a BH model of productivity. Increasing the flexibility of the production function by moving toward three parameter models would release the underlying structural limitations (Mangel et al., 2013) that cause these RP biases in the first place. Punt and Cope (2019) considers a suite of possible three parameter curves which could be used instead of current two parameter curves. For all of their benefits, three parameter production functions have their own complicating factors, and the structure present in the Schnute model explored here makes it an intuitive bridge model for developing three parameter models going forward.

- [show a schnute fit to data?](#) (Yeakel & Mangel, 2015) Prior

- 894 • summary of σ over RP space comparing between models (PT, Schnute, Schnute DD)

895 to show areas of model breakdown.

 - 896 – miss-identifying signal for noise.
 - 897 – It happens more as the dynamics get more complex.
 - 898 – point to the full age structured models.
- 899 • show the constrained BH space over a grid of $M, \kappa, \omega, W_\infty$
- 900 • Show that the constrained spaces vary only slightly as compared with the consequences

901 of misspecifying the functional form.
- 902 • estimating these other quantities (while they can create quite different Biomass series)

903 can only do so much to improve (expand) RP inference as compared with correctly

904 modeling P .
- 905 • mapping distance as a function of contrast at (3.5, 0.5)
- 906 • for LHS grid locations show $\frac{B^*}{B_0}$ and F^* biases for grids in $M \in (0, 0.5)$ For sure in High

907 Contrast, maybe also in Low??.

908 Chapter 3

909 A Delay Differential Model

910 • Introduction

- 911 – piggy back intro off of simpleModel
- 912 – problem statement and motivation
- 913 – introduce reference point and management decision making
- 914 – new dynamics of cohorting.

915 • Methods

- 916 – state and describe model
- 917 – Reference Point Derivation
- 918 – layout data generation/space filling problem
- 919 – how far to get the math for inputting into CAS
- 920 – method of CAS.
- 921 – describe and plot ζ .
- 922 – constrained BH space (method for visualizing)
- 923 – appendix for RP CAS calculation

924 • Results

- 925 • summary of σ over RP space comparing between models (PT, Schnute, Schnute DD)
- 926 to show areas of model breakdown.
- 927 – miss-identifying signal for noise.
- 928 – It happens more as the dynamics get more complex.
- 929 – point to the full age structured models.
- 930 • Show that the constrained spaces vary only slightly as compared with the consequences
- 931 of misspecifying the functional form.
- 932 • ?Discussion?

- 933 • summary of σ over RP space comparing between models (PT, Schnute, Schnute DD)
- 934 to show areas of model breakdown.
- 935 – miss-identifying signal for noise.
- 936 – It happens more as the dynamics get more complex.
- 937 – point to the full age structured models.
- 938 • show the constrained BH space over a grid of $M, \kappa, \omega, W_\infty$
- 939 • Show that the constrained spaces vary only slightly as compared with the consequences
- 940 of misspecifying the functional form.
- 941 • estimating these other quantities (while they can create quite different Biomass series)
- 942 can only do so much to improve (expand) RP inference as compared with correctly
- 943 modeling P .

1 Introduction

- the delay model: [J. Schnute \(1985\)](#) [J. Schnute \(1987\)](#) [Fournier and Doonan \(1987\)](#).
- discrete: [Hilborn and Walters \(1992, pg. 334\)](#)
- [Walters \(2020\)](#)
- automatic accounting for cohort cycles

2 Methods

2.1 Delay Differential Model

Age structured fisheries models typically assume [Von Bertalanffy \(1938, VB\)](#) growth in length with age. To model weight the assumption of VB growth in length is composed with a power law relating length to weight, $w = al^b$. Since b is usually ~ 3 this composition of assumed functional forms typically results in a monotonically increasing sigmoidal curve of weight with age. When $b \leq 1$ weight at age takes a VB-like form with $b = 1$ resulting in an exact correspondence of simultaneous VB-growth in length and weight.

The delay model slightly abridges these relationships by directly assuming VB growth in weight as follows,

$$w(a) = w_{\infty}(1 - e^{-\kappa(a-a_0)}). \quad (3.1)$$

κ is a parameter that controls the instantaneous rate of individual growth (in weight) with age. w_{∞} is the maximum weight of individuals in the population, and $w(a)$ is the average




Figure 3.1: The typical composition of allometric weight ($b = 3$) with VB growth in length, as approximated by VB growth in weight directly.

weight of an individual at age a . The parameter a_0 controls the age at which individuals are assumed to have zero weight; by letting $a_0 < 0$ this allows fish of age zero to have positive weight. Rather than taking a sigmoidally increasing function, VB growth directly in weight results in an monotonically inceasing curve that asymptotes with a strictly decreasing growth rate with age. (only a good approximation for older ages where growth begins to decline)

Together with VB growth, the delay model is derived from the assumption that both natural mortality and fishing selectivity are separately propotional to a common heavyside step function with age. That is to say, before a threshold age of selectivity, a_s , the population is assumed not to experience any mortality whatsoever, but all fish older then a_s experience the same rate of natural mortaility. Simulaneously all fish older than a_s are equally vulnerable to fishing (i.e. knife edge selectivity at age a_s), although fishing effort may vary from through time.

Walters (2020) shows that within these assumptions the following delay differential system of equations exactly models the population dynamics of the total exploitable biomass $B(t)$ and number of individuals $N(t)$ through time.

$$\frac{dB}{dt} = w(a_s)R(B; \theta) + \kappa [w_\infty N - B] - (M + F)B \quad (3.2)$$

$$\frac{dN}{dt} = R(B; \theta) - (M + F)N \quad (3.3)$$

This formulation separates the number of individuals in the population from the biomass of the population. The dynamics of N , as seen in Eq (3.3), are very similar to that of the production models previously presented, however the role of the production function is now filled by a "recruitment" function, $R(B)$, which describes the number of new individuals recruiting into the expoitable population as a function of exploitable biomass. In turn, the biomass dynamics are coupled to the numbers dynamics by the assumption of VB growth with growth parameters appearing in Eq (3.2), converting population numbers into biomass and accounting for the growth of biomass with age.

Eq (3.2) of the above model expands the notion of biomass production into the processes of recruitment, individual growth, and maturity. The term $w(a_s)R(B; \theta)$ represents the

987 biomass of new recruits; with $w(a_s)$ representing the weight of individuals at the age of
 988 maturity, a_s , and $R(B; \theta)$ representing the number of new recruits entering the exploitable
 989 population at time t . The negative term, $(M + F)B$, represents all causes of mortality as
 990 it is applied to biomass. Finally, the term $\kappa[w_\infty N - B]$ accounts for the net growth of the
 991 existing biomass by discounting the limiting maximal individual growth rate by metabolic
 992 weight loss proportional to $B(t)$. This term, together with the delay structure in R , provides
 993 the major computational savings of the delay differential setting, as compared with full age
 994 structured models, by automatically keeping track of changes in the mean size and growth
 995 associated with changes in recruitment as cohorts mature into the population.

Often a BH functional form is assumed for the stock recruitment relationship, but any adequately flexible family of functions may model this relationship. For the sake of evaluating the adequacy of assumed BH recruitment the simulation setting below is derived for the delay model under the assumption of the generalized three parameter Schnute recruitment as follows.

$$R(B; [\alpha, \beta, \gamma]') = \alpha B(t - a_s)(1 - \beta \gamma B(t - a_s))^{\frac{1}{\gamma}} \quad (3.4)$$

996 The parameters $\theta' = [\alpha, \beta, \gamma]$ function similarly in this setting as previously described in
 997 Section (??). That said, since the delay model explicitly parses out growth in it's dynamics,
 998 these parameters only describe the net processes of larval production, and maturation into
 999 the population, where as the production model used these parameters to also model the net
 1000 effects of growth on biomass production. The γ parameter generalizes the family to model
 1001 varying degrees of decreasing recruitment for large biomasses as γ increases. The Schnute
 1002 function is exactly equivalent to BH recruitment at the special case when $\gamma = -1$, it passes
 1003 through the Ricker model as $\gamma \rightarrow 0$, and Logistic recruitment occurs when $\gamma = 1$.

1004 Since the delay model assumes knife edge selectivity, at age a_s , the term $B(t - a_s)$ appears
 1005 in R . That is to say fish recruiting into the exploitable population are the result of larval
 1006 production of biomass a_s time units in the past. This is because fishing selectivity is only
 1007 assumed to occur for fish that are at least a_s time units old and thus fish younger than a_s
 1008 are not exploitable. This waiting period requires that new recruits be the result of spawning

1009 biomass a_s time units in the past. Modeling maturity in this way results in dynamics
 1010 equations which are a system of delay differential equations as opposed to the simple ODEs
 1011 that arise in the production model setting.

1012 \sim interpretation of recruitment (larval production, recruitment) [growth external] vs.
 1013 production (larval production, recruitment, growth)

1014 • general structure: [Walters \(2020\)](#) [Hilborn and Walters \(1992, pg. 334\)](#)

1015 • growth: [Von Bertalanffy \(1938\)](#)

1016 • recruitment: [J. Schnute \(1985\)](#); [J. T. Schnute and Richards \(1998\)](#)

1017 2.2 Reference Points

1018 Deriving reference points for the delay model under Schnute recruitment is conceptually
 1019 similar to the production model setting. The additional nonlinear VB growth assumptions
 1020 along side Schnute recruitment quickly make the expressions look somewhat unweildy, al-
 1021 though analytical solutions can still be derived for most of the same quantities (although
 1022 complicated by growth parameters).

Starting from Eqs. (3.2) and (3.3), setting both $\frac{dB}{dt}$ and $\frac{dN}{dt}$ simultaneously equal to zero, and solving for B and N as a function of fishing, gives the equilibrium biomass and numbers equations.

$$\bar{B}(F) = \frac{1}{\beta\gamma} \left(1 - \left(\frac{(F+M)(F+M+\kappa)}{\alpha w(a_s)(F+M+\frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma \right) \quad (3.5)$$

$$\bar{N}(F) = \frac{\alpha \bar{B}(F)(1 - \beta\gamma \bar{B}(F))^{1/\gamma}}{F+M} \quad (3.6)$$

1023 Eq. (3.6) is just $\frac{R(\bar{B})}{F+M}$, and is coupled to $\bar{B}(F)$ where most of the dynamics appear. Eq.
 1024 (3.5) resembles Eq (3.27) from the simple production model setting although the growth
 1025 parameters κ , w_∞ and $w(a_s)$, make slight adjustments to the balance of the maximum rate
 1026 of recruitment and mortality rate to give an expression for equilibrium biomass that accounts
 1027 for the factors of individual growth.

Expressions for B_0 and B^* are attained by evaluating $\bar{B}(F)$ at $F = 0$ and $F = F^*$ respectively. Calculation of F^* typically involves maximization of equilibrium yield, $\bar{Y} = F\bar{B}(F)$. While it was not possible to analytically maximize \bar{Y} , stable numerical solutions for calculating F^* were obtained by numerically solving for the roots of the analytical derivative of equilibrium yield with respect to F . Below a greatly simplified expression for $\frac{d\bar{Y}}{dF}$ is shown; the substitution $Z = F + M$ (total mortality rate) has been made to produce a more compact expression.

$$\frac{d\bar{Y}}{dF} = \frac{1}{\beta\gamma} \left[1 - \left(\frac{Z(Z + \kappa)}{\alpha w(a_s)(Z + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma - \left(\frac{\gamma F}{\alpha w(a_s)} \right) \left(\frac{Z(Z + \kappa)}{\alpha w(a_s)(Z + \frac{\kappa w_\infty}{w(a_s)})} \right)^{\gamma-1} \left(1 + \frac{\left(\frac{\kappa w_\infty}{w(a_s)} \right) \left(\kappa - \frac{\kappa w_\infty}{w(a_s)} \right)}{\left(Z + \frac{\kappa w_\infty}{w(a_s)} \right)^2} \right) \right] \quad (3.7)$$

F^* is calculated as the numerical root, w.r.t. F , of the above expression. The numerical root is calculated using the base R uniroot function which employs a derivative free search given by [Brent \(1973\)](#).

BH Constraint

In the simple production model the BH constrained RPs are fixed to $\frac{1}{x+2}$. In the delay differential modeling setting the constrained BH RP set is complicated the growth parameters a_s and κ . Under BH recruitment these parameters of the delay model slightly influence this relationship as seen in Figure (3.12). That said, the influence of a_s and κ on RPs is still largely limited to a confined region of reference point space which resembles the $\frac{1}{x+2}$ form. In fact the confined region of RPs is bounded above by $\frac{1}{x+2}$. In Figure (3.12) notice that for high values of κ and small values of a_s (red region) the BH RP space converges to $\frac{1}{x+2}$ as derived in the simple production model setting. The opposite limit with low values of κ and high values a_s




Figure 3.2: The space of BH RPs for the delay model as a function of κ and a_s . The RP space is plotted for 80×80 combinations of $\kappa \in [0.1, 2]$ and $a_s \in [0.1, 10]$. The color drawn is the resulting value of $w(a_s)$ mapped between blue and red. $\frac{1}{x+2}$ is plotted in black for reference.

1055 (blue region) depresses RPs away from $\frac{1}{x+2}$.

1056 2.3 Delay Differential Integration

1057 The delay model belongs to a class of differential equations known as delay differential
1058 equations (DDE). The delay arises from the $B(t - a_s)$ terms found in the recruitment
1059 function. Solving DDEs require special care which depends on the nature of the time delay.
1060 The addition of time-varying delays, many different delays, or very small delays (delays
1061 below the step size of the numerical integrator) results in some of the more challenging
1062 settings for solving DDEs. However with a single stationary model of the age of selectivity,
1063 the delay model in this setting represents one of the most straight forward DDE structures.
1064 The most numerically challenging case presented here arises in the case of the limiting
1065 production model when $a_s \rightarrow 0$ while $\kappa \rightarrow \infty$. That said the limiting production model can
1066 be approximated for values of $a_s \approx 0.1$, and it was straightforward to ensure that the step
1067 size of the integrator remained reasonably below 0.1.

1068 The DDE presented here is integrated with the initial values fixed at B_0 and N_0 as given
1069 by Eqs. (3.5) and (3.6) with $F = 0$ at any given configuration of θ and growth parameters.
1070 The system given in Eqs. (3.2) and (3.3) are then solved numerically using the implicit
1071 Livermore Solver (lsode) as implemented in the `dede` function of the R package `deSolve`
1072 (Soetaert et al., 2010). The `dede` solver provides many methods for integrating DDEs, but
1073 lsode was chosen because it is an implicit method that runs relatively quickly with a relatively
1074 smaller footprint in system memory as compared with other methods. The radau method
1075 was also tried in more computationally challenging settings with good results (albeit running
1076 more slowly than lsode). Ultimately the simulated parameter space did not produce DDEs
1077 that require the more expensive radau integrator to solve accurately.

1078 2.4 Simulation Design

1079 Similarly as previously described in Section (5) the relationship between RPs $\mapsto \theta$ cannot be
1080 fully expressed analytically for the Schnute delay model. However, just as in the production
1081 model setting, simulation only requires enough knowledge of these mappings to gather a list
1082 of (α, β, γ) tuples and the corresponding RPs in some reasonable space-filling design over

1083 RP space.

1084 In the delay model a partial mapping for $(F^*, B_0) \mapsto (\alpha(\cdot, \gamma), \beta(\cdot, \cdot, \gamma))$ can be derived
 1085 analytically in terms of RPs and γ . The substitution $Z^* = F^* + M$ is made where F^* and
 1086 M appear together to produce a more compact expression.

$$\alpha = \left[\left(\frac{Z^*(Z^* + \kappa)}{w(a_s)(Z^* + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma + \left(\frac{\gamma F^*}{w(a_s)} \right) \left(\frac{Z^*(Z^* + \kappa)}{w(a_s)(Z^* + \frac{\kappa w_\infty}{w(a_s)})} \right)^{\gamma-1} \left(1 + \frac{\left(\frac{\kappa w_\infty}{w(a_s)} \right) \left(\kappa - \frac{\kappa w_\infty}{w(a_s)} \right)}{(Z^* + \frac{\kappa w_\infty}{w(a_s)})^2} \right) \right]^{\frac{1}{\gamma}} \quad (3.8)$$

$$\beta = \frac{1}{\gamma B_0} \left(1 - \left(\frac{M(M + \kappa)}{\alpha w(a_s)(M + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma \right) \quad (3.9)$$

Above Eq. (3.8) results from setting Eq. (3.31) equal to zero and solving for α , and
 Eq. (3.9) results from solving the $\bar{B}(0)$ expression, as derived from Eq. (3.5), for β . The
 system is completed by further working with the $\frac{\bar{B}(F^*)}{\bar{B}(0)}$ expression, as seen below, to identify
 γ .

$$\frac{B^*}{B_0} = \frac{1 - \left(\frac{(F^* + M)(F^* + M + \kappa)}{\alpha w(a_s)(F^* + M + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma}{1 - \left(\frac{M(M + \kappa)}{\alpha w(a_s)(M + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma} \quad (3.10)$$

1087 The system formed by collecting Eqs. (3.8), (3.9), and (3.10) can be navigated similarly
 1088 to Eq. (3.33) in the Schnute production model setting. For a population experiencing
 1089 natural mortality M , VB growth with paramters κ and w_∞ , and age of selectivity a_s the
 1090 above system can fully specify α and β for a given γ , by fixing F^* , B_0 , and $\frac{B^*}{B_0}$. For a given γ
 1091 a cascade of closed form solutions for α and β can be obtained, just as in Section (5). First
 1092 $\alpha(\gamma)$ can be computed, and then $\beta(\alpha(\gamma), \gamma)$ can be computed. If $\alpha(\gamma)$ is filled back into the
 1093 expression for $\frac{B^*}{B_0}$, the system collapses into a single onerous expression for $\frac{B^*}{B_0}(\alpha(\gamma), \gamma)$. For
 1094 brevity, define the function $\zeta(\gamma) = \frac{B^*}{B_0}(\alpha(\gamma), \gamma, F^*, M)$ based on Eq. (3.10).

1095 Again rather than inverting $\zeta(\gamma)$ for γ , γ is the sampled so that the overall simulation
 1096 design is space filling as described in Section (5 .3). Given the sampled γ , the cascade of
 1097 $\alpha(\gamma)$, and then $\beta(\alpha(\gamma), \gamma)$, can be computed, and the Schnute delay model is fully defined
 1098 by a given $(\frac{F^*}{M}, \frac{B^*}{B_0})$. While conceputally this framing is similar to the Schnute production
 1099 model, the analytical expressions are more complex, and numerically trecherous, since growth
 1100 parameters appear explicitly here. Other ways of navigating the RPs $\mapsto \theta$ system are possible,

1101 but for the sake of numerical stability this strategy has proven the most reliably accurate by
 1102 limiting exposure to numerical error propagation.

1103 Each design location defines a complete Schnute delay differential model with the given
 1104 RP values. Indices of abundance are simulated from the Schnute model at each design
 1105 location, a small amount of residual variation, $\sigma = 0.01$, is added to the simulated index,
 1106 and the data are then fit with a misspecified BH model. The design captures various degrees
 1107 of model misspecification relative to the BH model, so as to observe the effect of recruitment
 1108 misspecification upon RP inference.

1109 point to catch, and LHS design, and Metamodel.

1110 2 .5 Parmeter Estimation

- 1111 • I use B only here
- 1112 • quick statement of inference, and reference to previous section

Let I_t , $t \in \{1, 2, 3, \dots, T\}$, be a series of indicies of abundance, proportional to biomass, as simulated from the Schnute Delay model. These data are modelled with the following log-normal observation model that has been intentionally constrained to BH recruitment,

$$I_t \sim LN(qB_t(\boldsymbol{\theta}, \boldsymbol{\phi}), \sigma^2). \quad (3.11)$$

1113 $B_t(\boldsymbol{\theta}, \boldsymbol{\phi})$ is the biomass solution of the BH constrained DDE system. The BH constraint
 1114 isimplemented by fixing $\gamma = -1$ so that $\boldsymbol{\theta}' = [\alpha, \beta, \gamma = -1]$. $\boldsymbol{\phi}$ is a vector of growth and
 1115 maturity parameters, $\boldsymbol{\phi}' = [\kappa, w_\infty, a_0, a_s]$. The nuisance parameter q models the proportion-
 1116 ality constant of the index with process biomass, and σ^2 models residual variation of the
 1117 index.

1118 In this setting, $\boldsymbol{\phi}$ and q are fixed to focus on the inferential affects of model misspecifi-
 1119 cation on recruitment parameters and RPs. Without an explicite mechanism for the delay
 1120 model to incorporate age data, under the BH model $\boldsymbol{\phi}$ is not well informed and would typically
 1121 be estimated externally for data limited stocks. Under BH recruitment $\boldsymbol{\phi}$ can only slightly
 1122 impact RPs as seen in Figure (3.12).

1123 σ^2 and θ are reparameterized to the log scale and fit via MLE. Reparameterizing the

parameters to the log scale improves the reliability of optimization, in addition to facilitating the use of Hessian information for estimating MLE standard errors. Given that the biological parameters enter the likelihood via a nonlinear differential equation, and further the parameters themselves are related to each other nonlinearly, the likelihood function can often be difficult to optimize. A hybrid optimization scheme is used to maximize the log likelihood to ensure that a global MLE solution is found. The R package GA (Scrucca, 2013, 2017) is used to run a genetic algorithm to explore parameter space globally. Optimization periodically jumps into the L-BFGS-B local optimizer to refine optima within a local mode. The scheme functions by searching globally, with the genetic algorithm, across many initial values for starting the local gradient-based optimizer. The genetic algorithm serves to iteratively improve hot starts for the local gradient-based optimizer. Additionally, optimization is only considered to be converged when the optimum results in an invertible Hessian at the found MLE.

- fixed $M = 0.2$, $a_0 = -1$, $w_\infty = 1$
- play with κ and age of selectivity a_s

Numbers Indices

While not utilized here, age structured models may commonly model indices as proportional to numbers rather than (or simultaneously to) biomass. When solving the DDE, Eq. (3.3) points out that the full DDE solution will expose a numbers solution simultaneously with a biomass solution that may be used for these purposes. These solutions are often quite similar since the main driver of process behavior comes from the form of R which is shared among N and B . However, it is common on the west coast of the US that indices derived from commercial fisheries are measured as weights while indices derived from recreational fisheries are often measured as counts. If a numbers index, J_t , is observed alongside the previously mentioned biomass index, the following likelihood component is often added as a conditionally independent component of the likelihood,

$$J_t \sim LN(pN_t(\boldsymbol{\theta}, \boldsymbol{\phi}), \tau^2). \quad (3.12)$$

1140 $N_t(\boldsymbol{\theta}, \boldsymbol{\phi})$ is the numbers solution of the DDE system. $\boldsymbol{\theta}$ and $\boldsymbol{\phi}$ are the productivity and
 1141 growth parameters shared in common with the biomass component. p and τ^2 are then the
 1142 analogous proportionality constant and residual variation of the numbers index respectively.

1143 2.6 GP Metamodel

1144 point to catch, and LHS design, and Metamodel.

1145 3 Results




Figure 3.3: Biomass dynamics of BH (*left*), Ricker (*center*), and Logistic (*right*) delay differential models in the low contrast simulation setting. In all cases $\alpha = 1.2$ and β is chosen so that each model shares the same B_{MSY} within each given γ .

1146 Figure (3.4) demonstrates a range of behaviors that the Schnute delay model can display.
 1147 The three special cases of $\gamma = -1$ (BH), $\gamma \rightarrow 0$ (Ricker), and $\gamma = 1$ (Logistic) recruit-
 1148 ment are shown with varying configurations of growth ranging from the simple (no growth)
 1149 production model setting when $a_s \rightarrow 0$ and $\kappa \rightarrow \infty$, to an opposing case ($a_s = 10$ and
 1150 $\kappa = 0.1$) representing extremely lagged selectivity and slow individual-growth. By reference
 1151 with Figure (3.12), notice that the latter individual-growth/maturity configuration repre-
 1152 sents a RP set in the deep blue regim, thus this case represents dynamics where RPs are
 1153 heavily influenced by individual-growth/maturity, rather than solely determined by the as-
 1154 sumed functional form of recruitment as is the case with the simple production model. The
 1155 case $a_s = 2$ and $\kappa = 0.1$ is shown as an example of dynamics where RPs are more moderatly
 1156 influenced by individual-growth/maturity.

1157 Notice under the extreme $a_s = 10$ and $\kappa = 0.1$ setting, biomass comes into equilibrium at
 1158 B_{MSY} as oscillating curves. This effect is most noticable in the Logistic model here due to
 1159 its steeply right leaning yeild curve interacting with the large lag in selectivity as the sudden
 1160 onset of fishing produces a shock that oscillates over the steepest regions of the yeild curve.
 1161 One may exagorate the oscillations of a BH model similarly, in this heavily lagged setting,
 1162 by shocking the population over the steepest portion of its yeild curve (a sudden release in
 1163 fishing applied to a heavily fished population at low equilibrium biomass).

1164 • RP curve triptic (BH, Ricker, Logistic) each at the three examples

1165 3 .1 Simple Production Model Limit




Figure 3.4: Biomass dynamics of BH (*left*), Ricker (*center*), and Logistic (*right*) delay differential models in the low contrast simulation setting. In all cases $\alpha = 1.2$ and β is chosen so that each model shares the same B_{MSY} within each given γ .




Figure 3.5:

3.2 Dramatic Growth




Figure 3.6:

3.3 Oscillatory Growth Influence




Figure 3.7:




Figure 3.8:

- Selectivity and Growth cannot account for misspecification of Recruitment.




Figure 3.9:

- Selectivity and Growth are influenced by the proper specification of recruitment.
If properly specified Selectivity and Growth can be estimated, along with all three parameters of the Schnute model.
- show production model limit (contrast
 - $a_s \rightarrow 0$: instant maturity
 - $\kappa \rightarrow \infty$: recruit as an adult ()
- describe second order shapes of growth/maturity (and cause)
 - weight of recruits => scaling biomass (q , β , and w_∞)
 -
- describe RP bias
- flat




Figure 3.10: This is only a test




Figure 3.11: This is only a test

1180

why not here?




Figure 3.12. This is only a test

4 Introduction

The most fundamental model in modern fisheries management is the surplus-production model. These models focus on modeling population growth via nonlinear parametric ordinary differential equations (ODE). Key management quantities called reference points (RPs) are commonly derived from the ODE equilibrium equations and depend upon the parameterization of biomass production. Two-parameter forms of the production function have been shown to limit the theoretical domain of RPs (Mangel et al., 2013). The limited RP-space of two parameter models are a major source of model misspecification for RPs and thus induce bias in RP estimation. The behavior of RP estimation bias is not well understood and as a result often underappreciated. A metamodeling approach is developed here to describe RP biases and explore mechanisms of model failure under the most common two parameter models.

Data for a typical surplus-production model comes in the form of an index of abundance through time which is assumed to be proportional to the reproducing biomass for the population of interest. The index is often observed alongside a variety of other known quantities, but at a minimum, each observed index will be observed in the presence of some known catch for the period. Figure (3.13) shows the classic Namibian Hake dataset exemplifying the form.

Indices are assumed to have multiplicative log-normal errors, and thus the following observation model arises naturally,

$$I_t = qB_t e^\epsilon \quad \epsilon \sim N(0, \sigma^2). \quad (3.13)$$

Above q is often referred to as the “catchability parameter”; it serves as the proportionality constant mapping between the observed index of abundance and biomass. σ^2 models residual variation. Biologically speaking q and σ^2 are often treated as nuisance parameters with the “biological parameters” entering the model through a process model on biomass.

Biomass is assumed to evolve as an ODE; in this case I focus on the following form

$$\frac{dB}{dt} = P(B(t); \boldsymbol{\theta}) - Z(t)B(t). \quad (3.14)$$




Figure 3.13: *left*: An index of abundance data, catch per unit effort (CPUE), for Namibian Hake from 1965 to 1987 (Hilborn & Mangel, 1997). *right*: The associated catch data for Namibian Hake over the same time period.

Here biomass is assumed to change in time by two processes, net production of biomass into the population, $P(B)$, and various sources of biomass removal, Z , from the population.

Firstly, the population grows through a production function, $P(B)$. Production in this setting is defined as the net biomass increase due to all reproduction and maturation processes. The production function is assumed to be a parametric (generally non-linear) function relating the current biomass of the population to an aggregate production of biomass.

Secondly, the population decreases as biomass is removed by various sources that are assumed to remove biomass linearly with biomass. Above, $Z(t)$, is an aggregate rate of removal. When the fishing rate, $F(t)$, is the only source of removal $Z(t) = F(t)$, however often models will also included other linear terms in $Z(t)$. Commonly the rate of “natural mortality”, M , is also included as an additional term so that $Z(t) = M + F(t)$.

From a management perspective a major goal of modeling is to accurately infer a quantity known as *maximum sustainable yield* (MSY). One could maximize simple yield at a particular moment in time (and only for that moment) by fishing all available biomass in that moment. This strategy is penny-wise but pound-foolish (not to mention ecologically devastating) since it doesn’t leave biomass in the population to reproduce in the future. We seek to fish in a way that allows (or even encourages) future productivity in the population. This is accomplished by maximizing the equilibrium level of catch over time. Equilibrium yield is considered by

replacing the steady state biomass (\bar{B}) in the assumed form for catch, so that $\bar{Y} = F\bar{B}(F)$, where $\bar{\cdot}$ indicates a value at steady state. MSY is found by maximizing $\bar{Y}(F)$ with respect to F , and F^* is the fishing rate at MSY. Going forward let $*$ decorate any value derived under the condition of MSY.

Fisheries are very often managed based upon reference points which serve as simplified heuristic measures of population behavior. The mathematical form of RPs depends upon the model assumptions through the production function. While a number of different RPs exist which describe the population in different (but related) ways, the most common RPs revolve around the concept of MSY (or robust ways of measuring MSY (Hilborn, 2010; Punt et al., 2016)). Here the focus is primarily on the RPs $\frac{B^*}{B(0)}$ and F^* ($\frac{F^*}{M}$ when appropriate) for their pervasive use in modern fisheries (Punt & Cope, 2019).

F^* is the afore mentioned fishing rate which results in MSY. $\frac{B^*}{B(0)}$ is the depletion of the stock at MSY. That is to say $\frac{B^*}{B(0)}$ describes the fraction of the unfished population biomass that will remain in the equilibrium at MSY. In general $F^* \in \mathbb{R}^+$ and $\frac{B^*}{B(0)} \in (0, 1)$, however under the under the assumption of a two parameter production function production models will be structurally unable to capture the full theoretical range of RPs.

Many of the most commonly used production functions depend only on two parameters. For example, the Schaefer model depends only on the biological parameters r and K , and limits RP inference so that under the Schaefer model $\left(F^*, \frac{B^*}{B(0)}\right) \in \left(\mathbb{R}^+, \frac{1}{2}\right)$. The two parameter Fox model (Fox Jr., 1970) limits $\left(F^*, \frac{B^*}{B(0)}\right) \in \left(\mathbb{R}^+, \frac{1}{e}\right)$. Similarly the two parameter Cushing (Cushing, 1971), Beverton-Holt (Beverton & Holt, 1957, BH) and Ricker (Ricker, 1954) production functions do not model the full theoretical space of RPs (Mangel et al., 2013; Yeakel & Mangel, 2015).

The bias-variance trade-off (Ramasubramanian & Singh, 2017) makes it clear that the addition of a third parameter in the production function will necessarily reduce estimation bias. However the utility of this bias reduction is still under debate because the particular mechanisms and behavior (direction and magnitude) of these biases for key management quantities are not fully understood or described. Lee et al. (2012) provides some evidence that estimation of productivity parameters are dependent on biomass contrast as well as model specification. Conn et al. (2010) comes to similar conclusions via calibration modeling

1252 techniques. These studies indicate important factors that contribute to inferential failure.
 1253 However they do not offer mechanisms of model failure, nor do their experimental designs
 1254 allow for the control of different types of model misspecification.

1255 In this study I consider the behavior of inference when index data are simulated from
 1256 three parameter PT and Schnute production models, but the simulated data are fit using
 1257 intentionally misspecified two parameter logistic or BH production models. The work begins
 1258 with a derivation of RPs under the three parameter models. A method is then presented
 1259 for generating simulation designs based on the parametric form of RPs which serves as a
 1260 control on the nature of simulated model misspecification. Finally a Gaussian Process (GP)
 1261 metamodel ([Gramacy, 2020](#)) is constructed for exploration and analysis of RP biases.

1262 A key insight of this approach is that bias is considered broadly across RP-space to
 1263 uncover patterns and correlations between RPs. The GP metamodel is explicit about trade-
 1264 offs between RPs so as to inform the full utility of reducing bias, as well as to suggest
 1265 mechanisms for understanding what causes bias. Further, the effect of contrast on estimation
 1266 is considered together with model misspecification.

1267 5 Methods

1268 5.1 Pella-Tomlinson Model

The three parameter Pella-Tomlinson (PT) family has a convenient form that includes, among others ([Fox Jr., 1970](#); [Rankin & Lemos, 2015](#)), the logistic production function as a special case. PT production function is parameterized so that $\boldsymbol{\theta} = [r, K, \gamma]$ and the family takes the following form,

$$P_p(B; [r, K, \gamma]) = \frac{rB}{\gamma - 1} \left(1 - \left(\frac{B}{K} \right)^{(\gamma-1)} \right). \quad (3.15)$$

1269 γ is a parameter which breaks PT out of the
 1270 restrictive symmetry of the logistic curve. In gen-
 1271 eral $\gamma \in (1, \infty)$, with the logistic model appear-
 1272 ing in the special case of $\gamma = 2$, and the Fox
 1273 model appearing as a limiting case as $\gamma \rightarrow 1$. The
 1274 parameter r controls the maximum reproductive
 1275 rate of the population in the absence of compe-
 1276 tition for resources (i.e. the slope of production
 1277 function at the origin). K is the so called "car-
 1278 rying capacity" of the population. In this con-
 1279 text the carrying capacity can be formally stated
 1280 as steady state biomass in the absence of fishing
 1281 (i.e. $\bar{B}(0) = K$). In Figure (3.14) PT recruitment
 1282 is shown for a range of parameter values so as to
 1283 demonstrate the various recruitment shapes that
 1284 can be achieved by PT recruitment.

1285 While the form of the PT curve produces
 1286 some limitations (Fletcher, 1978), importantly
 1287 the introduction of a third parameter allows enough flexibility to fully describe the space
 1288 of reference points used in management. To see this, the reference points are analytically
 1289 derived for the PT model below.

1290 PT Reference Points

1291 With $B(t)$ representing biomass at time t , under PT production, the dynamics of biomass
 1292 are defined by the following ODE,

$$\frac{dB}{dt} = \frac{rB}{\gamma - 1} \left(1 - \left(\frac{B}{K} \right)^{\gamma-1} \right) - FB. \quad (3.16)$$

An expression for the equilibrium biomass is attained by setting Eq (3.16) equal to zero,
 and rearranging the resulting equation to solve for B . Thinking of the result as a function




Figure 3.14: The Pella-Tomlinson production function plotted across a variety of parameter values. The special cases of Logistic production is shown in black, and the left-leaning and right-leaning regimes are shown in blue and red respectively.

of F gives,

$$\bar{B}(F) = K \left(1 - \frac{F(\gamma - 1)}{r} \right)^{\frac{1}{\gamma-1}}. \quad (3.17)$$

At this point it is convenient to notice that $\bar{B}(0) = K$. The expression for B^* is given by evaluating Eq (3.17) at F^* . To get an expression for F^* , the equilibrium yield is maximized with respect to F ,

$$F^* = \operatorname{argmax}_F F \bar{B}(F). \quad (3.18)$$

In the case of PT production this maximization can be done analytically, by differentiating the equilibrium yield with respect to F as follows,

$$\frac{d\bar{Y}}{dF} = \bar{B}(F) + F \frac{d\bar{B}}{dF} \quad (3.19)$$

$$\frac{d\bar{B}}{dF} = -\frac{K}{r} \left(1 - \frac{F(\gamma - 1)}{r} \right)^{\frac{1}{\gamma-1}-1}. \quad (3.20)$$

Setting Eq (3.19) equal to 0, substituting $\bar{B}(F)$ and $\frac{d\bar{B}}{dF}$ by Equations (3.17) and (3.20) respectively, and solving for F produces the following expression for the fishing rate required to produce MSY,

$$F^* = \frac{r}{\gamma} \quad (3.21)$$

Plugging the above expression for F^* back into Eq (3.17) gives the following expression for biomass at MSY,

$$B^* = K \left(\frac{1}{\gamma} \right)^{\frac{1}{\gamma-1}}. \quad (3.22)$$

The above derived expressions for $\bar{B}(0)$, B^* , and F^* can then be used to build a specific analytical form for the biological reference points in terms of only productivity parameters.

$$F^* = \frac{r}{\gamma} \qquad \frac{B^*}{\bar{B}(0)} = \left(\frac{1}{\gamma} \right)^{\frac{1}{\gamma-1}} \quad (3.23)$$

1296 Simulation

Generating simulated indices of abundance from the PT model requires inverting the relationship between $\left(F^*, \frac{B^*}{B(0)}\right)$, and (r, γ) . It is not generally possible to analytically invert this relationship for many three parameter production functions (Punt & Cope, 2019; J. T. Schnute & Richards, 1998). Most three parameter production functions lead to RPs that require expensive numerical methods to invert; more over the numerical inversion procedure can often be unstable. That said, for the case of PT this relationship is analytically invertible, and leads to the following relationship

$$r = \gamma F^* \qquad \gamma = \frac{W\left(\frac{B^*}{B(0)} \log\left(\frac{B^*}{B(0)}\right)\right)}{\log\left(\frac{B^*}{B(0)}\right)}. \quad (3.24)$$

1297 Above W is the Lambert product logarithm function. More details about this derivation,
1298 and the Lambert product logarithm, are given in Appendix (7).

1299 Using Eq. (3.24) to obtain production parameters, a PT production model can be fully
1300 defined for any combination of the RPs F^* and $\frac{B^*}{B(0)}$. Since K does not enter the RP
1301 calculation its value is fixed arbitrarily at 10000.

1302 Indices of abundance are simulated from the three parameter PT production model
1303 broadly over the space of F^* and $\frac{B^*}{B(0)}$ via a space filling design as described in Section
1304 (5 .3). A small amount of residual variation, $\sigma = 0.01$, is added to the simulated index, and
1305 these data are then fit with a Schaefer model, at various degrees of misspecification, so as to
1306 observe the effect of productivity model misspecification upon RP inference.

1307 5 .2 Schnute Model

The Schnute production function is a three parameter generalization of many of the most common two parameter production functions (Deriso, 1980; J. Schnute, 1985). It can be written in the following form, with parameters α , β , and γ ,

$$P_s(B; [\alpha, \beta, \gamma]) = \alpha B(1 - \beta\gamma B)^{\frac{1}{\gamma}}. \quad (3.25)$$

1308 The BH and Logistic production func-
 1309 tions arise when γ is fixed to -1 or 1 respec-
 1310 tively. The Ricker model is a limiting case
 1311 as $\gamma \rightarrow 0$. For $\gamma < -1$ a family of strictly in-
 1312 creasing Cushing-like curves arise, culminat-
 1313 ing in linear production as $\gamma \rightarrow -\infty$. These
 1314 special cases form natural regimes of simi-
 1315 larly behaving production functions as seen
 1316 in Figure (3.15).

1317 The behavior of RP inference under the
 1318 BH model is of particular interest due to the
 1319 overwhelming popularity of the BH assump-
 1320 tion in fisheries models. Since Schnute pro-
 1321 duction models can represent a quantifiably

1322 wide variety of possible productivity behaviors, they present an ideal simulation environment
 1323 for inquiry of the reliability of inference under the BH assumption.

Under Schnute production, biomass dynamics evolve according to the following ODE,

$$\frac{dB}{dt} = P_s(B; \theta) - (M + F)B. \quad (3.26)$$

1324 This equation largely takes the same form as previously described, except that P_s is the
 1325 Schnute production function and natural mortality, M , is modeled explicitly here. Natural
 1326 mortality models the instantaneous rate of mortality from all causes outside of fishing. Ex-
 1327 plicitly modeling natural mortality is not only a typical assumption of fisheries models, but
 1328 is also key to the making RPs well defined over the relevant domain of γ .

The derivation of RPs under Eq. (3.26) follows a similar logic as under the PT model. An expression for equilibrium biomass is attained by setting $\frac{dB}{dt} = 0$ and rearranging the resulting expression to solve for B

$$\bar{B}(F) = \frac{1}{\gamma\beta} \left(1 - \left(\frac{M + F}{\alpha} \right)^\gamma \right). \quad (3.27)$$




Figure 3.15: The Schnute production function plotted across a variety of parameter values. Regimes of similarly behaving curves are grouped by color.

The above expression quickly yields B_0 , B^* by evaluation at $F = 0$ and F^* respectively,

$$B_0 = \frac{1}{\gamma\beta} \left(1 - \left(\frac{M}{\alpha} \right)^\gamma \right) \quad (3.28)$$

$$\frac{B^*}{B_0} = \frac{1 - \left(\frac{M+F^*}{\alpha} \right)^\gamma}{1 - \left(\frac{M}{\alpha} \right)^\gamma}. \quad (3.29)$$

Attaining an expression for F^* requires maximization of equilibrium yield, $\bar{Y} = F\bar{B}(F)$, with respect to F . Analytically maximizing proceeds by differentiating \bar{Y} to produce

$$\frac{d\bar{Y}}{dF} = \bar{B}(F) + F \frac{d\bar{B}}{dF} \quad (3.30)$$

$$\frac{d\bar{B}}{dF} = -\frac{1}{\beta} \left(\frac{\left(\frac{M+F}{\alpha} \right)^\gamma}{F+M} \right). \quad (3.31)$$

Setting $\frac{d\bar{Y}}{dF} = 0$, filling in the expressions for $\bar{B}(F)$ and $\frac{d\bar{B}}{dF}$, then rearranging to solve for F^* is less yielding here than it was in the case of the PT model. This procedure falls short of providing an analytical solution for F^* directly in terms of θ , but rather shows that F^* must respect the following expression,

$$0 = \frac{1}{\gamma} - \left(\frac{1}{\gamma} + \frac{F^*}{F^* + M} \right) \left(\frac{F^* + M}{\alpha} \right)^\gamma. \quad (3.32)$$

The lack of an analytical solution here is understood. [J. T. Schnute and Richards \(1998](#), pg. 519) specifically points out that F^* cannot be expressed analytically in terms of productivity parameters, but rather gives a partial analytical expression for the inverse relationship. Although parameterized slightly differently, [J. T. Schnute and Richards \(1998\)](#) derives expressions for α and β as a function of RPs and γ .

Since RPs are left without a closed form expression, computing RPs from productivity parameters amounts to numerically solving the system formed by collecting the expressions (3.32), (3.28), and (3.29).

Simulation

For the purposed of simulation, it is not necessary to completely know the precise relationships mapping RPs $\mapsto \theta$ or $\theta \mapsto$ RPs. Simulation only requires enough knowledge of these

1340 mappings to gather a list of (α, β, γ) tuples, for data generation under the Schnute model,
 1341 and the corresponding RPs in some reasonable space-filling design over RP space.

Similarly to [J. T. Schnute and Richards \(1998\)](#), expressions (3.32) and (3.28) are solved for α and β respectively. This leads to the partial mapping $(F^*, B_0) \mapsto (\alpha(\cdot, \gamma), \beta(\cdot, \cdot, \gamma))$ in terms of RPs and γ . By further working with Eq. (3.29), to identify γ , the following system is obtained,

$$\begin{aligned}\alpha &= (M + F^*) \left(1 + \frac{\gamma F^*}{M + F^*} \right)^{1/\gamma} \\ \beta &= \frac{1}{\gamma B_0} \left(1 - \left(\frac{M}{\alpha} \right)^\gamma \right) \\ \frac{B^*}{B_0} &= \frac{1 - \left(\frac{M + F^*}{\alpha} \right)^\gamma}{1 - \left(\frac{M}{\alpha} \right)^\gamma}.\end{aligned}\tag{3.33}$$

1342 For a population experiencing natural mortality M , by fixing F^* , B_0 , and $\frac{B^*}{B_0}$ the above
 1343 system can fully specify α and β for a given γ . Notice for a given γ a cascade of closed
 1344 form solutions for α and β can be obtained. First $\alpha(\gamma)$ can be computed, and then
 1345 $\beta(\alpha(\gamma), \gamma)$ can be computed. If $\alpha(\gamma)$ is filled back into the expression for $\frac{B^*}{B_0}$, the system
 1346 collapses into a single onerous expression for $\frac{B^*}{B_0}(\alpha(\gamma), \gamma)$. For brevity, define the function
 1347 $\zeta(\gamma) = \frac{B^*}{B_0}(\alpha(\gamma), \gamma, F^*, M)$ based on Eq. (3.29).

1348 Inverting $\zeta(\gamma)$ for γ , and computing the cascade of $\alpha(\gamma)$, and then $\beta(\alpha(\gamma), \gamma)$, fully defines
 1349 the Schnute model for a given $(\frac{F^*}{M}, \frac{B^*}{B_0})$. However inverting ζ accurately is extremely difficult.
 1350 Inverting ζ analytically is not feasible, and numerical methods for inverting ζ are unstable
 1351 and can be computationally expensive. Rather than numerically invert precise values of $\zeta(\gamma)$,
 1352 γ is sampled so that the overall simulation design is space filling as described in Section (5
 1353 .3).

1354 Each design location defines a complete Schnute production model with the given RP
 1355 values. Indices of abundance are simulated from the Schnute model at each design location,
 1356 a small amount of residual variation, $\sigma = 0.01$, is added to the simulated index, and the data
 1357 are then fit with a misspecified BH production model. The design at large captures various
 1358 degrees of model misspecification relative to the BH model, so as to observe the effect of
 1359 productivity model misspecification upon RP inference.

5.3 Latin Hypercube Sampling

The goal of space filling design in this setting is to extend the notion of the random sample (and its desirable parameter estimation properties) across the simulated RP domain so as to represent the simulated space as well as possible (Gramacy, 2020). The simple random sample is the gold standard of classical unbiased parameter estimation, however simple randomness is patchy, often sampling some regions of design space quite densely, while leaving other regions of design space empty. Space filling designs aim to preserve (or enhance) parameter estimation properties across the simulated domain (Devon Lin & Tang, 2015; Stein, 1987), while constraining samples to be spaced in some notion of spread over the entire space. Latin hypercube sampling (McKay et al., 2000, LHS) is among the most foundational of space filling designs used in computer experiments.

A LHS of size n , in the 2 dimensional space defined by RPs, distributes samples so as to spread points across a design region in a broadly representative way. A LHS design extends the notion of a univariate random uniform sample across multiple dimensions so that each margin of the design space enjoys a uniform distribution.

LHS designs achieve this notion of uniformity by first partitioning each dimension of the design space into regular grids of size n . By intersecting the grids of each dimension, cells are produced that evenly partition the design space. In two dimensions n^2 cells are produced, from which a total of n samples are taken. Crucially only one sample is taken from a given element of each grid in each dimension so as to reduce clumping of the n samples across the design space.




Figure 3.16: LHS grids. Intersecting \mathcal{F} and \mathcal{B} produces n^2 cells; a particular cell $\mathcal{F}_i \times \mathcal{B}_j$ is shown in grey. Maybe just show points.

PT Design

Letting \mathcal{F} and \mathcal{B} be regular grids, of size $n = 100$, on $F^* \in (0.1, 0.7)$ and $\frac{B^*}{B_0} \in (0.2, 0.6)$ respectively, a LHS design of size 100 is collected among the cells produced by $\mathcal{F} \times \mathcal{B}$.

Each of the sampled LHS design locations represent a unique PT model with the sampled RP values. Since the relationship mapping RPs analytically to productivity parameters can be found for the PT model, LHS designs the the PT model are computed directly in RP space and Eq. (3.24) is used to map the sampled RP design locations to PT productivity parameters.

Schnute Design

Due to the lack of an analytical relationship mapping RPs $\mapsto \theta$, analogous to the PT model's Eq. (3.24), producing a LHS design over Schnute RPs requires a more tactful approach. The structured relationship between the RPs and productivity parameters, described in Section (5), allows an approximate LHS to be obtained by a careful navigation of the system of equations seen in Eq. (3.33).

Under the Schnute model, let \mathcal{F} and \mathcal{B} represent regular grids on $\frac{F^*}{M} \in (0.25, 4)$ and $\frac{B^*}{B_0} \in (0.15, 0.7)$ respectively which can serve as the scaffolding for computing an approximate LHS

Since it is not practical to invert $\zeta(\gamma)$, a uniform sample in $\frac{B^*}{B_0}$ can be obtained by modeling γ as a random variable, with realization γ^* , and thinking of $\zeta(\gamma)$ as its cumulative distribution function (CDF). The aim is to model γ as an easily sampled random variable with a CDF that closely approximates ζ , so that $\zeta(\gamma^*) \sim U(\zeta_{min}, 1)$ as closely as possible. There may be many good models for the distribution of γ , but in this setting the

Given B_0 , M , and F^* :

- 1) Draw $\gamma^* \sim \gamma|F^*, M$.
- 2) Compute $\frac{B^*}{B_0} = \zeta(\gamma^*)$
- 3) Compute $\alpha^* = \alpha(\gamma^*, F^*, M)$
- 4) Compute $\beta^* = \beta(\alpha^*, \gamma^*, M, B_0)$

Figure 3.17: An outline of the sampling procedure for γ given B_0 , M , and F^* .

following distribution is very effective,

$$\gamma \sim \zeta_{min} \delta(\gamma_{min}) + t(\mu, \sigma, \nu) \mathbf{1}_{\gamma > \gamma_{min}}. \quad (3.34)$$

1408 Above, t is the density of the three pa-
 1409 rameter location-scale family Student's t dis-
 1410 tribution with location μ , scale σ , and de-
 1411 grees of freedom ν . $\mathbf{1}_{\gamma > \gamma_{min}}$ is an indica-
 1412 tor function that serves to truncate Stu-
 1413 dent's t distribution at the lower bound γ_{min} .
 1414 $\delta(\gamma_{min})$ is the Dirac delta function evaluated
 1415 at γ_{min} , which is scaled by the known value
 1416 ζ_{min} ; this places probability mass ζ_{min} at
 1417 the point γ_{min} . Since sampling from Stu-
 1418 dent's t distribution is readily doable, sam-
 1419 pling from a truncated Student's t mixture
 1420 only requires slight modification.

Let T be the CDF of the modeled distri-
 bution of γ . Since the point $(\gamma_{min}, \zeta_{min})$ is
 known from the dynamics of the Schnute model at a given RP, full specification of Eq. (3.34)
 only requires determining the values for μ , σ , and ν which make T best approximate $\zeta(\gamma)$.
 Thus, the values of μ , σ , and ν are chosen by minimizing the L^2 distance between $T(\gamma)$ and
 $\zeta(\gamma)$.




Figure 3.18: $\zeta(\gamma)$ Plotted for $F^* = 0.1$ and $M = 0.2$. The point $(\gamma_{min}, \zeta_{min})$ shows the lowest biologically meaningful value of γ ; below which productivity is negative.

$$[\hat{\mu}, \hat{\sigma}, \hat{\nu}] = \arg \min_{[\mu, \sigma, \nu]} \int_{\Gamma} (T(\gamma; \mu, \sigma, \nu) - \zeta(\gamma))^2 d\gamma \quad (3.35)$$

1421 Fitting the distribution $T(\gamma|\hat{\mu}, \hat{\sigma}, \hat{\nu})$ for
 1422 use generating γ^* values at a specific F^* and
 1423 M releases the need to invert ζ . $T(\gamma|\hat{\mu}, \hat{\sigma}, \hat{\nu})$,
 1424 together with the structure in Eq. (3.33),
 1425 allows for the collection of an approximate
 1426 LHS sample via the algorithm seen in Algo-
 1427 rithm (2).

1428 $\frac{F^*}{M}$ is drawn uniformly from \mathcal{F}_i . Con-
 1429 ditioning on the sample of F^* , and M ,
 1430 $T(\gamma|\hat{\mu}, \hat{\sigma}, \hat{\nu})$ is fit and γ^* is sampled. ζ^* is
 1431 then computed and placed into the appropri-
 1432 ate grid element \mathcal{B}_j . Given γ^* , the cascade
 1433 $\alpha(\gamma^*)$, and $\beta(\alpha(\gamma^*), \gamma^*)$, can be computed.
 1434 The algorithm continues until all of the de-
 1435 sign elements, $(\frac{F^*}{M}, \zeta^*) \Leftrightarrow (\alpha^*, \beta^*, \gamma^*)$, have
 1436 been computed for all $i \in [1, \dots, n]$.

Algorithm 2 LHS of size n on rectangle R .

```

1: procedure  $LHS_n(R)$ 
2:   Define  $n$ -grids  $\mathcal{F}, \mathcal{B} \in R$ 
3:   for each grid element  $i$  do
4:     Draw  $\frac{F^*}{M} \sim Unif(\mathcal{F}_i)$ 
5:     Compute  $[\hat{\mu}, \hat{\sigma}, \hat{\nu}]$  given  $F^*$  &  $M$ 
6:     while  $\mathcal{B}_j$  not sampled do
7:       Draw  $\gamma^* \sim T(\gamma|\hat{\mu}, \hat{\sigma}, \hat{\nu})$ 
8:       Compute  $\zeta^* = \zeta(\gamma^*)$ 
9:       Compute  $j$  such that  $\zeta^* \in \mathcal{B}_j$ 
10:    end while
11:    Compute  $\alpha^* = \alpha(\gamma^*, F^*, M)$ 
12:    Compute  $\beta^* = \beta(\alpha^*, \gamma^*, M, B_0)$ 
13:    Save  $(\frac{F^*}{M}, \zeta^*) \Leftrightarrow (\alpha^*, \beta^*, \gamma^*)$  in  $\mathcal{F}_i \times \mathcal{B}_j$ 
14:  end for
15: end procedure

```

1437 Design Refinement

1438 Since the behavior of RP inference, under misspecified models, will vary in yet-unknown
 1439 ways, the exact sampling design density may be hard to know a priori. Several factors,
 1440 including the particular level of observation uncertainty, high variance (i.e. hard to resolve)
 1441 features of the response surface, or simply "gappy" instantiations of the initial LHS design
 1442 may necessitate adaptive design refinement, to accurately describe RP biases. Given the
 1443 temperamental relationship between RPs and productivity parameters in the Schnute model,
 1444 a recursive refinement algorithm, that makes use of the previously described LHS routine, is
 1445 developed.

1446 While LHS ensures uniformity in the design margins, and a certain degree of spread, it
 1447 is widely recognized that particular LHS instantiations may leave substantive gaps in the
 1448 simulation design. To correct this, LHS is often paired with design elements of maximin

1449 design (Morris & Mitchell, 1995; Devon Lin & Tang, 2015). Maximin designs sample the
 1450 design space by maximizing the minimum distance between sampled points. This has the
 1451 advantage of definitionally filling holes in the design, however because no points are ever
 1452 drawn outside of the design domain, samples tend to clump around edges (particularly
 1453 corners) of the design domain. Since LHS ensures uniformity in the margins and maximin
 1454 designs enjoys a certain sense of optimality in how they define and fill gaps (Johnson et al.,
 1455 1990), the methods are quite complimentary when combined.

Making use of this complimentary relationship, holes in the existing LHS design of RPs
 are identified based on maximin design principles. New design points are collected based on
 areas of the RP design space which maximizes the minimum distance between all pairs of
 points in the current design, based on the following distance function

$$d(\mathbf{x}, \mathbf{x}') = \sqrt{(\mathbf{x} - \mathbf{x}')^T \mathbf{D}^{-1} (\mathbf{x} - \mathbf{x}')} \quad (3.36)$$

$$\mathbf{D} = \text{diag} \left[(\max(\mathcal{F}) - \min(\mathcal{F}))^2, (\max(\mathcal{B}) - \min(\mathcal{B}))^2 \right].$$

1456 Above, d is a scaled distance function that defines the distance between points in the
 1457 differing scales of $\frac{B^*}{B_0}$ and $\frac{F^*}{M}$. \mathbf{D} is a diagonal matrix that measures the squared size of the
 1458 domain in each axis of so as to normalize distances to a common scale.

If \mathbf{X}_n is the initial design, computed on R_{full} , let \mathbf{x}_a be the augmenting point which
 maximizes the minimum distance between all of the existing design points,

$$\mathbf{x}_a = \underset{\mathbf{x}'}{\text{argmax}} \min \{d(\mathbf{x}_i, \mathbf{x}') : i = 1, \dots, n\}. \quad (3.37)$$

1459 The point \mathbf{x}_a is used as an anchor for augmenting \mathbf{X}_n . An additional $LHS_{n'}$ (via
 1460 Algorithm (2)) is collected, adding n' design points, centered around \mathbf{x}_a , to the overall
 1461 design. The augmenting region, $R_{(x_a, d_a)}$, for collecting $LHS_{n'}$ is defined based on the square
 1462 centered at \mathbf{x}_a with side length $2d_a$, where $d_a = \min \{d(\mathbf{x}_i, \mathbf{x}_a) : i = 1, \dots, n\}$, in the space
 1463 defined by the metric d .

1464 Due to the tendency of maximin sampling to cluster augmenting points on the edges of
 1465 the design space, $R_{(x_a, d_a)}$ is truncated by the outer most limits of R_{full} so as to focus design

1466 augmentation within the specified domain of the simulation. Furthermore, since the design
 1467 space has a nonlinear constraint at low values of $\frac{B^*}{B_0}$, the calculation of x_a is further truncated
 1468 based on a convex hull defined by the existing samples in the overall design.

1469 Design refinement then proceeds as follows. An initial design is computed, $X_n = LHS_n(R_{full})$,
 1470 based on an overall simulated region of RPs R_{full} . The maximin augmenting point, x_a , is
 1471 computed at a maximin distance of d_a from the existing samples. An augmenting design
 1472 $X_{n'} = LHS_{n'}(R_{(x_a, d_a)})$ is collected and added to X_n . Design refinement carries on recursively
 1473 collecting augmenting designs in this way until the maximin distance falls below the desired
 1474 level.

1475 5 .4 Gaussian Process Metamodel

1476 At its core, a metamodel is simply a model of some mapping of inputs to outputs (the
 1477 mapping itself is typically defined by a computer model). By modeling the mapping with a
 1478 statistical model (that explicitly defines the relevant features of the mapping) a metamodel
 1479 defines a specific ontology for the mapping. By simulating examples of the mapping, the
 1480 inferential infrastructure of the statistical model is used to empirically learn an effective
 1481 emulation of the mapping within the ontology defined by the statistical model. The pre-
 1482 dictive infrastructure of the statistical model is then useful as an approximate abstraction
 1483 of the system itself to better understand the system through further data collection, cheap
 1484 approximation of the mapping, and/or study of the mapping itself.

1485 In this setting, the aim of metamodeling is to study how well RPs are inferred when typical
 1486 two parameter models of productivity (Logistic and BH) are misspecified for populations
 1487 that are actually driven by more complicated dynamics. The simulation design, \mathbf{X} , provides
 1488 a sample of different population dynamics that are driven by three parameter production
 1489 functions broadly in RP space. By simulating index of abundance data from the three
 1490 parameter model, and fitting those data with the two parameter production model, we
 1491 observe particular instances of how well RPs are inferred at the given misspecification of the
 1492 two parameter model relative to the true three parameter production model. By gathering
 1493 all of the simulated instances of how RPs are inferred (under the two parameter model),
 1494 we form a set of example mappings to train a metamodel which represents the mapping

1495 of true RPs (under the three parameter model) to estimates of RPs under the misspecified
 1496 two parameter production model. The metamodel is essentially a surrogate for inference
 1497 under the misspecified two parameter production model that controls for the specific degree
 1498 of model misspecification.

1499 A flexible GP model is assumed for the structure of the metamodel to describe the map-
 1500 ping of RPs under misspecified two parameter models of productivity. A GP is a stochastic
 1501 process generalizing the multivariate normal distribution to an infinite dimensional analog.
 1502 GP models are often specified primarily through the choice of a covariance (or correlation)
 1503 function which defines the relationship between locations in the input space. Typically corre-
 1504 lation functions are specified so that points closely related in space result in correlated effects
 1505 in the model. In this setting the inputs to the GP metamodel are the space of reference points
 1506 with define the simulated three parameter production models.

While index of abundance data are generated from three parameter models, at each
 design location of the simulation, fitting the restricted two parameter model results in a
 maximum likelihood estimate (MLE; and associated estimation uncertainty) of each of the
 productivity parameters (i.e. Schaefer:[$\log(r)$, $\log(K)$], BH:[$\log(\alpha)$, $\log(\beta)$]). To simplify
 the specification of the metamodel, let \mathbf{y} be a vector collecting the fitted MLEs for one of
 the productivity parameters, and let $\boldsymbol{\omega}$ be a vector of estimates of the estimator variances
 (via the inverted Fisher information) at each \mathbf{y} . Each of the fitted productivity parameter
 estimates are then modeled using independent instances of the following GP metamodel.

$$\begin{aligned}\mathbf{y} &= \beta_0 + \mathbf{X}\boldsymbol{\beta} + \mathbf{v} + \boldsymbol{\epsilon} \\ \mathbf{v} &\sim N_n(\mathbf{0}, \tau^2 \mathbf{R}_\ell) \\ \boldsymbol{\epsilon} &\sim N_n(\mathbf{0}, \boldsymbol{\omega}'\mathbf{I})\end{aligned}\tag{3.38}$$

1507 \mathbf{X} is the $n \times 2$ LHS design matrix of RPs for each simulated three parameter data
 1508 generating model as described in Section (5.3). ϵ models independent normally distributed
 1509 error, which provides an ideal mechanism for propagating uncertainty from inference in the
 1510 simulation step into the metamodel. By matching each \mathbf{y}_i with an observed ω_i variance term,
 1511 ϵ serves to down weight the influence of each \mathbf{y}_i in proportion to the inferred production model

1512 sampling distribution uncertainty. This has the effect of smoothing the GP model in a way
 1513 similar to the nugget effect (Gramacy & Lee, 2012), although the application here models
 1514 this effect heterogeneously.

The term, \mathbf{v} , contains spatially correlated GP effects. The correlation matrix, \mathbf{R}_ℓ describes how RPs close together in the simulation design are more correlated than those that are far away. This spatial effect is modeled with a squared exponential correlation function,

$$R(\mathbf{x}, \tilde{\mathbf{x}}) = \exp \left(\sum_{i=1}^2 \frac{-(x_i - \tilde{x}_i)^2}{2\ell_j^2} \right). \quad (3.39)$$

1515 R has an anisotropic separable form which allows for differing length scales, ℓ_1 and ℓ_2 ,
 1516 in the different RP axes. The flexibility to model correlations separately in the different
 1517 RP axes is key due to the differences in the extent of the RP domains marginally. The
 1518 metamodel parameters β_0 , $\boldsymbol{\beta}$, τ^2 , ℓ_1 and ℓ_2 are fit via MLE against the observations \mathbf{y} , \mathbf{X} ,
 1519 and $\boldsymbol{\omega}$ from simulation fits.

1520 Fitting the metamodel allows for a full predictive description of inference under the
 1521 misspecified restricted models. Predictive estimates are obtained via kriging (Cressie, 2015)

$$\hat{y}(\mathbf{x}) = \beta_0 + \mathbf{x}\boldsymbol{\beta} + \mathbf{r}(\mathbf{x})'\mathbf{R}_\ell^{-1}(\mathbf{y} - (\beta_0 + \mathbf{X}\boldsymbol{\beta})) \quad (3.40)$$

1522 $\hat{y}(\mathbf{x})$ is the predicted value of the modeled productivity parameter MLE under the two
 1523 parameter production model, when the index of abundance is generated from the three
 1524 parameter production model at RP location \mathbf{x} . $\mathbf{r}(\mathbf{x})$ is a vector-valued function of correlation
 1525 function evaluations for the predictive location \mathbf{x} against all observations in \mathbf{X} (i.e. $\mathbf{r}(\mathbf{x}) =$
 1526 $\mathbf{R}(\mathbf{x}, \mathbf{x}_i) \forall \mathbf{x}_i \in \mathbf{X}$).

1527 While metamodeling occurs on the inferred productivity parameters of the restricted
 1528 production model, the metamodel can also be used to build estimates of major biological
 1529 RPs. For the BH model the relevant transformations for relating productivity parameters
 1530 with RPs are given in Eqs. (3.29, 3.32) with γ fixed to -1; for the Schaefer model $\hat{B}^* = \frac{\hat{K}}{2}$ and
 1531 $\hat{F}^* = \frac{\hat{r}}{2}$. Applying the metamodel predictive surfaces on the scale of RP estimates allows for
 1532 the quantification of estimation bias that is induced by fitting a misspecified two parameter

production model to indices of abundance generated under three parameter productivity.

5.5 Catch

It is known that contrast in the observed index and catch time series can effect inference on the productivity parameters (Hilborn & Walters, 1992). In this setting contrast refers to changes in the long term trends of index data. Figure (3.19, *right*) demonstrates an example of biomass that includes contrast induced by catch. It is not well understood how contrast may factor into inferential failure induced by model misspecification. Thus catch is parameterized so as to allow for a spectrum of possible contrast simulation settings.

Catch is parameterized so that $F(t)$ can be controlled with respect to F^* . Recall that catch is assumed to be proportional to biomass, so that $C(t) = F(t)B(t)$. To control $F(t)$ with respect to F^* , $C(t)$ is specified by defining the quantity $\frac{F(t)}{F^*}$ as the relative fishing rate. $B(t)$ is defined by the solution of the ODE, and F^* is defined by the biological parameters of the model. By defining $\frac{F(t)}{F^*}$, catch can then be written as $C(t) = F^* \left(\frac{F(t)}{F^*} \right) B(t)$.

Intuitively $\frac{F(t)}{F^*}$ describes the fraction of F^* that $F(t)$ is specified to for the current $B(t)$. When $\frac{F(t)}{F^*} = 1$, $F(t)$ will be held at F^* , and the solution of the ODE brings $B(t)$ into equilibrium at B^* . When $\frac{F(t)}{F^*}$ is held constant in time biomass comes to equilibrium as an exponential decay from K approaching B^* . When $\frac{F(t)}{F^*} < 1$, $F(t)$ is lower than F^* and $B(t)$ is pushed toward $\bar{B} > B^*$. Contrarily, when $\frac{F(t)}{F^*} > 1$, $F(t)$ is higher than F^* and $B(t)$ is pushed toward $\bar{B} < B^*$; the precise values of \bar{B} can be calculated from the steady state biomass equations provided above and depend upon the specific form of the production function.

For the simulations presented here, a family of fishing behaviors are considered where the fishing rate accelerates as technology and fishing techniques improve rapidly until management practices are applied, which ultimately brings fishing into equilibrium at F^* . This is parameterized as three distinct phases, over a total of 45 units of time, with each phase lasting 15 time units. The specific form is given below.

$$\frac{F(t)}{F^*} = ae^{bt}\mathbf{1}_{0 \leq t < 15} + (d - ct)\mathbf{1}_{15 \leq t < 30} + \mathbf{1}_{30 \leq t \leq 45} \quad (3.41)$$

The first term of Eq(3.41) is an exponential increase in fishing, the second term is a linear

decline in relative fishing as initial management practices are applied, and the third term, $\mathbf{1}_{30 \leq t \leq 45}$, simply holds the fishing rate at F^* there after. These three phases are controlled by the four parameters a , b , c , and d . By enforcing that the interface of the phases meet at χ_{max} and 1 respectively the relative fishing series is reduced to a two parameter family.

$$a = e^{\log(\chi_{max}) - 15b} \quad b = \frac{1}{t - 15} \log \left(\frac{\chi_{min}}{\chi_{max}} \right) \quad (3.42)$$

$$c = \frac{\chi_{max} - 1}{15 - 1} \quad d = 15c + \chi_{max} \quad (3.43)$$

1553 By further specifying $\chi_{max} = 1.6^\chi$ and $\chi_{min} = 0.4^\chi$ the two parameters χ_{max} , and χ_{min}
 1554 can be reduced to the single parameter χ . The tuning parameter χ then singularly controls
 1555 contrast that appears in time series data.




Figure 3.19: (left) Relative fishing with low, medium, and high contrast. (right) Population biomass and catch at each associated level of contrast.

1556 When $\chi = 0$, the relative fishing rate is a constant at 1 to create a low contrast simulation
 1557 environment. As χ increases Eq (3.41) induces more and more contrast in the observed index
 1558 and catch time series until $\chi = 1$ which produces a high contrast simulation environment.
 1559 Figure (3.19) demonstrates a spectrum of contrast simulation environments as well as the
 1560 time series data they induce in the solution of the production model ODE.

5.6 Two Parameter Production Model Inference

The simulated mapping results from fitting an intentionally misspecified two parameter production model to index of abundance data that are generated from a more complex three parameter model of productivity. Thus, let I_t be an index of abundance simulated from the three parameter PT or Schnute production models at time $t \in \{1, 2, 3, \dots, T\}$. However the fitted model is specified to be intentionally misspecified so that the fitted model is driven by a two parameter Schaefer, or BH production model respectively.

The observation model for the fitted model is log-normal such that,

$$I_t|q, \sigma^2, \boldsymbol{\theta} \sim LN(qB_t(\boldsymbol{\theta}), \sigma^2). \quad (3.44)$$

$B_t(\boldsymbol{\theta})$ is defined by the solution of the ODEs defined by the Schaefer, or BH models. For the Schaefer model $\boldsymbol{\theta} = [r, K]$, and for the BH model $\boldsymbol{\theta} = [\alpha, \beta]$. From the perspective of the fitted model, the observed I_t are assumed independent conditional on q , σ^2 , r , K and the two parameter ODE model for biomass. Thus the log likelihood can be written as

$$\log \mathcal{L}(q, \sigma^2, \boldsymbol{\theta}; I) = -\frac{T}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_t \log \left(\frac{I_t}{qB_t(\boldsymbol{\theta})} \right)^2. \quad (3.45)$$

In this setting, q is fixed at the true value of 0.0005 to focus on the inferential effects of model misspecification on biological parameters. σ^2 and $\boldsymbol{\theta}$ are reparameterized to the log scale and fit via MLE. Reparameterizing the parameters to the log scale improves the reliability of optimization, in addition to facilitating the use of Hessian information for estimating MLE standard errors.

Given that the biological parameters enter the likelihood via a nonlinear ODE, and further the parameters themselves are related to each other nonlinearly, the likelihood function can often be difficult to optimize. A hybrid optimization scheme is used to maximize the log likelihood to ensure that a global MLE solution is found. The R package GA ([Scrucca, 2013, 2017](#)) is used to run a genetic algorithm to explore parameter space globally. Optimization periodically jumps into the L-BFGS-B local optimizer to refine optima within a local mode. The scheme functions by searching globally, with the genetic algorithm, across many initial

values for starting the local gradient-based optimizer. The genetic algorithm serves to iteratively improve hot starts for the local gradient-based optimizer. Additionally, optimization is only considered to be converged when the optimum results in an invertible Hessian at the found MLE.

5.7 Continuous model formulation

An important (and often overlooked) implementation detail is the solution to the ODE which defines the progression of biomass through time. As a statistical model it is of paramount importance that this ODE not only have a solution, but also that the solution be unique.

If the form of $\frac{dB}{dt}$ is at least Lipschitz continuous, then the Cauchy-Lipschitz-Picard theorem provides local existence and uniqueness of $B(t)$. Recall from Eq(3.14) that $\frac{dB}{dt}$ is separated into a term for biomass production, $P(B)$, and a term for removals, $Z(t)B(t)$. For determining Lipschitz continuity of $\frac{dB}{dt}$, the smallest Lipschitz constant of $\frac{dB}{dt}$ will be the sum of the constants for each of the terms $P(B)$ and $Z(t)B(t)$ separately. Typically any choice of $P(B)$ will be continuously differentiable, which implies Lipschitz continuity. At a minimum $Z(t)$ typically contains fishing mortality as a function of time $F(t)$ to model catch in time as $C(t) = F(t)B(t)$. $Z(t)$ may or may not contain M , but typically M is modeled as stationary in time and does not pose a continuity issue, unlike some potential assumptions for $C(t)$.

In practice $C(t)$ is determined by a series of observed, assumed known, catches. Catch observations are typically observed on a quarterly basis, but in practice may not be complete for every quarter of the modeled period. It is overwhelmingly common to discretized the ODE via Euler's method with integration step sizes to match the observation frequency of the modeled data. This is often convenient but can present several issues. This strategy often pushes the assumption of catch continuity under the rug, but for regularity of the statistical model an implicit assumption of continuity of the catches is required. While mechanistically at the finest scale fishers must only catch discrete packets of biomass (i.e. individual fish), it is sensible to consider catches as accruing in a continuous way. Furthermore any assumption of continuity will be required to be at least Lipschitz continuous for the required regularity of the model.

Here I assume catches accrue linearly between observed catches. This assumption defines

1609 the catch function as a piecewise linear function of time, with the smallest Lipschitz constant
1610 for the catch term defined by the steepest segment of the catch function. This assumption
1611 represents one of the simplest ways of handling catch, while retaining Lipschitz continuity
1612 overall. Furthermore linearly interpolated catch is adequately parsimonious for the typical
1613 handling of catches.

1614 **Integration and Stiffness**

1615 As previously mentioned, the overwhelming majority of implementations of population dy-
1616 namics models discretized the ODE using Euler's method with the integration step sized
1617 fixed so as to match the observation frequency. In this setting we explore model parameter-
1618 izations that explore the full extent of biologically relevant reference points. This exercise
1619 produces some combinations of parameters that result in numerically stiff ODEs.

1620 The concept of stiffness in ODEs is hard to precisely characterize. [Wanner and Hairer](#)
1621 [\(1996, p.2\)](#) describe stiffness in the following pragmatic sense, "Stiff equations are problems
1622 for which explicit methods don't work". It is hard to make this definition more mathemati-
1623 cally precise, but this a consistent issue for models of fast growing species in the low contrast
1624 simulation. Euler's method, as often implemented, is particularly poorly suited for these
1625 stiff regions of parameter space. In these stiff regions it is necessary to integrate the ODE
1626 with an implicate integration method.

1627 Several of the most common implicate methods were tried including the Livermore Solver
1628 for ODEs (lsode), and the Variable Coefficient ODE Solver (vode) as implemented in the
1629 deSolve package of R ([Soetaert et al., 2010](#)). The difference between implicit solvers is
1630 negligible, while explicit methods result in wildly varying solutions to the ODE in stiff
1631 regions of parameter space. Results shown here are computed using the lsode integration
1632 since it runs relatively quickly and has a relatively smaller footprint in system memory.

6 Results

6.1 PT/Schaefer

An MSY -Optimal Catch History

When $F(t)$ is held constant at F^* , as it is in the "low contrast" simulation setting, $B(t)$ comes to equilibrium as an exponential decay from K to B^* . Understanding model misspecification bias is simplified in this setting due to the relative simplicity that this induces in $B(t)$. However this simplicity is known to poorly inform estimates of r , and thus F^* , due to the limited range of the production function that is observed (Hilborn & Walters, 1992).

Figure (3.20) shows four of the most misspecified example production function fits as compared to the true data generating PT production functions. The rug plots below each set of curves show how the observed biomasses decay exponentially from K to B^* in each case. In particular, notice how observations only exist where the PT biomass is greater than B^* . Due to the leaning of the true PT curves, and the symmetry of the logistic parabola, the logistic curve only observes information about its slope at the origin from data observed on the right portion of the PT curves. The top two panels of Figure (3.20) shows PT data generated such that $\frac{B^*}{B(0)} > 0.5$; in these cases PT is steeper to the right of B^* than it is on the left, and so the the logistic curve over-estimates r , and consequently also over-estimates F^* . The bottom two panels of Figure (3.20) show PT




Figure 3.20: A comparison of the true PT production function (in black) and the estimated logistic curve (in red) with 95% CI shown. The examples shown represent the four corners of maximum model misspecification in the simulated RP-space. Observed biomasses are plotted in the rug plots below the curves.

data generated with $\frac{B^*}{B(0)} < 0.5$ and where the vice versa phenomena occurs. PT is shallower

to the right of B^* than it is on the left and so the logistic parabola estimate tends to under estimate F^* .

Metamodeled Trends

Each point in the space of the RPs F^* and $\frac{B^*}{B(0)}$ uniquely identifies a complete PT model with different combinations of parameters values. Recall that when $\gamma = 2$ for the PT model, the PT curve becomes a parabola and is equivalent to the logistic curve of the Schaefer model. Since the logistic curve is symmetric about B^* , the Schaefer model must fix the value of $\frac{B^*}{B(0)}$ at the constant 0.5 for any value of F^* . So the line through RP space defined by $\frac{B^*}{B(0)} = 0.5 \quad \forall \quad F^*$, defines the subset of RP space where $\gamma = 2$ and where the PT model is equivalent to the Schaefer model. For brevity this subset of RP where $\frac{B^*}{B(0)} = 0.5$ will be referred to as the ‘‘Schaefer set’’. Thus simulated data that are generated along the Schaefer set will be the only data that are not misspecified relative to the Schaefer model; as PT data are simulated farther and farther away from this line at $\frac{B^*}{B(0)} = 0.5$ model misspecification of the Schaefer model becomes worse and worse.

While Figure (3.20) demonstrates a real trend in simulation results, individual simulation runs will at best show jittery trends due to the stochastic nature of statistical inference. The GP process metamodel accounts for this stochasticity to focus analysis on the signal in the simulation results. Recall that metamodeling occurs on the scale of the inferred productivity parameters of the restricted production model, by transforming metamodel predictions via Eq. (3.23), metamodeled predictions are obtained for Schaefer RPs. By further subtracting the true data generating PT RPs from the predicted Schaefer RPs at each point in RP space a pattern of inferential RP bias, induced by model misspecification of the Schaefer model, can be seen to be seen.

Figure (3.21) shows the pattern of biases the Schaefer model creates when fit to PT data generated at each point of RP space. An equivalent way to think of Figure (3.21) is that since the Schaefer model must estimate RPs in the Schaefer set, the metamodel arrows indicate the mapping that is created by inferring RPs under a misspecified Schaefer model fit to PT data generated at each point over the pictured region.

Since $\frac{B^*}{B_0}$ must be 0.5 under the Schaefer model, biases in the $\frac{B^*}{B_0}$ direction must simply




Figure 3.21: Joint bias direction for $(F^*, \frac{B^*}{B_0})$ estimates under the misspecified Schaefer Model. The intensity of color represents the excess bias relative to the shortest possible mapping. Results in the low contrast setting are shown *left*, and the high contrast setting is shown *right*.

map vertically onto the Schaefer set. Due to this simplified RP geometry under the Schaefer model, the degree of bias in $\frac{B^*}{B_0}$ estimation is entirely defined solely by the degree of model misspecification irrespective of F^* . Furthermore, the closest possible point along the Schaefer set that Schaefer model inference could map RPs would be the perfectly vertical mapping. This pattern only contains the strictly necessary bias present in $\frac{B^*}{B_0}$, and zero bias in F^* . Any deviation from this minimal bias pattern necessarily to be due to added bias in F^* .

The two simulation settings shown in Figure (3.21) are identical except for the amount of contrast present in the simulated index. The left panel of Figure (3.21) shows RP biases in the low contrast setting, while the right panel shows the high contrast setting. Notice that in the low contrast setting the RP bias pattern is far from the minimum distance mapping, however when contrast is added the mapping becomes much closer to a minimal bias mapping. In the low contrast setting the observed bias is consistent with the pattern and mechanism described in Figure (3.20), where F^* is underestimated for data generated below the Schaefer line and overestimated above the Schaefer set. In the high contrast simulation the mapping is nearly minimal distance with the exception of PT data generated with simultaneously low $\frac{B^*}{B_0}$ and high F^* .




Figure 3.22: Bias in F^* under the Schaefer model when PT data are generated with increasing contrast so that F^* and $\frac{B^*}{B_0}$ are fixed at 0.699 and 0.201 respectively.

1707 Figure (6 .1) demonstrates how bias in F^* estimation decreases as contrast is added to
 1708 PT data as generated in the low $\frac{B^*}{B_0}$ and high F^* regime. By including additional contrast
 1709 F^* bias is decreased, however parameterizing contrast so as to fully extinguish F^* bias may
 1710 require a more complex model of fishing.

- 1711 • summary of σ over RP space comparing between models (PT, Schnute, Schnute DD)
- 1712 to show areas of model breakdown.
- 1713 – miss-identifying signal for noise.
- 1714 – It happens more as the dynamics get more complex.
- 1715 – point to the full age structured models.
- 1716 • show the constrained BH space over a grid of $M, \kappa, \omega, W_\infty$
- 1717 • Show that the constrained spaces vary only slightly as compared with the consequences
- 1718 of misspecifying the functional form.
- 1719 • estimating these other quantities (while they can create quite different Biomass series)
- 1720 can only do so much to improve (expand) RP inference as compared with correctly
- 1721 modeling P .
- 1722 • mapping distance as a function of contrast at (3.5, 0.5)
- 1723 • for LHS grid locations show $\frac{B^*}{B_0}$ and F^* biases for grids in $M \in (0, 0.5)$ For sure in High
- 1724 Contrast, maybe also in Low??.

1725 7 Appendix: Inverting $\frac{B^*}{B(0)}$ and γ for the PT Model

For brevity let $\zeta = \frac{B^*}{B(0)}$.

$$\begin{aligned}\zeta &= \left(\frac{1}{\gamma}\right)^{\frac{1}{\gamma-1}} \\ \zeta &= \gamma \zeta^\gamma \\ \zeta &= \gamma e^{\gamma \log(\zeta)} \\ \zeta \log(\zeta) &= \gamma \log(\zeta) e^{\gamma \log(\zeta)}\end{aligned}$$

The Lambert product logarithm, W , is defined as the inverse function of $z = xe^x$ such that $x = W(z)$. Applying this definition allows for the isolation of γ .

$$\begin{aligned}\gamma \log(\zeta) &= W(\zeta \log(\zeta)) \\ \gamma &= \frac{W(\zeta \log(\zeta))}{\log(\zeta)}\end{aligned}\tag{3.46}$$

1726 The Lambert product logarithm is a multivalued function with a branch point at $-\frac{1}{e}$. The
1727 principal branch, $W_0(z)$, is defined on $z \in (-\frac{1}{e}, \infty)$, and the lower branch, $W_{-1}(z)$, is
1728 defined on $z \in (-\frac{1}{e}, 0)$. Taken individually, each respective branch is analytic, but cannot
1729 be expressed in terms of elementary functions.

1730 When $\zeta \in (0, \frac{1}{e})$ the solution of interest in Eq. (3.24) comes from W_0 . When $\zeta \rightarrow \frac{1}{e}$, the
1731 Fox Model emerges as $\gamma \rightarrow 1$. When $\zeta \in (\frac{1}{e}, 1)$ the solution of interest comes from W_{-1} . For
1732 the use case presented here, Eq. (3.24) is to be interpreted as,

$$\gamma = \begin{cases} \frac{W_0(\zeta \log(\zeta))}{\log(\zeta)} & \zeta \in (0, \frac{1}{e}) \\ \frac{W_{-1}(\zeta \log(\zeta))}{\log(\zeta)} & \zeta \in (\frac{1}{e}, 1) \end{cases}.\tag{3.47}$$

1733 Prager 2002, Figure(2).

1734 [https://math.stackexchange.com/questions/3004835/is-the-lambert-w-function-analytic-](https://math.stackexchange.com/questions/3004835/is-the-lambert-w-function-analytic-if-not-everywhere-then-on-what-set-is-it-an)
1735 [if-not-everywhere-then-on-what-set-is-it-an](https://math.stackexchange.com/questions/3004835/is-the-lambert-w-function-analytic-if-not-everywhere-then-on-what-set-is-it-an) [https://researchportal.bath.ac.uk/en/publications/algebraic-](https://researchportal.bath.ac.uk/en/publications/algebraic-properties-of-the-lambert-w-function-from-a-result-of-r)
1736 [properties-of-the-lambert-w-function-from-a-result-of-r](https://researchportal.bath.ac.uk/en/publications/algebraic-properties-of-the-lambert-w-function-from-a-result-of-r)

References

- Beverton, R. J., & Holt, S. J. (1957). *On the dynamics of exploited fish populations* (Vol. 11). Springer Science & Business Media.
- Brent, R. P. (1973). Chapter 4: An Algorithm with Guaranteed Convergence for Finding a Zero of a Function. In *Algorithms for minimization without derivatives*. Courier Corporation.
- Conn, P. B., Williams, E. H., & Shertzer, K. W. (2010). When can we reliably estimate the productivity of fish stocks? *Canadian Journal of Fisheries and Aquatic Sciences*, 67(3), 511–523.
- Cressie, N. (2015). *Statistics for spatial data*. John Wiley & Sons.
- Cushing, D. H. (1971, May). The Dependence of Recruitment on Parent Stock in Different Groups of Fishes. *ICES Journal of Marine Science*, 33(3), 340–362. Retrieved 2023-06-03, from <https://doi.org/10.1093/icesjms/33.3.340> doi: 10.1093/icesjms/33.3.340
- Deriso, R. B. (1980, February). Harvesting Strategies and Parameter Estimation for an Age-Structured Model. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(2), 268–282. Retrieved 2020-05-13, from <https://www.nrcresearchpress.com/doi/abs/10.1139/f80-034> doi: 10.1139/f80-034
- Devon Lin, C., & Tang, B. (2015). Latin Hypercubes and Space-filling Designs. In *Handbook of Design and Analysis of Experiments*.
- Fletcher, R. I. (1978). On the restructuring of the Pella-Tomlinson system. *Fish. Bull*, 76(3), 515–521.
- Fournier, D. A., & Doonan, I. J. (1987). A length-based stock assessment method utilizing a generalized delay-difference model. *Canadian Journal of Fisheries and Aquatic*

- Sciences*, 44(2), 422–437. (Publisher: NRC Research Press Ottawa, Canada)
- Fox Jr., W. W. (1970). An Exponential Surplus-Yield Model for Optimizing Exploited Fish Populations. *Transactions of the American Fisheries Society*, 99(1), 80–88. Retrieved 2022-02-17, from <https://onlinelibrary.wiley.com/doi/abs/10.1577/1548-8659%281970%2999%3C80%3AAESMFO%3E2.0.CO%3B2> (_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1577/1548-8659%281970%2999%3C80%3AAESMFO%3E2.0.CO%3B2>) doi: 10.1577/1548-8659(1970)99<80:AESMFO>2.0.CO;2
- Gramacy, R. B. (2020). *Surrogates: Gaussian process modeling, design, and optimization for the applied sciences*. Chapman and Hall/CRC.
- Gramacy, R. B., & Lee, H. K. (2012). Cases for the nugget in modeling computer experiments. *Statistics and Computing*, 22(3), 713–722. (Publisher: Springer)
- Hilborn, R. (2010). Pretty good yield and exploited fishes. *Marine Policy*, 34(1), 193–196. (Publisher: Elsevier)
- Hilborn, R., & Mangel, M. (1997). *The Ecological Detective: Confronting Models with Data*. Princeton University Press.
- Hilborn, R., & Walters, C. J. (1992). Quantitative Fisheries, Stock Assessment: Choice Dynamics, and Uncertainty Chapman and Hall. *New York*.
- Johnson, M. E., Moore, L. M., & Ylvisaker, D. (1990). Minimax and maximin distance designs. *Journal of statistical planning and inference*, 26(2), 131–148. (Publisher: Elsevier)
- Lee, H.-H., Maunder, M. N., Piner, K. R., & Methot, R. D. (2012, August). Can steepness of the stock–recruitment relationship be estimated in fishery stock assessment models? *Fisheries Research*, 125-126, 254–261. Retrieved 2022-01-29, from <https://linkinghub.elsevier.com/retrieve/pii/S0165783612001099> doi: 10.1016/j.fishres.2012.03.001
- Magnusson, A., & Hilborn, R. (2007). What makes fisheries data informative? *Fish and Fisheries*, 8(4), 337–358. (Publisher: Wiley Online Library)
- Mangel, M., MacCall, A. D., Brodziak, J., Dick, E., Forrest, R. E., Pourzand, R., & Ralston, S. (2013, April). A perspective on steepness, reference points, and stock assessment.

1792 *Canadian Journal of Fisheries and Aquatic Sciences*, 70(6), 930–940. Retrieved 2019-
 1793 07-03, from <https://www.nrcresearchpress.com/doi/10.1139/cjfas-2012-0372>
 1794 doi: 10.1139/cjfas-2012-0372

1795 McKay, M. D., Beckman, R. J., & Conover, W. J. (2000). A comparison of three methods
 1796 for selecting values of input variables in the analysis of output from a computer code.
 1797 *Technometrics*, 42(1), 55–61. (Publisher: Taylor & Francis)

1798 Morris, M. D., & Mitchell, T. J. (1995, February). Exploratory designs for computa-
 1799 tional experiments. *Journal of Statistical Planning and Inference*, 43(3), 381–402. Re-
 1800 trieved 2023-05-28, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S037837589400035T)
 1801 [037837589400035T](https://www.sciencedirect.com/science/article/pii/S037837589400035T) doi: 10.1016/0378-3758(94)00035-T

1802 Punt, A. E., Butterworth, D. S., Moor, C. L. d., Oliveira, J. A. A. D., & Haddon, M. (2016).
 1803 Management strategy evaluation: best practices. *Fish and Fisheries*, 17(2), 303–334.
 1804 Retrieved 2018-12-13, from [https://onlinelibrary.wiley.com/doi/abs/10.1111/](https://onlinelibrary.wiley.com/doi/abs/10.1111/faf.12104)
 1805 [faf.12104](https://onlinelibrary.wiley.com/doi/abs/10.1111/faf.12104) doi: 10.1111/faf.12104

1806 Punt, A. E., & Cope, J. M. (2019, September). Extending integrated stock assessment mod-
 1807 els to use non-depensatory three-parameter stock-recruitment relationships. *Fisheries*
 1808 *Research*, 217, 46–57. Retrieved 2019-07-19, from [http://www.sciencedirect.com/](http://www.sciencedirect.com/science/article/pii/S0165783617301819)
 1809 [science/article/pii/S0165783617301819](http://www.sciencedirect.com/science/article/pii/S0165783617301819) doi: 10.1016/j.fishres.2017.07.007

1810 Ramasubramanian, K., & Singh, A. (2017). *Machine learning using R* (No. 1). Springer.

1811 Rankin, P. S., & Lemos, R. T. (2015, October). An alternative surplus production
 1812 model. *Ecological Modelling*, 313, 109–126. Retrieved 2022-02-11, from [https://](https://www.sciencedirect.com/science/article/pii/S0304380015002732)
 1813 www.sciencedirect.com/science/article/pii/S0304380015002732 doi: 10.1016/
 1814 [j.ecolmodel.2015.06.024](https://www.sciencedirect.com/science/article/pii/S0304380015002732)

1815 Ricker, W. E. (1954). Stock and recruitment. *Journal of the Fisheries Board of Canada*,
 1816 11(5), 559–623. (Publisher: NRC Research Press Ottawa, Canada)

1817 Schnute, J. (1985, March). A General Theory for Analysis of Catch and Effort Data.
 1818 *Canadian Journal of Fisheries and Aquatic Sciences*, 42(3), 414–429. Retrieved 2020-
 1819 05-13, from <https://www.nrcresearchpress.com/doi/abs/10.1139/f85-057> doi:
 1820 10.1139/f85-057

1821 Schnute, J. (1987). A general fishery model for a size-structured fish population. *Canadian*

- Journal of Fisheries and Aquatic Sciences*, 44(5), 924–940. (Publisher: NRC Research Press Ottawa, Canada)
- Schnute, J. T., & Richards, L. J. (1998, February). Analytical models for fishery reference points. *Canadian Journal of Fisheries and Aquatic Sciences*, 55(2), 515–528. Retrieved 2020-01-14, from <https://www.nrcresearchpress.com/doi/abs/10.1139/f97-212> doi: 10.1139/f97-212
- Scrucca, L. (2013, April). GA: A Package for Genetic Algorithms in R. *Journal of Statistical Software*, 53, 1–37. Retrieved 2022-01-17, from <https://doi.org/10.18637/jss.v053.i04> doi: 10.18637/jss.v053.i04
- Scrucca, L. (2017). On Some Extensions to GA Package: Hybrid Optimisation, Parallelisation and Islands Evolution. On some extensions to GA package: hybrid optimisation, parallelisation and islands evolution. *The R Journal*, 9(1), 187–206. Retrieved 2022-01-17, from <https://journal.r-project.org/archive/2017/RJ-2017-008/index.html>
- Soetaert, K., Petzoldt, T., & Setzer, R. W. (2010, February). Solving Differential Equations in R: Package deSolve. *Journal of Statistical Software*, 33, 1–25. Retrieved 2023-05-29, from <https://doi.org/10.18637/jss.v033.i09> doi: 10.18637/jss.v033.i09
- Stein, M. (1987). Large sample properties of simulations using Latin hypercube sampling. *Technometrics*, 29(2), 143–151. (Publisher: Taylor & Francis)
- Von Bertalanffy, L. (1938). A quantitative theory of organic growth (inquiries on growth laws. II). *Human biology*, 10(2), 181–213. (Publisher: JSTOR)
- Walters, C. J. (2020). The continuous time Schnute-Deriso delay-difference model for age-structured population dynamics, with example application to the Peru anchoveta stock.
- Wanner, G., & Hairer, E. (1996). *Solving ordinary differential equations II* (Vol. 375). Springer Berlin Heidelberg.
- Yeakel, J. D., & Mangel, M. (2015, February). A generalized perturbation approach for exploring stock recruitment relationships. *Theoretical Ecology*, 8(1), 1–13. Retrieved 2023-06-03, from <https://doi.org/10.1007/s12080-014-0230-z> doi: 10.1007/s12080-014-0230-z