

Metamodeling for Bias Estimation of Biological Reference Points Under Two-Parameter SRRs

Nick Grunloh



13 August 2024



Outline

1 Introduction

2 The Schaefer Model

3 The Beverton-Holt Model

4 Delay Differential Growth Extension

5 End

Outline

- 1** Introduction
 - 2** The Schaefer Model
 - 3** The Beverton-Holt Model
 - 4** Delay Differential Growth Extension
 - 5** End

- MSA?
 - RPs
 - Steepness paper

Surplus Production Model General Structure

$$I_t = qB_t e^\epsilon \quad \epsilon \sim N(0, \sigma^2). \quad (1)$$

$$\frac{dB}{dt} = P(B(t); \theta) - Z(t)B(t). \quad (2)$$

Outline

1 Introduction

2 The Schaefer Model

3 The Beverton-Holt Model

4 Delay Differential Growth Extension

5 End


$$P_p(B; [r, K]) = rB \left(1 - \left(\frac{B}{K}\right)\right)$$

$$F^* = \frac{r}{2}$$


$$B^* = \frac{K}{2}$$

$$B_0 = K$$

$$\frac{B^*}{B_0} = \frac{1}{2}$$



$$P_p(B; [r, K, \gamma]) = \frac{rB}{\gamma - 1} \left(1 - \left(\frac{B}{K} \right)^{(\gamma-1)} \right). \quad (3)$$



$$\frac{dB}{dt} = \frac{rB}{\gamma - 1} \left(1 - \left(\frac{B}{K} \right)^{\gamma-1} \right) - FB. \quad (4)$$

$$\bar{B}(F) = K \left(1 - \frac{F(\gamma - 1)}{r} \right)^{\frac{1}{(\gamma-1)}}. \quad (5)$$

$$F^* = \frac{r}{\gamma} \qquad \qquad \frac{B^*}{\bar{B}(0)} = \left(\frac{1}{\gamma} \right)^{\frac{1}{\gamma-1}} \quad (6)$$

$$F^* = \frac{r}{\gamma} \quad \frac{B^*}{\bar{B}(0)} = \left(\frac{1}{\gamma} \right)^{\frac{1}{\gamma-1}} \quad (7)$$

$$r = \gamma F^* \quad \gamma = \frac{W \left(\frac{B^*}{\bar{B}(0)} \log \left(\frac{B^*}{\bar{B}(0)} \right) \right)}{\log \left(\frac{B^*}{\bar{B}(0)} \right)} \quad (8)$$








Figure: (left) Relative fishing with low, medium, and high contrast.
(right) Population biomass and catch at each associated level of contrast.







Figure: Joint bias direction for $(F^*, \frac{B^*}{B_0})$ estimates under the misspecified Schaefer Model. The intensity of color represents the excess bias relative to the shortest possible mapping. Results in the low contrast setting are shown *right*, and the high contrast setting is shown *left*.



Outline

1 Introduction

2 The Schaefer Model



3 The Beverton-Holt Model

4 Delay Differential Growth Extension

5 End


$$P(B) = \frac{\alpha B}{1 + \beta B}. \quad (9)$$

$$\frac{dB}{dt} = P(B; \theta) - MB - FB \quad (10)$$



$$R(B; \alpha, \beta, \gamma) = \alpha B (1 - \beta \gamma B)^{\frac{1}{\gamma}}$$

Schnute Production



$$\bar{B}(F) = \frac{1}{\gamma\beta} \left(1 - \left(\frac{M+F}{\alpha} \right)^\gamma \right). \quad (11)$$

$$B_0 = \frac{1}{\gamma\beta} \left(1 - \left(\frac{M}{\alpha} \right)^\gamma \right) \quad (12)$$


$$\frac{B^*}{B_0} = \frac{1 - \left(\frac{M+F^*}{\alpha}\right)^\gamma}{1 - \left(\frac{M}{\alpha}\right)^\gamma}. \quad (13)$$

$$\frac{d\bar{Y}}{dF} = \bar{B}(F) + F \frac{d\bar{B}}{dF} \quad (14)$$

$$\frac{d\bar{B}}{dF} = -\frac{1}{\beta} \left(\frac{\left(\frac{M+F}{\alpha}\right)^\gamma}{F+M} \right). \quad (15)$$

$$\begin{aligned}\alpha &= (M + F^*) \left(1 + \frac{\gamma F^*}{M + F^*}\right)^{1/\gamma} \\ \beta &= \frac{1}{\gamma B_0} \left(1 - \left(\frac{M}{\alpha}\right)^\gamma\right) \\ \frac{B^*}{B_0} &= \frac{1 - \left(\frac{M+F^*}{\alpha}\right)^\gamma}{1 - \left(\frac{M}{\alpha}\right)^\gamma}. \end{aligned} \tag{17}$$

$$\gamma' \sim \zeta_{min}\delta(\gamma_{min}) + t(\mu, \sigma, \nu)\mathbf{1}_{\gamma > \gamma_{min}} \quad (18)$$



Schnute Recruitment

$$R(B; \alpha, \beta, \gamma) = \alpha B_{t-a_s} (1 - \beta \gamma B_{t-a_s})^{\frac{1}{\gamma}}$$

Logistic


$$\gamma = 1$$

Ricker

$$\gamma \rightarrow 0$$

Beverton-Holt

$$\gamma = -1$$






Relative Bias in Estimated B_{MSY}/B_0 High Contrast
Bias Direction for $(F_{MSY}/M, B_{MSY}/B_0)$ Jointly

Figure: Heatplots showing the bias in RP estimation induced by model misspecification of the BH model in the high contrast simulation setting. In all cases



Add Yield curves from AFS presentation.

Outline

1 Introduction

2 The Schaefer Model

3 The Beverton-Holt Model

4 Delay Differential Growth Extension

5 End

General Modeling Structure

$$\begin{aligned}\frac{dB}{dt} &= \overbrace{w(a_s)R(B; \theta)}^{\text{Recruitment Biomass}} + \overbrace{\kappa [w_\infty N - B]}^{\text{Net Growth}} - \overbrace{(M + F)B}^{\text{Mortality}} \\ \frac{dN}{dt} &= R(B; \theta) - (M + F)N\end{aligned}$$

$R(B_{t-a_s}; \alpha, \beta, \gamma)$: Schnute Three Parameter Recruitment

$w(a)$: Von Bertalanffy Individual Growth

Knife-Edge Selectivity at age a_s

Schnute Recruitment

$$R(B; \alpha, \beta, \gamma) = \alpha B_{t-a_s} (1 - \beta \gamma B_{t-a_s})^{\frac{1}{\gamma}}$$

Logistic


$$\gamma = 1$$

Ricker

$$\gamma \rightarrow 0$$

Beverton-Holt

$$\gamma = -1$$



Schnute Recruitment

$$R(B; \alpha, \beta, \gamma) = \alpha B_{t-a_s} (1 - \beta \gamma B_{t-a_s})^{\frac{1}{\gamma}}$$

Logistic


$$\gamma = 1$$

Ricker

$$\gamma \rightarrow 0$$

Beverton-Holt

$$\gamma = -1$$




Von Bertalanffy Individual Growth

$$w(a) = w_\infty(1 - e^{-\kappa(a-a_0)})$$

- Instant Growth:
(Production Model)
 $a_s \rightarrow 0$ $\kappa \rightarrow \infty$

- Dynamic Growth:
 a_s : Not Instant
 κ : Peakness of Dynamics






$$\bar{B}(F) = \frac{1}{\beta\gamma} \left(1 - \left(\frac{(F+M)(F+M+\kappa)}{\alpha w(a_s)(F+M + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma \right) \quad (19)$$

$$\bar{N}(F) = \frac{\alpha \bar{B}(F)(1 - \beta\gamma \bar{B}(F))^{1/\gamma}}{F+M} \quad (20)$$



$$\alpha = \left[\left(\frac{Z^*(Z^* + \kappa)}{w(a_s)(Z^* + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma + \left(\frac{\gamma F^*}{w(a_s)} \right) \left(\frac{Z^*(Z^* + \kappa)}{w(a_s)(Z^* + \frac{\kappa w_\infty}{w(a_s)})} \right)^{\gamma-1} \left(1 + \frac{\left(\frac{\kappa w_\infty}{w(a_s)} \right) \left(\kappa - \frac{\kappa w_\infty}{w(a_s)} \right)}{(Z^* + \frac{\kappa w_\infty}{w(a_s)})^2} \right) \right]^{\frac{1}{\gamma}} \quad (21)$$


$$\beta = \frac{1}{\gamma B_0} \left(1 - \left(\frac{M(M+\kappa)}{\alpha w(a_s)(M + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma \right) \quad (22)$$

$$\frac{B^*}{B_0} = \frac{1 - \left(\frac{(F^*+M)(F^*+M+\kappa)}{\alpha w(a_s)(F^*+M + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma}{1 - \left(\frac{M(M+\kappa)}{\alpha w(a_s)(M + \frac{\kappa w_\infty}{w(a_s)})} \right)^\gamma} \quad (23)$$



$$w(a) = w_x(1 - e^{-\kappa(a+1)})$$





Many Thanks:

- Dr. Marc Mangel
- Collaborators at NOAA
- NMFS Sea Grant



Metamodel Details


$$\hat{\mu} = \widehat{\log(r)} \quad -\text{or}- \quad \hat{\mu} = \widehat{\log(K)}$$

$$\mathbf{x} = \left(F_{MSY}, \frac{B_{MSY}}{\bar{B}(0)} \right)$$


$$\begin{aligned}\hat{\mu} &= \beta_0 + \boldsymbol{\beta}' \mathbf{x} + f(\mathbf{x}) + \epsilon \\ f(\mathbf{x}) &\sim \text{GP}(0, \tau^2 R(\mathbf{x}, \mathbf{x}')) \\ \epsilon_i &\sim \mathcal{N}(0, \hat{\omega}_i).\end{aligned}$$

$$R(\mathbf{x}, \mathbf{x}') = \exp \left(\sum_{j=1}^2 \frac{-(x_j - x'_j)^2}{2\ell_j^2} \right)$$


High Contrast PT $\sigma = 0.12$ Data



1x Samples




2x Samples




4x Samples

Contrast




No Contrast




Mangel et al.

Fig. 4. DeYoreo et al. (2012) used both a BH-SRR and three-parameter SRR, similar to the S-SRR in a stock assessment of cowcod (*Sebastodes levis*). We show samples from posterior distributions arising from different values of steepness. Unlike most stock assessments, we plot B_{MSY}/B_0 versus F_{MSY}/M . The grey circles show the results for the BH-SRR. This curve is another way of representing the constraint placed on a stock assessment by using a BH-SRR and specifying steepness — results must lie along this curve. The black circles represent the outcome of the three-parameter SRR. The black X represents the result when steepness is asserted to be 0.6.



Logistic



Schnute, J. T., & Kronlund, A. R. (2002). Estimating salmon stock recruitment relationships from catch and escape-
ment data. Canadian Journal of Fisheries and Aquatic Sciences, 59(3), 433–449.