MIZORAM PUBLIC SERVICE COMMISSION

TECHNICAL COMPETITIVE EXAMINATION FOR JUNIOR GRADE OF MIZORAM HEALTH SERVICES (COMBINED)

(AYUSH SUB-CADRE)

UNDER HEALTH & FAMILY WELFARE DEPARTMENT, GOVERNMENT OF MIZORAM.

(SEPTEMBER-2025)

TECHNICAL SUBJECT PAPER-I
(ALLIED SUBJECT)

Time Allowed : 2 hours FM : 200

All questions carry equal marks of 2 each.
Attempt all questions.

1. Dysmenorrhea is caused by:
(a) Sudden increase in progesterone (b) Prolactin
(c) Secretory endometrium (d) Venous congestion

2. Most common type of fibroid:
(a) Cervical (b) Subserosal
(c) Intramural (d) Submucosal

3. Most common ovarian cyst to undergo torsion is:
(a) Ovarian fibroma (b) Brenners tumour
(c) Benign cystic teratoma (d) Serous cyst adenoma

4. Duration of second stage of labour in primigravida is:
(a) Less than 1 hour (b) 1-2hours
(c) 2-3 hours (d) 3-4 hours

5. MTP act allows abortion till?
(a) 12 weeks (b) 24 weeks
(c) 15 weeks (d) 20 weeks

6. Fetal heart sound can be auscultated at:
(a) 8-10 weeks (b) 18-20 weeks
(c) 10-12 weeks (d) 14-16 weeks

7. Softening of the cervix during pregnancy is called:
(a) Chadwick's sign (b) Goodell's sign
(c) Jacquemier's sign (d) Piskacek"s sign

8. Most specific marker of neural tube defects is:
(a) Alpha fetoprotein (b) Pseudocholinesterase
(c) Acetylcholinesterase (d) Inhibin

9. Engagement of foetal head is with reference to:
(a) Suboccipito frontal diameter
(b) Occipito frontal diameter
(c) Biparietal diameter
(d) Bitemporal diameter

10. Lochia rubra is seen upto:
(a) 5 days
(b) 10 days
(c) 15 days
(d) 20 days

11. Most common cause of postpartum haemorrhage:
(a) Traumatic
(b) Blood coagulopathy
(c) Atonic uterus
(d) Sheehan syndrome

12. Caput succedeum is said to occur in baby:
(a) Within 24 hours
(b) Within 48 hours
(c) 2-3 weeks
(d) 2-3 months

13. Assessment of progress labour is best done by:
(a) Rupture of membrane
(b) Uterine contraction
(c) Doppler
(d) Partograph

14. Normal Crown-Rump length at birth is:
(a) 20-25 cm
(b) 38-50 cm
(c) 60-72 cm
(d) 78-85 cm

15. Antepartum haemorrhage is caused by all, except:
(a) Placenta previa
(b) Abruptio placenta
(c) Circumvallate placenta
(d) Placenta acreta

16. The most common type of female pelvis is:
(a) Gynecoid
(b) Android
(c) Anthropoid
(d) Platypelloid

17. Shortest diameter of Gynecoid pelvis is:
(a) Bitemporal
(b) Posterior sagittal
(c) Diagonal conjugate
(d) Obstetric conjugate

18. Still birth rate includes babies dead after:
(a) 20 weeks
(b) 24 weeks
(c) 28 weeks
(d) 30 weeks

19. Precocious puberty is those who menstruate in:
(a) Before 10 years
(b) 10-12 years
(c) 12-14 years
(d) 14-16 years

20. Ashermann's syndrome is characterized by:
(a) Amenorrhoea
(b) Dysmenorrhoea
(c) Leucorrhoea
(d) Metrorrhagia

21. Menorrhagia is when blood loss is more than:
(a) 50 ml
(b) 80 ml
(c) 110 ml
(d) 150 ml

22. Episiotomy is best done:
(a) Medially
(b) Laterally
(c) J Shaped
(d) Mediolaterally

23. Antenatally the most useful measurement in ultrasound is:
(a) Crown rump length
(b) Femur length
(c) Biparietal diameter
(d) Abdominal circumference

24. Face to pubis delivery is common in which type of pelvis:
(a) Platypelloid (b) Anthrapoid
(c) Android (d) Gynaecoid

25. The following is a characteristic symptom of chronic pelvic inflammatory disease:
(a) Dysuria (b) Constipation
(c) Backache (d) Fever

26. Which classification is used for burns in children:
(a) Wallace (b) Wilson
(c) Lund and Browder (d) Rule of nine

27. Concentration of sodium in RL is:
(a) 109 (b) 120
(c) 154 (d) 130

28. What is the most important aspect of management of burn injury in the first 24 hours?
(a) Antibiotics (b) Dressing
(c) Fluid resuscitation (d) Escharotomy

29. Best prognostics factor for head injury is:
(a) Glasgow coma scale (b) Mode of injury
(c) CT (d) None of the above

30. Most common cause of upper GI bleed.
(a) Esophageal varices (b) Peptic ulcer
(c) Trauma (d) Erosive gastritis

31. Kehr's sign is:
(a) Pain in the left hypochondrium due to splenic injury
(b) Carpopedal spasm of the hand and wrist
(c) Purple or bluish discoloration of periumbilical area
(d) Acute pain in tip of left shoulder due to ruptured spleen

32. Which renal stone is radiolucent?
(a) Uric acid
(b) Calcium oxalate
(c) Calcium phosphate
(d) Struvite

33. Diagnosis of a patients presenting with sudden headache, vomiting and unconsciousness.
(a) Subarachnoid haemorrhage (b) Subdural haemorrhage
(c) Extradural haemorrhage (d) Intracerebral hemorrhage

34. Risus sardonicus is seen in all except:
(a) Tetanus (b) Wilsons disease
(c) Strychnine poisoning (d) Epilepsy

35. Types of gastric cancer are the following except:
(a) Adenocarcinoma (b) Lymphoma
(c) Leiomyosarsoma (d) Astrocytoma

36. Mallory weiss tear occur at:
(a) Duodeno-jejunal flexure (b) Ileo-caecal junction
(c) Gastro-oesophageal junction (d) Colo-rectal junction

37. Stress fracture usually occurs at:
(a) 1st metacarpal (b) 2nd metacarpal
(c) Tibia (d) 2nd metatarsal

38. Carpal bone which fractures commonly:
(a) Scaphoid (b) Lunate
(c) Hamate (d) Pisiform

39. Most common joint to undergo dislocation is:
(a) Shoulder (b) Hip
(c) Patello femoral (d) Radio-ulnar

40. Pond fracture is seen in:
(a) Children (b) Adolescents
(c) Adults (d) Elderly

41. Curling's ulcer is a complication of:
(a) Gastrojejunostomy (b) Severe burns
(c) Old scar of burns (d) Head injury

42. Sun burns are:
(a) First degree burns (b) Second degree burns
(c) Third degree burns (d) Could be any of the above

43. Most common cause of surgical wound infection is:
(a) Staph aureus (b) Streptococcus
(c) Pseudomonas (d) Anaerobes

44. "Golf hole' ureter is seen in:
(a) Ureteric calculus (b) Ureteral polyp
(c) Tuberculosis of ureter (d) Retroperitoneal fibrosis

45. Commonest presentation of bilateral ureteric stones:
(a) Chronic renal failure (b) Urinary tract infection
(c) Pain (d) Hematuria

46. Murphy's triad suggests the diagnosis of:
(a) Acute appendicitis (b) Acute cholecystitis
(c) Acute pancreatitis (d) Perforated peptic ulcer

47. Commonest cause of death in peptic ulcer patient is:
(a) Perforation (b) Haemorrhage
(c) Pyloric stenosis (d) Malignancy

48. Carcinoma rectum is suspected when:
(a) Recurrent alteration of bowel habits (b) Weight reduction
(c) Presence of occult blood in stools (d) profuse bleeding per rectum

49. Triad of jaundice, chills and fever occur in:
(a) Common bile duct stone (b) Post cholecystectomy
(c) Acute hepatitis (d) Pancreatitis

50. Epistaxis in elderly patients is most often due to:
(a) Nasopharyngeal carcinoma (b) Foreign body
(c) Bleeding disorder (d) Hypertension

51. Match List-I (Deficiency) with List-II(Disease)-

List-I (Deficiency)	**List-II** (Disease)
A-Zinc	1-Follicular Keratosis
B-Essential fatty acid	2-Acrodermatitis Enteropathica
C-Folic acid	3-Neural tube defect
D-Factor 4	4-Tetany

(a) A-4, B-2, C-1, D-3 (b) A-3, B-2, C1, D-4
(c) A-2, B-1, C-3, D-4 (d) A-1, B-2, C-4, D-3

52. Most common cause of hyperthyroidism:
(a) Thyroid adenoma (b) Thyroid hyperplasia
(c) Thyroid carcinoma (d) Graves' disease

53. Hirsutism may be found in any of these disorders, except:
(a) Cushing's syndrome (b) Hypothyroidism
(c) Congenital adrenal hyperplasia (d) Polycystic Ovarian syndrome

54. Dawn phenomenon is a feature of:
(a) Post prandial coma (b) Breakfast hypoglycemia
(c) Early morning hypoglycemia (d) Early morning hyperglycemia

55. Pemberton sign is seen in:
(a) Retrosternal goiter (b) Graves opthalmopathy
(c) Thyroid crisis (d) Addison crisis

56. HbA1c level in blood explains:
(a) Hepatorenal syndrome (b) Iron deficiency anaemia
(c) Long term status of blood sugar (d) Acute rise of sugar

57. Cushing's disease presents by:
(a) Increased ACTH and decreased cortisol (b) Decreased ACTH and decrease cortisol
(c) Increased ACTH and increased cortisol (d) Increased catecholamines

58. Renal vein thrombosis is most commonly associated with:
(a) Membranous glomerulopathy (b) Diabetic nephropathy
(c) Minimal change disease (d) Membrano-proliferative glomerulonephritis

59. Marker of acute renal injury are all except:
(a) Clusterin (b) Acid phosphatase
(c) Osteopontin (d) Alanine aminopeptidase

60. RBC cast is seen in:
(a) Acute tubular nephritis (b) Acute pyelonephritis
(c) Acute interstitial nephritis (d) Acute glomerulonephritis

61. Conjugated hyperbilirubinaemia is seen in:
(a) Gilbert's syndrome (b) Dubin Jhonson syndrome
(c) Criggler Najjar syndrome (d) Lucy Driscoll syndrome

62. Which of the following hepatitis viruses has significant perinatal transmission?
(a) Hepatitis A virus (b) Hepatitis B virus
(c) Hepatitis C virus (d) Hepatitis E virus

63. Most common cause of LVH:
(a) MS (b) AR
(c) Mitral valve prolapse (d) Hypertension

64. Water hammer pulse is seen in:
(a) Aortic regurgitation (b) Aortic stenosis
(c) Mitral regurgitation (d) Aortic stenosis and mitral regurgitation

65. Carpal tunnel syndrome is due to compression of:
(a) Radial nerve (b) Median nerve
(c) Ulnar nerve (d) Flexor tendons

66. Koplik's spots are seen in:
(a) Measles (b) Mumps
(c) Diptheria (d) Rubella

67. The following parasitic infection are commonly seen in India- except:
(a) Schistosomiasis (b) Pinworms
(c) Trichomoniasis (d) Microfilaria

68. Cysticercosis is a cyst of:
(a) Echinococcus (b) Trichinella
(c) Taenia solium (d) Dracunlus

69. Reiter's syndrome is a triad of the following symptoms:
(a) Urethritis, Cervicitis, and Salpingitis (b) Urethritis, Conjunctivitis and Arthritis
(c) Urethritis, Cystitis and Pyelitis (d) Urethritis, Cystitis and Proctitis

70. Macrocytic hypochromic anaemia is associated with:
(a) Vitamin A deficiency (b) Iron deficiency anaemia
(c) Hypoplastic anaemia (d) Vit B12 deficiency

71. Fasciculation is seen in:
(a) LMN type of lesion (b) UMN type of lesion
(c) Myoneural junction (d) Peripheral neuropathy

72. Amyotrophic lateral sclerosis involve:
(a) Dorsal root ganglion (b) Ventral root ganglion
(c) Posterior horn cells (d) Anterior horn cells

73. Which of the following is the most common location of hypertensive haemorrhage:
(a) Pons (b) Putamen
(c) Thalamus (d) Cerebellum

74. In Alzheimer's disease which part if the brain is involved:
(a) Cerebral cortex (b) Cerebellum
(c) Pons (d) Hypothalamus

75. Which immunoglobulin is elevated in a case of chronic allergy hypersensitivity?
(a) Ig-E (b) Ig-M
(c) Ig-A (d) Ig-G

76. First enzyme to be raised in MI is:
(a) Troponin-I (b) LDH
(c) CPK-MB (d) Myoglobin

77. Most common type of schizophrenia:
(a) Paranoid (b) Catatonic
(c) Residual (d) Undifferentiated

78. Pterygium of nail is characteristically seen in:
(a) Psoriasis (b) Lichen planus
(c) Tinea unguim (d) None of the above

79. Alopecia areata is due to:
(a) Autoimmune (b) Lichenoid reaction
(c) Estrogen stimulation (d) Androgen stimulation

80. Scrub typhus is caused by:
(a) Coxiella burnetti (b) Salmonella Typhi
(c) Rickettsia tsutsugamushi (d) Rickettsia prowazekii

81. All of the following CSF findings are present in tuberculous meningitis, except:
(a) Raised sugar levels (b) Cob web formation
(c) Low chloride levels (d) Raised protein levels

82. Autism is:
(a) Social and language communication problem (b) Metabolic disease
(c) Neurodevelopmental disorder (d) Mostly due to hypothalamus damage

83. Genital warts are caused by:
(a) Human Papilloma Virus (b) Herpes simples virus
(c) Esptein Barr virus (d) Lymphogranuloma venereum

84. HLA-B27 is associated with all except:
(a) Ankylosing spondylitis (b) Reiters syndrome
(c) Behcets syndrome (d) Psoriatic spondylitis

85. Bence Jones proteins are:
(a) Heavy chain IgG (b) Light chain IgG
(c) Seen in lymphoma (d) Present in bone marrow

86. All are seen in congestive cardiac failure except:
(a) Kerley B lines (b) Pleural effusion
(c) Ulcerative colitis (d) Bronchiectasis

87. Vocal resonance is increased in:
(a) Collapse (b) Pneumothorax
(c) Pleural effusion (d) Consolidation

88. The minimum duration of symptoms to be associated with chronic bronchitis is:
(a) 6 months (b) 1 year
(c) > 2 years (d) > 5 years

89. In benign hypertension, commonest vascular pathology is:
(a) Fatty infiltration of intima (b) Hyaline arteriosclerosis
(c) Fibrinoid necrosis (d) Atherosclerosis

90. Most specific enzyme for Myocardial infraction is:
(a) CPK-MM (b) CPK-BB
(c) CPK-MB (d) LDH

91. Zinc deficiency has all the manifestations except:
(a) Alopecia (b) Cardiomyopathy
(c) Growth retardation (d) Hypogonadism

92. Generalized oedema can be produced by deficiency of:
(a) Vit A (b) Thiamine
(c) Vit D (d) Folic acid

93. Hypervitaminosis A can cause all except:
(a) Anorexia (b) Loss of hair
(c) Peripheral neuritis (d) Hypotension

94. Most common presentation of endemic goitre is:
(a) Hyperthyroidism (b) Hypothyroidism
(c) Hoarseness of voice (d) Swelling

95. Which is not seen in Irritable bowel syndrome:
(a) Weight loss (b) Mucus per rectum
(c) Altered bowel habit (d) Abdominal pain

96. Most common presentation of primary biliary cirrhosis is:
(a) Jaundice (b) Abdominal mass
(c) Pruritus (d) Pain

97. ABO haemolytic disease of new born occurs when the blood group of mother is:
(a) A (b) B
(c) AB (d) O

98. Earliest change in Iron deficiency anemia is:
(a) Decreased serum iron (b) Decreased serum ferritin
(c) Decreased TIBC (d) Decreased transferrin saturation

99. Auer rods are seen in which type of leukaemia:
(a) ALL (b) AML
(c) CLL (d) CML

100. Satellite lesion is seen in which leprosy:
(a) Tuberculoid (b) Borderline tuberculoid
(c) Lepromatous (d) Histoid

* * * * * * *