MIZORAM PUBLIC SERVICE COMMISSION

TECHNICAL COMPETITIVE EXAMINATION FOR JUNIOR GRADE OF MIZORAM HEALTH SERVICES (COMBINED) (GDMO SUB-CADRE) UNDER HEALTH & FAMILY WELFARE DEPARTMENT, GOVERNMENT OF MIZORAM. (SEPTEMBER-2025)

TECHNICAL SUBJECT PAPER-I (ALLOPATHY)

Time Allowed : 2 hours FM : 200

All questions carry equal marks of 2 each.
Attempt all questions.

1. All are common causes of persistent diarrhoea except:
(a) Giardia (b) Rotavirus
(c) E. histolytica (d) C. difficile

2. Which is not a cause of metabolic acidosis?
(a) Alcoholism (b) Uncontrolled hyperglycemia
(c) Starvation (d) Liddle Syndrome

3. Geriatric giants do not include:
(a) Urinary incontinence (b) Frequent falls
(c) Social isolation (d) Cognitive impairment

4. Angiotensinogen is converted to angiotensin in kidney by:
(a) ACE (b) Aldosterone
(c) Renin (d) BNP

5. Activation of RAAS results in:
(a) increased afterload (b) increased preload
(c) decreased afterload (d) decreased preload

6. First antigen to appear in Hep B infection is:
(a) HBcAg (b) HBeAg
(c) HBsAg (d) HBxAg

7. Which is a cause of conjugated hyperbilirubinemia?
(a) Gilbert's Syndome (b) Rotor's syndrome
(c) Crigler-Najjar syndrome 2 (d) Haemolytic disease

8. Pathophysiology of right sided heart failure include all except:
(a) Increased afterload (b) Increased preload
(c) Decreased contractility (d) Decreased preload

9. Most common cause of upper Gi bleed is:
(a) Oesophageal varix (b) Peptic ulcer
(c) Upper GI tumour (d) Mallory-Weiss tear

10. Most common route of Hepatitis C infection transmission is:
(a) Faecal-orally (b) Percutaneous
(c) Sexual (d) Perinatal

11. Serological test for Hep B includes all except:
(a) Anti-HBcAg (b) HBeAg
(c) HBcAg (d) Anti-HBeAg

12. Anemia is a well recognised side effect of:
(a) Efavirenz (b) Tenofovir Disoproxil fumarate
(c) Lopinavir/ritonavir (d) Zidovudine

13. Infection which can cause Q fever?
(a) Enterohemorrhagic E. Coli (0157:H7) (b) Campylobacter
(c) Salmonella (d) Coxiella burnetii

14. All of the following are water soluble vitamin except:
(a) Vitamin B (b) Vitamin C
(c) Vitamin D (d) All of the above

15. Mid systolic murmur is a feature of all except.
(a) Pulmonic stenosis (b) HOCM
(c) Mitral valvulitis (d) Aortic stenosis

16. CREST consist of all except:
(a) Calcinosis (b) Raynauds phenomenon
(c) Sclerodactyly (d) Tinea

17. Which of the following is not associated with erythema nodosum?
(a) Sarcoidosis (b) Hepatitis B
(c) Crohn's Disease (d) Leukemia

18. All Anti-tubercular drugs are hepatotoxic except:
(a) Pyrazinamide (b) Rifampicin
(c) Ethambutol (d) Isoniazid

19. Fondaparinux is an anticoagulant with a mechanism of action as:
(a) Factor Xa inhibitor (b) Direct thrombin inhibition
(c) Vitamin K antagonist (d) Antiplatelet activity

20. What term is used to describe loss of muscle mass and strength that occurs with aging, often leading to functional decline and increased risk of falls?
(a) Sarcopenia (b) Osteoporosis
(c) Atherosclerosis (d) Hypertension

21. What is the temperature at which platelet is stored under standard blood banking conditions?
(a) 20-24°C (b) 2-6°C
(c) -40°C (d) None of the above

22. A 50 year old man presents with a characteristic sudden onset '*thunderclap headache*'. What is the most likely diagnosis for this condition?
(a) Sub-arachnoid hemorrhage (b) Sub-dural hemorrhage
(c) Extra-dural hemorrhage (d) None of the above

23. Root value of bicep jerk is:
(a) C5-C6 (b) L5-S1
(c) L3-L4 (d) D12-L1

24. The most common cause of chronic kidney disease is:
(a) Diabetes
(b) Hypertension
(c) Glomerulonephritis
(d) Polycystic kidney disease

25. Which of the following produces antibodies in the body?
(a) Null cells
(b) Helper T-cells
(c) Killer T-cells.
(d) Plasma cells

26. In gene therapy; which infective agent is used as a vehicle?
(a) E.coli
(b) Pneumococcus
(c) Corynebacterium
(d) Adenovirus

27. Which of the following is not an autoimmune disorder?
(a) Pemphigus vulgaris
(b) Anti phospholipid syndrome
(c) Wilson's disease
(d) Hashimoto's thyroiditis

28. Vitamin C toxicity may result in:
(a) High uric acid
(b) Oxalate stones in kidney
(c) Loss of libido
(d) Sideroblastic anaemia

29. Serum calcium level is lowered by the deficiency of:
(a) Calcitonin
(b) Thyroxine
(c) Parathormone
(d) Both calcitonin and parathormone

30. Raspberry tongue is found in:
(a) Scarlet fever
(b) Glandular fever
(c) Yellow fever
(d) Rheumatic fever

31. Regarding melaena, which statement is false?
(a) At least 60ml of blood is required
(b) Blood should remain at least 4 hours within the gut
(c) Black tarry semi solid stool
(d) Offensive in odour

32. The most common gastrointestinal disorder in a community is:
(a) Diverticulitis
(b) Duodenal ulcer
(c) Reflux oesophagitis
(d) Irritable bowel syndrome

33. Central cyanosis is not found in:
(a) Acute pulmonary oedema
(b) Fallot's tetralogy
(c) Left-to-right shunt
(d) Transposition of great vessels

34. Left parasternal heave is diagnostic of:
(a) Left ventricular hypertrophy
(b) Right atrial hypertrophy
(c) Right ventricular hypertrophy
(d) Hypertrophic cardiomyopathy

35. Woody dullness in percussion over chest is classically found in:
(a) Thickened pleura
(b) Consolidation
(c) Collapse of the lung
(d) Pleural effusion

36. Bronchial breath sound is found in all *except:*
(a) Collapse with patent bronchus
(b) Bronchial asthma
(c) Superficial, big, empty cavity with patent bronchus
(d) Broncho pleural fistula

37. Which is not a symptom of raised intracranial tension?
(a) Altered consciousness (b) Headache
(c) Non-projectile vomiting (d) Convulsions

38. Neck rigidity is not found in:
(a) After epileptic seizure (b) Meningism
(c) Hysteria (d) Tetanus

39. Crossed hemiplegia indicates the site of lesion in:
(a) Internal capsule (b) Cortex
(c) Cervical spine (d) Brainstem

40. Which of the following is not a feature of UMN palsy?
(a) Spasticity (b) Babinski's sign
(c) Clonus (d) Fasciculations

41. Commonest cerebrovascular accident (CVA) is:
(a) Cerebral haemorrhage (b) Cerebral thrombosis
(c) Cerebral embolism. (d) Subarachnoid haemorrhage

42. Which of the following is not a criteria for diagnosis of nephrotic syndrome?
(a) Hypertension (b) Massive proteinuria
(c) Hyperlipidaemia (d) Anasarca

43. Oliguria is:
(a) <50 ml urine/24hrs (b) <100 ml urine/24hrs
(c) <200 ml urine/24hrs. (d) <400 mL urine/24hr

44. Renal vein thrombosis in adults is seen in:
(a) Amyloidosis (b) Acute glomerulonephritis
(c) Interstitial nephritis (d) Horse shoe kidney

45. Auer rods are found in:
(a) Acute myeloid leukaemia (AML) (b) Blast crisis of CML
(c) Acute lymphatic leukaemia (ALL) (d) Blast crisis of CLL

46. Commonest cause of jaundice in thalassaemia is:
(a) Viral hepatitis C (b) Iron deposition in liver
(c) Viral hepatitis B (d) Haemolysis

47. Stevens-Johnson syndrome is classically seen in all *except:*
(a) Carbamazepine (b) Cotrimoxazole
(c) Corticosteroids (d) Thiacetazones

48. Which of the following is a viral infection?
(a) Lichen planus (b) Lupus vulgaris
(c) Psoriasis (d) Molluscum contagiosum

49. Raynaud's phenomenon is not a feature of:
(a) Hyper viscosity syndrome (b) Ergot ingestion
(c) Coarctation of aorta (d) Dermatomyositis

50. Commonest cause of coma in a diabetic is:
(a) Diabetic ketoacidosis (b) Lactic acidosis
(c) Hyperosmolar non-ketotic coma. (d) Hypoglycaemia

51. The first line drug used to treat seizure in neonate is:
(a) Phenytoin (b) Diazepam
(c) Levetiracetam (d) Phenobarbitone

52. During positive pressure ventilation in newborn resuscitation, heart rate assessment is done every
(a) 10 seconds (b) 20 seconds
(c) 30 seconds (d) 60 seconds

53. A 5 year old boy presenting with fever for 1 week with fatiguability and skin bleeds, and noted to have hepatosplenomegaly is more likely to have :
(a) Aplastic anemia (b) Immune thrombocytopenia
(c) Acute leukemia (d) Drug induced thrombocytopenia

54. The characteristic coagulation abnormality seen in haemophilia is :
(a) Prolonged PT, APTT (b) Normal PT, prolonged APTT
(c) Normal APTT, prolonged PT (d) Normal PT and APTT, but low platelets

55. Which of the following viral infections is associated with a strong predisposition to Hodgkin lymphoma?
(a) Herpes simplex infection (b) Varicella
(c) Parvovirus infection (d) Epstein-Barr virus infection

56. Which of the following drugs can be used for intermittent prophylaxis in febrile seizure?
(a) Valparin (b) Phenytoin
(c) Clobazam (d) Levipil

57. The recommended vitamin D intake in children < 1 year is :
(a) 200 IU/day (b) 400 IU/day
(c) 600 IU/day (d) 800 IU/day

58. The most profound consequence of undernutrition is :
(a) Premature death (b) Repeated infections
(c) Stunting (d) Cell damage

59. The preferable antibiotic in neonate with pertussis is:
(a) Azithromycin (b) Erythromycin
(c) Clarithromycin (d) Amoxycillin

60. Hand foot mouth disease is one of the more distinctive rash syndromes, it is MOST frequently caused by:
(a) Coxsackievirus A6 (b) Coxsackievirus A16
(c) Coxsackievirus B2 (d) Coxsackievirus B12

61. Gastroesophageal reflux disease is the most common esophageal disorder in children of all ages. Of the following, the most common symptom in infancy is:
(a) Excessive crying (b) Regurgitation
(c) Failure to thrive (d) Sleeping disturbances

62. Guillain- barre syndrome can be a sequel in the following foodborne bacterial illness:
(a) Bacillus anthracis (b) Bacillus cereus
(c) Campylobacter jejuni (d) Clostridium botulinum

63. The low osmolality WHO oral rehydration solution has the following per liter EXCEPT:
(a) 75 mEq of sodium (b) 50 mEq of chloride
(c) 20 mEq of potassium (d) Total osmolality of 245 mOsm

64. There are many anomalies associated with anorectal malformations. Of the following, the MOST common are:
(a) Kidney anomalies
(b) Cardiac anomalies
(c) Esophageal atresia
(d) Spina bifida

65. Biliary sludge or cholelithiasis can be detected in > 40 % of children who are treated with prolonged course of:
(a) Cefotaxime
(b) Meropenem
(c) Erythromycin
(d) Ceftriaxone

66. Habit cough (psychogenic cough) must be considered in any child with a cough that has lasted for weeks or months, that has been refractory to treatment and is:
(a) Abrupt
(b) Barking quality
(c) All day, never during sleep
(d) Paroxysmal

67. Recurrent pneumonia is defined as-
(a) 2 or more episodes in a single year, with radiographic clearing between occurences
(b) 2 or more episodes in a single year, without radiographic clearing between occurences
(c) 3 or more episodes in a single year, with radiographic clearing between occurences
(d) 4 or more episodes in a single year

68. The most consistent clinical manifestation of pneumonia is-
(a) Subcostal retraction
(b) Tachypnea
(c) Fever
(d) Cough

69. The most common cyanotic cardiac lesion to present in the newborn period is:
(a) Dextroposed transposition of the great arteries
(b) Hypoplastic left heart syndrome
(c) Tetralogy of fallot
(d) Truncus arteriosus

70. The most common cause of syncope in children is:
(a) Wolff Parkinson white syndrome
(b) Prolonged QT syndrome
(c) Atrioventricular block
(d) Neurocardiogenic syncope

71. Surgical repair of VSD at the time of diagnosis should be considered in:
(a) Small supracristal VSD
(b) Nonrestrictive VSD
(c) Hemodynamically significant VSD
(d) Restrictive VSD

72. The best assessment of iron overload for patients with thalassemia major is achieved by:
(a) Liver MRI
(b) Bone marrow biopsy
(c) Serum iron
(d) Serum ferritin

73. Which of the following viruses is more likely to be associated with chronic immune thrombocytopenia?
(a) Cytomegalovirus
(b) Parvovirus
(c) Influenza virus
(d) Human immune deficiency virus

74. In hematuria, renal biopsy is indicated in all the following EXCEPT:
(a) Persistent microscopic hematuria
(b) Decreased renal function
(c) Hypertension
(d) Persistent high serum complement C3

75. The most common cause of permanent Congenital hypothyroidism is:
(a) Dyshormonogenesis
(b) Thyroid agenesis
(c) Iodine deficiency
(d) Defect of iodide transport

76. The following are true about pathological jaundice EXCEPT:
(a) Clinical jaundice detected before 48 hours of age
(b) Rise in serum bilirubin by more than 5 mg/ dL/ day
(c) Serum bilirubin more than 15 mg/dL
(d) Clay/white coloured stool

77. All are included under '*initial steps*' in neonatal resuscitation EXCEPT:
(a) Dry baby and remove wet linen
(b) Cut the cord and shift to radiant warmer
(c) Stimulate baby by rubbing
(d) Initiate breastfeeding

78. According to Neonatal resuscitation protocol (NRP), initial positive pressure ventilation (PPV) in a term baby should be done using:
(a) Oxygen at Fio2 of 50 %
(b) Oxygen at FiO2 of 21-30%
(c) Oxygen at FiO2 of 100%
(d) Oxygen in room air

79. All are common complications associated with malnutrition EXCEPT:
(a) Encephalopathy
(b) Dehydration
(c) Hypoglycaemia
(d) Infection

80. Stunting in a child is:
(a) Weight/height < -2SD
(b) Height/weight < -2SD
(c) Height/age < -2SD
(d) BMI < -2SD

81. How much measurement of mid-upper arm circumference is considered as Severe acute malnutrition (SAM)?
(a) Less than 12 cm
(b) Less than 11.5 cm
(c) Less than 11 cm
(d) Less than 10.5 cm

82. False about National Tuberculosis Eradication Program (NTEP):
(a) Program aims to eliminate tuberculosis in India by 2028
(b) Operates on four pillars: Detect, Treat, Prevent and Build
(c) The program collaborates with the private sector to enhance TB detection and management
(d) Provides free TB testing and treatment

83. False about birth dose vaccine is:
(a) BCG can be given till 1 year of age
(b) OPV is given as 5 drops orally
(c) Hepatitis B should be given at birth or within 24 hours of birth
(d) BCG is given intradermally

84. Marasmus is characterized by the following EXCEPT:
(a) Severe wasting
(b) Baggy pants appearance
(c) Monkey facies
(d) Mild to moderate edema

85. Polymorphic rash is seen in:
(a) Measles
(b) Mumps
(c) Scarlet fever
(d) Chicken pox

86. Viral hepatitis in children is most commonly caused by:
(a) Hepatitis A
(b) Hepatitis B
(c) Hepatitis C
(d) Hepatitis D

87. False about Dengue fever:
(a) Caused by arbovirus of family Flaviviridae
(b) Spread by mosquito of the species Aedes aegypti
(c) Positive tourniquet test may be seen
(d) Febrile phase is characterised by shock

88. Persistent diarrhea is:
(a) Diarrhea lasting for more than 7 days
(b) Diarrhea lasting for more than 14 days
(c) Diarrhea lasting for more than 3 weeks
(d) None of the above

89. Common organisms causing pneumonia in children are all EXCEPT:
(a) Pneumococcus
(b) Hemophilus influenza type b
(c) Rotavirus
(d) Respiratory syncytial virus

90. Conjugated hyperbilirubinemia is seen in:
(a) Breastmilk jaundice
(b) Neonatal cholestasis
(c) Gilbert syndrome
(d) Hypothyroidism

91. A 12-month-old infant with cold and cough has SpO2 of 88% in room air. According to IMNCI, this infant will be classified as having:
(a) Pneumonia
(b) Septicemia
(c) Severe Pneumonia
(d) Mild Pneumonia

92. For diagnosing Acute Rheumatic fever, which one of the following is a minor criteria?
(a) Carditis
(b) Arthralgia
(c) Polyarthritis
(d) Subcutaneous nodules

93. Which of the following is a cyanotic heart disease:
(a) VSD
(b) ASD
(c) PDS
(d) Tricuspid Atresia

94. Which of the following is a constant feature seen in Downs syndrome?
(a) Hypotonia
(b) Acute leukaemia
(c) Congenital heart defects
(d) Brachycephaly

95. Fever, barking cough and stridor is a common presentation of:
(a) Pneumonia
(b) Bronchiolitis
(c) Croup
(d) Empyema

96. All of the following signs are seen in a child with pneumothorax EXCEPT:
(a) Decreased breath sound
(b) Tympanic note on percussion
(c) Mediastinal shift to the opposite site
(d) Crowding of ribs

97. Which of the following is NOT a usual characteristic of cyanotic congenital heart disease?
(a) Elevated blood pressure in arms
(b) Digital clubbing
(c) Polycythaemia
(d) Exercise intolerance

98. Minimal change nephrotic syndrome is characterized by all of the following EXCEPT:
(a) Edema
(b) Hypercholesterolemia
(c) Hypertension
(d) Hypoproteinaemia

99. In febrile seizure, the following statements are true EXCEPT:
(a) Giving paracetamol does not prevent recurrence
(b) Genetic predisposition
(c) 2 types - simple and complex febrile seizure
(d) Occur in newborn

100. Seizure during neonatal period can be caused by:
(a) Hypoglycaemia
(b) Hypocalcaemia
(c) Hypoxia
(d) All of the above

* * * * * * *