

## Research Article

# The Effect of Smoking on COVID-19 Symptom Severity: Systematic Review and Meta-Analysis

Askin Gülsen <sup>1</sup>, Burcu Arpinar Yigitbas <sup>2</sup>, Berat Uslu <sup>2</sup>, Daniel Drömann,<sup>1</sup>  
and Oguz Kilinc <sup>3</sup>

<sup>1</sup>Department of Pneumology, Airway Research Center North (ARC�), Member of the German Center for Lung Research (DZL), University of Luebeck, Germany

<sup>2</sup>Department of Pneumology, Yedikule Chest Disease and Thoracic Surgery Hospital, Istanbul, Turkey

<sup>3</sup>Department of Pneumology, Dokuz Eylül University, Izmir, Turkey

Correspondence should be addressed to Askin Gülsen; askingulsen@hotmail.com

Received 2 June 2020; Accepted 19 August 2020; Published 9 September 2020

Academic Editor: Roberto Walter Dal Negro

Copyright © 2020 Askin Gülsen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** Coronavirus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome coronavirus 2 (SAR2-COV-2) and was first identified in Wuhan, China, in December of 2019, but quickly spread to the rest of the world, causing a pandemic. While some studies have found no link between smoking status and severe COVID-19, others demonstrated a significant one. The present study aimed to determine the relationship between smoking and clinical COVID-19 severity via a systematic meta-analysis approach. **Methods.** We searched the Google Scholar, PubMed, Scopus, Web of Science, and Embase databases to identify clinical studies suitable for inclusion in this meta-analysis. Studies reporting smoking status and comparing nonsevere and severe patients were included. Nonsevere cases were described as mild, common type, nonintensive care unit (ICU) treatment, survivors, and severe cases as critical, need for ICU, refractory, and nonsurvivors. **Results.** A total of 16 articles detailing 11322 COVID-19 patients were included. Our meta-analysis revealed a relationship between a history of smoking and severe COVID-19 cases (OR = 2.17; 95% CI: 1.37–3.46;  $P < .001$ ). Additionally, we found an association between the current smoking status and severe COVID-19 (OR = 1.51; 95% CI: 1.12–2.05;  $P < .008$ ). In 10.7% (978/9067) of nonsmokers, COVID-19 was severe, while in active smokers, severe COVID-19 occurred in 21.2% (65/305) of cases. **Conclusion.** Active smoking and a history of smoking are clearly associated with severe COVID-19. The SARS-COV-2 epidemic should serve as an impetus for patients and those at risk to maintain good health practices and discontinue smoking. The trial is registered with the International Prospective Register of Systematic Reviews (PROSPERO) CRD42020180173.

## 1. Introduction

Coronavirus disease 2019 (COVID-19), which is caused by severe acute respiratory syndrome coronavirus 2 (SAR2-COV-2), was first identified in Wuhan, China, in December of 2019. It has subsequently spread across the world, causing a global pandemic. This highly contagious disease has thus far infected 23.4 million people worldwide and killed approximately 808000 patients, yielding a case fatality rate (CFR) that varies between 0.7 and 12.7% (average: 3.4%) [1, 2].

COVID-19 primarily targets lung epithelial cells, causing viral pneumonia and acute respiratory distress syndrome (ARDS), especially in elderly patients. Therefore, mortality is higher in the elderly and in patients with at least one accompanying comorbid disease [3]. In the last report issued by the Centers for Disease Control and Prevention Institute, the incidence of respiratory disease was 9.2% in patients diagnosed with a severe COVID-19 clinical course [4]. Chronic obstructive pulmonary disease (COPD) and asthma are also common comorbidities in severe cases and are reported in

carrying cigarette packets in the pockets, and blowing of smoke. Similarly, exposure to passive smoke can alter ACE-2 gene expression and cause immune system changes. Naturally, these patients also have the potential to have severe COVID-19 symptoms, and smoke exposure is a potential risk factor for those around the patient, including their friends and family.

Future studies should continue to collect nicotine consumption information, including the number of cigarettes smoked per day, passive exposure, and degree of COPD, and should evaluate the dynamics of interactions between cigarette smoking and COVID-19. In addition, the effects of various smoking habits (e.g., mild versus heavy consumption, water pipe use, and electronic cigarettes use) on the transmission of SARS-CoV-2, the clinical severity of COVID-19, and the clinical progression of COVID-19 should be investigated. Finally, the relationship between COPD severity and COVID-19, and the potential therapeutic effect of nicotine on severe COVID patients should also be examined and clarified. In addition, clinicians can pay more attention to the history of smoking of COVID-19 patients, and more further research may aim to determine mechanisms that drive or decrease this risk.

## 5. Conclusion

The present meta-analysis revealed that active smoking and a history of smoking are significantly associated with increased COVID-19 symptom severity. The SARS-CoV-2 epidemic should serve as an impetus for patients and those at risk to maintain good health practices and discontinue smoking.

## Data Availability

The data [data.oma and data.xls] used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors have stated explicitly that there are no conflicts of interest in connection with this article.

## Authors' Contributions

AG performed the concept of the manuscript, data collection, design of tables and figures, analysis and interpretation of data, and writing and drafting of the manuscript; BU performed the data collection and quality analysis; BAY performed the acquisition of data, quality analysis, and writing and revising the manuscript; DD and OK performed the major role in reviewing and revising the manuscript.

## Acknowledgments

Funding of the Federal Ministry for Science and Education (BMBF), German Center for Lung Research, is gratefully acknowledged.

## References

- [1] World Health Organisation, "Coronavirus disease (COVID-2019) situation reports," [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200824-weekly-epi-update.pdf?sfvrsn=806986d1\\_4](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200824-weekly-epi-update.pdf?sfvrsn=806986d1_4).
- [2] Coronavirus Worldometer, "Country-specific case fatality rates (CFR) of COVID-19 patients," August 2020 <https://www.worldometers.info/coronavirus/>.
- [3] J. Yang, Y. Zheng, X. Gou et al., "Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis," *International Journal of Infectious Diseases*, vol. 94, pp. 91–95, 2020.
- [4] CDC COVID-19 Response Team, CDC COVID-19 Response Team, N. Chow et al., "Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019—United States, February 12–March 28, 2020," *MMWR. Morbidity and Mortality Weekly Report*, vol. 69, no. 13, pp. 382–386, 2020.
- [5] S. Garg, L. Kim, M. Whitaker et al., "Hospitalization Rates and Characteristics of Patients Hospitalized with Laboratory-Confirmed Coronavirus Disease 2019 — COVID-NET, 14 States, March 1–30, 2020," *MMWR. Morbidity and Mortality Weekly Report*, vol. 69, no. 15, pp. 458–464, 2020.
- [6] Q. Zhao, M. Meng, R. Kumar et al., "The impact of COPD and smoking history on the severity of COVID-19: A systemic review and meta-analysis," *Journal of Medical Virology*, 2020.
- [7] G. Lippi and B. M. Henry, "Active smoking is not associated with severity of coronavirus disease 2019 (COVID-19)," *European Journal of Internal Medicine*, vol. 75, pp. 107–108, 2020.
- [8] D. Moher, A. Liberati, J. Tetzlaff, D. G. Altman, and for the PRISMA Group, "Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement," *BMJ*, vol. 339, article b2535, 2009.
- [9] G. A. Wells, B. Shea, D. O'Connell et al., *The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses* Ottawa Hospital Research Institute May 2020 [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp).
- [10] Q. Chen, Z. Zheng, C. Zhang et al., "Clinical characteristics of 145 patients with corona virus disease 2019 (COVID-19) in Taizhou, Zhejiang, China," *Infection*, vol. 48, no. 4, pp. 543–551, 2020.
- [11] Y. Feng, Y. Ling, T. Bai et al., "COVID-19 with Different Severities: A Multicenter Study of Clinical Features," *American Journal of Respiratory and Critical Care Medicine*, vol. 201, no. 11, pp. 1380–1388, 2020.
- [12] P. Goyal, J. J. Choi, L. C. Pinheiro et al., "Clinical Characteristics of Covid-19 in New York City," *The New England Journal of Medicine*, vol. 382, no. 24, pp. 2372–2374, 2020.
- [13] W. J. Guan, Z. Y. Ni, Y. Hu et al., "Clinical Characteristics of Coronavirus Disease 2019 in China," *New England Journal of Medicine*, vol. 382, no. 18, pp. 1708–1720, 2020.
- [14] W. Liu, Z. W. Tao, L. Wang et al., "Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease," *Chinese Medical Journal*, vol. 133, no. 9, pp. 1032–1038, 2020.
- [15] P. Mo, Y. Xing, Y. Xiao et al., "Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China," *Clinical Infectious Diseases*, 2020.
- [16] Q. Di, X. Yan, X. Tang et al., "Epidemiological and clinical features of 2019-nCoV acute respiratory disease cases in