

Retrieve place value and rounding

Use the scaffold, show each step and ask for an adult check when marked.

Name: _____ Date: _____

1. Read the model: Order 6,908,210, 6,890,201 and 6,980,120, then round each to the nearest 100,000. Copy or complete the first step.

2. Find the interval from -18 to 27.

3. Give the range of whole numbers that round to 3,600,000 to the nearest 100,000.

4. Miro says: Miro applies one rounding rule to every requested place. Is this correct? Explain.

Teacher answers and checking prompts

1. Read the model: Order 6,908,210, 6,890,201 and 6,980,120, then round each to the nearest 100,000. Copy or complete the first step.
Answer Order: 6,890,201; 6,908,210; 6,980,120. Rounded: 6,900,000; 6,900,000; 7,000,000.
2. Find the interval from -18 to 27.
Answer 45.
3. Give the range of whole numbers that round to 3,600,000 to the nearest 100,000.
Answer 3,550,000 to 3,649,999.
4. Miro says: Miro applies one rounding rule to every requested place. Is this correct? Explain.
Answer Mark each rounding place separately and reconsider the deciding digit each time.

Retrieve place value and rounding

Work independently. Show a complete method and check at least one answer.

Name: _____ Date: _____

1. Find the interval from -18 to 27.

2. Give the range of whole numbers that round to 3,600,000 to the nearest 100,000.

3. Create a seven-digit number that rounds down to 5,000,000 to the nearest million and up to 5,400,000 to the nearest 100,000.

4. Identify and correct this misconception: Miro applies one rounding rule to every requested place.

5. Write a complete sentence explaining how you checked one answer.

Teacher answers and checking prompts

1. Find the interval from -18 to 27.

Answer 45.

2. Give the range of whole numbers that round to 3,600,000 to the nearest 100,000.

Answer 3,550,000 to 3,649,999.

3. Create a seven-digit number that rounds down to 5,000,000 to the nearest million and up to 5,400,000 to the nearest 100,000.

Answer Answers vary, for example 5,360,000.

4. Identify and correct this misconception: Miro applies one rounding rule to every requested place.

Answer Mark each rounding place separately and reconsider the deciding digit each time.

5. Write a complete sentence explaining how you checked one answer.

Answer Answer should name an inverse, estimate, boundary, known fact or equivalent representation.

Retrieve place value and rounding

Prove, compare or generalise. Write enough reasoning for another pupil to follow.

Name: _____ Date: _____

1. Create a seven-digit number that rounds down to 5,000,000 to the nearest million and up to 5,400,000 to the nearest 100,000.

2. Explain why this reasoning fails: Miro applies one rounding rule to every requested place.

3. Create a new example that tests the same idea as: Give the range of whole numbers that round to 3,600,000 to the nearest 100,000.

4. Find a second method or representation. Compare its efficiency with your first method.

5. Write one always, sometimes or never statement about today's learning and justify it.

Teacher answers and checking prompts

1. Create a seven-digit number that rounds down to 5,000,000 to the nearest million and up to 5,400,000 to the nearest 100,000.

Answer Answers vary, for example 5,360,000.

2. Explain why this reasoning fails: Miro applies one rounding rule to every requested place.

Answer Mark each rounding place separately and reconsider the deciding digit each time.

3. Create a new example that tests the same idea as: Give the range of whole numbers that round to 3,600,000 to the nearest 100,000.

Answer Answers vary. The example and solution must preserve the mathematical structure.

4. Find a second method or representation. Compare its efficiency with your first method.

Answer Answers vary. Comparison should refer to accuracy, number structure and clarity.

5. Write one always, sometimes or never statement about today's learning and justify it.

Answer Answers vary. A valid example and counterexample should support the classification.