

Find factor pairs systematically

Use the scaffold, show each step and ask for an adult check when marked.

Name: _____ Date: _____

1. Read the model: List every factor pair of 72. Copy or complete the first step.

2. List all factors of 48.

3. How many factors does 36 have?

4. Miro says: Miro stops checking factor pairs after reaching 5. Is this correct? Explain.

Teacher answers and checking prompts

1. Read the model: List every factor pair of 72. Copy or complete the first step.
Answer 1 and 72; 2 and 36; 3 and 24; 4 and 18; 6 and 12; 8 and 9.
2. List all factors of 48.
Answer 1, 2, 3, 4, 6, 8, 12, 16, 24, 48.
3. How many factors does 36 have?
Answer 9 factors.
4. Miro says: Miro stops checking factor pairs after reaching 5. Is this correct? Explain.
Answer Continue until the first factor meets or passes its paired factor.

Find factor pairs systematically

Work independently. Show a complete method and check at least one answer.

Name: _____ Date: _____

1. List all factors of 48.

2. How many factors does 36 have?

3. Find a number below 100 with exactly 10 factors.

4. Identify and correct this misconception: Miro stops checking factor pairs after reaching 5.

5. Write a complete sentence explaining how you checked one answer.

Teacher answers and checking prompts

1. List all factors of 48.

Answer 1, 2, 3, 4, 6, 8, 12, 16, 24, 48.

2. How many factors does 36 have?

Answer 9 factors.

3. Find a number below 100 with exactly 10 factors.

Answer 48 is one example.

4. Identify and correct this misconception: Miro stops checking factor pairs after reaching 5.

Answer Continue until the first factor meets or passes its paired factor.

5. Write a complete sentence explaining how you checked one answer.

Answer Answer should name an inverse, estimate, boundary, known fact or equivalent representation.

Find factor pairs systematically

Prove, compare or generalise. Write enough reasoning for another pupil to follow.

Name: _____ Date: _____

1. Find a number below 100 with exactly 10 factors.

2. Explain why this reasoning fails: Miro stops checking factor pairs after reaching 5.

3. Create a new example that tests the same idea as: How many factors does 36 have?

4. Find a second method or representation. Compare its efficiency with your first method.

5. Write one always, sometimes or never statement about today's learning and justify it.

Teacher answers and checking prompts

1. Find a number below 100 with exactly 10 factors.

Answer 48 is one example.

2. Explain why this reasoning fails: Miro stops checking factor pairs after reaching 5.

Answer Continue until the first factor meets or passes its paired factor.

3. Create a new example that tests the same idea as: How many factors does 36 have?

Answer Answers vary. The example and solution must preserve the mathematical structure.

4. Find a second method or representation. Compare its efficiency with your first method.

Answer Answers vary. Comparison should refer to accuracy, number structure and clarity.

5. Write one always, sometimes or never statement about today's learning and justify it.

Answer Answers vary. A valid example and counterexample should support the classification.