

Solve multi-step addition and subtraction problems

Use the scaffold, show each step and ask for an adult check when marked.

Name: _____ Date: _____

1. Read the model: A venue has 125,000 seats. It sells 48,675 morning tickets and 39,890 evening tickets. How many remain? Copy or complete the first step.

2. A charity target is £500,000. It raises £187,650 then £204,875. How much more is needed?

3. A reservoir holds 2,500,000 litres. It loses 384,750 litres and gains 275,600 litres. New volume?

4. Miro says: Miro subtracts the two ticket sales separately from the original but loses track of the first subtotal. Is this correct? Explain.

Teacher answers and checking prompts

1. Read the model: A venue has 125,000 seats. It sells 48,675 morning tickets and 39,890 evening tickets. How many remain? Copy or complete the first step.

Answer 36,435 seats.

2. A charity target is £500,000. It raises £187,650 then £204,875. How much more is needed?

Answer £107,475.

3. A reservoir holds 2,500,000 litres. It loses 384,750 litres and gains 275,600 litres. New volume?

Answer 2,390,850 litres.

4. Miro says: Miro subtracts the two ticket sales separately from the original but loses track of the first subtotal. Is this correct? Explain.

Answer Combine the tickets first or label each new subtotal before continuing.

Solve multi-step addition and subtraction problems

Work independently. Show a complete method and check at least one answer.

Name: _____ Date: _____

1. A charity target is £500,000. It raises £187,650 then £204,875. How much more is needed?

2. A reservoir holds 2,500,000 litres. It loses 384,750 litres and gains 275,600 litres. New volume?

3. Create a three-step addition and subtraction problem with final answer 250,000.

4. Identify and correct this misconception: Miro subtracts the two ticket sales separately from the original but loses track of the first subtotal.

5. Write a complete sentence explaining how you checked one answer.

Teacher answers and checking prompts

1. A charity target is £500,000. It raises £187,650 then £204,875. How much more is needed?

Answer £107,475.

2. A reservoir holds 2,500,000 litres. It loses 384,750 litres and gains 275,600 litres. New volume?

Answer 2,390,850 litres.

3. Create a three-step addition and subtraction problem with final answer 250,000.

Answer Answers vary. Every step must be labelled and accurate.

4. Identify and correct this misconception: Miro subtracts the two ticket sales separately from the original but loses track of the first subtotal.

Answer Combine the tickets first or label each new subtotal before continuing.

5. Write a complete sentence explaining how you checked one answer.

Answer Answer should name an inverse, estimate, boundary, known fact or equivalent representation.

Solve multi-step addition and subtraction problems

Prove, compare or generalise. Write enough reasoning for another pupil to follow.

Name: _____ Date: _____

1. Create a three-step addition and subtraction problem with final answer 250,000.

2. Explain why this reasoning fails: Miro subtracts the two ticket sales separately from the original but loses track of the first subtotal.

3. Create a new example that tests the same idea as: A reservoir holds 2,500,000 litres. It loses 384,750 litres and gains 275,600 litres. New volume?

4. Find a second method or representation. Compare its efficiency with your first method.

5. Write one always, sometimes or never statement about today's learning and justify it.

Teacher answers and checking prompts

1. Create a three-step addition and subtraction problem with final answer 250,000.
Answer Answers vary. Every step must be labelled and accurate.
2. Explain why this reasoning fails: Miro subtracts the two ticket sales separately from the original but loses track of the first subtotal.
Answer Combine the tickets first or label each new subtotal before continuing.
3. Create a new example that tests the same idea as: A reservoir holds 2,500,000 litres. It loses 384,750 litres and gains 275,600 litres. New volume?
Answer Answers vary. The example and solution must preserve the mathematical structure.
4. Find a second method or representation. Compare its efficiency with your first method.
Answer Answers vary. Comparison should refer to accuracy, number structure and clarity.
5. Write one always, sometimes or never statement about today's learning and justify it.
Answer Answers vary. A valid example and counterexample should support the classification.