

Revisit fraction calculation

Use the scaffold, show each step and ask for an adult check when marked.

Name: _____ Date: _____

1. Read the model: Calculate $\frac{3}{4} + \frac{5}{8}$. Copy or complete the first step.

2. Calculate $2\frac{1}{3} - \frac{5}{6}$.

3. Find $\frac{7}{12}$ of 360.

4. Miro says: Miro adds both numerators and denominators when adding unlike fractions. Is this correct? Explain.

Teacher answers and checking prompts

1. Read the model: Calculate $\frac{3}{4} + \frac{5}{8}$. Copy or complete the first step.

Answer $\frac{11}{8}$ or $1\frac{3}{8}$.

2. Calculate $2\frac{1}{3} - \frac{5}{6}$.

Answer $1\frac{1}{2}$.

3. Find $\frac{7}{12}$ of 360.

Answer 210.

4. Miro says: Miro adds both numerators and denominators when adding unlike fractions. Is this correct? Explain.

Answer Find equivalent fractions with a common denominator, then add the numerators.

Revisit fraction calculation

Work independently. Show a complete method and check at least one answer.

Name: _____ Date: _____

1. Calculate $2\frac{1}{3} - \frac{5}{6}$.

2. Find $\frac{7}{12}$ of 360.

3. Create two different fractions greater than 1 whose sum is 3.

4. Identify and correct this misconception: Miro adds both numerators and denominators when adding unlike fractions.

5. Write a complete sentence explaining how you checked one answer.

Teacher answers and checking prompts

1. Calculate $2\frac{1}{3} - \frac{5}{6}$.

Answer $1\frac{1}{2}$.

2. Find $\frac{7}{12}$ of 360.

Answer 210.

3. Create two different fractions greater than 1 whose sum is 3.

Answer Answers vary, for example $\frac{3}{2} + \frac{3}{2} = 3$.

4. Identify and correct this misconception: Miro adds both numerators and denominators when adding unlike fractions.

Answer Find equivalent fractions with a common denominator, then add the numerators.

5. Write a complete sentence explaining how you checked one answer.

Answer Answer should name an inverse, estimate, boundary, known fact or equivalent representation.

Revisit fraction calculation

Prove, compare or generalise. Write enough reasoning for another pupil to follow.

Name: _____ Date: _____

1. Create two different fractions greater than 1 whose sum is 3.

2. Explain why this reasoning fails: Miro adds both numerators and denominators when adding unlike fractions.

3. Create a new example that tests the same idea as: Find $\frac{7}{12}$ of 360.

4. Find a second method or representation. Compare its efficiency with your first method.

5. Write one always, sometimes or never statement about today's learning and justify it.

Teacher answers and checking prompts

1. Create two different fractions greater than 1 whose sum is 3.

Answer Answers vary, for example $3/2 + 3/2 = 3$.

2. Explain why this reasoning fails: Miro adds both numerators and denominators when adding unlike fractions.

Answer Find equivalent fractions with a common denominator, then add the numerators.

3. Create a new example that tests the same idea as: Find $7/12$ of 360.

Answer Answers vary. The example and solution must preserve the mathematical structure.

4. Find a second method or representation. Compare its efficiency with your first method.

Answer Answers vary. Comparison should refer to accuracy, number structure and clarity.

5. Write one always, sometimes or never statement about today's learning and justify it.

Answer Answers vary. A valid example and counterexample should support the classification.